

Guidelines for the Nurse in the School Setting

4th Edition 2022



EMSC
Emergency Medical
Services for Children

OSF HealthCare
**Children's Hospital
of Illinois**

Table of Contents

Acknowledgements	i
Introduction	ii
Assessment and Triage	1
Assessment is a Lifesaving Tool	1
Components of a Systemic Assessment.....	1
Scene Safety Assessment	3
Across-the-Room Assessment	4
Primary Assessment	5
Secondary Assessment	9
Triage.....	15
Ongoing Monitoring and Reassessment	15
Final Disposition	16
Completing the Process	16
Prevention	17
Students with Special Needs	18
Summary	18
References and Further Information.....	18
School Nurse Protocols	21
Systematic Assessment/Immediate Care	23
Assessment Tools	24
Triage.....	25
Abdominal Pain.....	26
Anaphylaxis/Allergic Reaction	27
Asthma Episode (Acute)/Reactive Airway Disease	28
Bites and Stings.....	29
Burns.....	30
Burn Assessment	31
Chest Pain	32
Chest Trauma	33
Child Maltreatment, Suspected	34
Indications of Child Maltreatment.....	35
Cold Related Injuries (Hypothermia/Frostbite)	36
Dental, Oral and Maxillofacial Emergencies.....	37
Diabetic Emergencies.....	38

Disaster/Emergency Response: Mass Casualty Incidents	39
START/JumpSTART® Algorithms for MCI Triage	40
Drowning/Submersion Injuries	41
Ear Emergencies	42
Eating Disorders	43
Eye Emergencies	44
Foreign Body Airway Obstruction	45
AHA Airway Clearing Maneuvers	46
Headache	47
Head/Spinal Cord Trauma	48
Heat Related Injuries	49
Hemophilia	50
Hemorrhage Control	51
Tourniquet Application	52
Increased Intracranial Pressure in a Student with VP Shunt	54
Lacerations/Abrasion	55
Musculoskeletal Injuries	56
Interventions for Musculoskeletal Injuries	57
Nose Emergencies	58
Obstetric Emergencies	59
Emergency Neonatal Care	60
Respiratory Distress	61
Seizures	62
Sexual Abuse/Sexual Assault and Teen Dating Violence	63
Sickle Cell Disease	64
Substance Abuse	65
Suicide Prevention	66
Syncope/Unconsciousness	67
Throat Emergencies	68
Toxic Emergencies (Ingestion/Environmental)	69
Toxidromes	70
Trauma	71
Interventions for Musculoskeletal Trauma	72
Violent Behavior	73
Violent Behavior: Safety Tips	74

Acknowledgements

Guidelines for the Nurse in the School Setting contains excerpts from the Illinois School Nurse Emergency Care (SNEC) Course manual, 7th Edition (2021), to provide a resource that school nurses can utilize in their practice. This document was developed under the direction of the Illinois EMSC School Nurse Committee. The Illinois Emergency Medical Services for Children Advisory Board acknowledges the commitment and dedication of the EMSC School Nurse Committee members. Their collective efforts have aided the Illinois EMSC program tremendously in striving toward the goal of improving pediatric emergency care within our state.

Special thanks are extended to the Illinois Association of School Nurses, the Illinois Department of Public Health School Health Program, and the Illinois State Board of Education for their ongoing guidance and support.

Guidelines for the Nurse in the School Setting is not copyrighted. Readers are free to duplicate and use all or part of the information in this publication as long as the work is properly cited, the use is educational or for medical care guidance, and is not for profit. In accordance with accepted publishing standards, we request acknowledgment for any information or content reproduced in any other form or publication.

Cite as: *Guidelines for the Nurse in the School Setting, 4th Edition*. OSF Children’s Hospital of Illinois/OSF Saint Francis Medical Center, Peoria, IL. Published: April 2022.

www.osfhealthcare.org/childrens/services/pediatric-readiness

Introduction

The state of Illinois recognizes the unique field of school nursing and the multiple roles that the nurse in the school environment is called upon to perform. The intent of the *Guidelines for the Nurse in the School Setting* is to provide nurses working in the school setting with a set of emergency care guidelines that can be utilized during the delivery of care to the ill or injured student.

It is important to note that there are variations in school settings, services and access to resources and personnel. These differences can impact the delivery of care. Illinois EMSC has made every effort to ensure that the information presented in this document is accurate and represents current accepted practice in the United States. However, the recommendations in this document are not intended to indicate an exclusive course of treatment or to be applicable in all circumstances or to serve as a substitute for the professional advice of a physician. It is recommended that care must always be based upon the student's clinical presentation and on authorized policies. We also recommend that you use the information in this document as a guide for developing local school policies and protocols. It is hoped that this document becomes a valuable addition to the nursing practice resources already available to school nurses.

Because schools provide services to students throughout the childhood and adolescent years, school nurses must be cognizant of pediatric developmental stages and be able to provide age appropriate services and information. A baseline understanding of pediatric developmental characteristics and a general knowledge of appropriate approaches in the assessment and management of the varied age groups will be beneficial. The assessment and treatment approach when dealing with an injured or ill kindergartner is quite different from that of a high-school student, particularly when considering student participation in the decision making process, provision of educational information, and level of parental involvement. In addition, children with special needs or chronic conditions may require more frequent emergency care interventions due to their physical or mental disability and/or chronic illness.

The school nurse has a key role not only in providing emergency care to students, but also in developing prevention strategies. Many emergencies are avoided in the schools each day because school nurses have assisted in the education of teachers/school staff, parents/guardians and students in prevention and early intervention techniques. Proactive development of individualized health care plans and emergency care plans can also play a key role in prevention.

By working collaboratively with school administrators/staff, medical advisors, local EMS agencies, local health care practitioners, and parents/guardians, the school nurse can be instrumental in establishing a comprehensive program of emergency care aimed at reducing student morbidity and mortality.

Keeping abreast of practice changes and trends within the field of school nursing is critical. It is also essential to maintain awareness of state and federal regulations and requirements. Resources that can assist in this effort are available through a variety of sources, particularly the [Illinois Association of](#)

[School Nurses](#), the [Illinois Department of Public Health School Health Program](#), the [Illinois State Board of Education School Health Program](#) and the [National Association of School Nurses](#).

Educational programs leading to certification in the specialty of school nursing can enhance the practice of school nursing and the management of a comprehensive school health program. Certification represents a national standard of preparation, knowledge, and practice. The National Board for Certification of School Nurses (NBCSN) endorses the concept of voluntary certification by examination for all school nurses. Professional certification in school nursing provides an ongoing, quality credentialing process for eligible school nurses. The NBCSN provides the opportunity for school nurses to set the standards for their specialty area through voluntary professional certification.

In addition, Professional Educator License Endorsement in School Nursing (PEL-SN) is available for school nurses in Illinois. There are currently four approved programs in Illinois through which the PEL-SN may be obtained. These are listed below, along with hyperlinks to their websites for additional information.

- DePaul University, Chicago, IL: [School Nurse Endorsement Program](#)
- Illinois State University – Mennonite College of Nursing, Normal, IL: [Graduate Level School Nurse Certification Program](#)
- Lewis University, Romeoville, IL: [School Nurse Non-Degree Program](#)
- University of Illinois - Chicago, Chicago, IL: [School Nurse Certificate Program](#)

There is also a need for ongoing education in order to refresh the school nurse's knowledge and skill base as well as to ensure clinical consistency with current practice guidelines. The Illinois EMSC *School Nurse Emergency Care (SNEC) Course* is a 3 day in-person course that is designed to enhance the assessment and appropriate triaging skills of the school nurse when confronted with the acutely ill or injured child. An on-line version of this course is also available at [Public Health Learning](#). With the increasing number of urgent health related conditions seen within the changing school environment and the need to assure emergency and disaster preparedness, this course supports the school nurse's core knowledge base in responding to these situations. The course is team taught by both school nurses and emergency department nurses since both perspectives are essential in delivering this emergency training. Course content contains lectures, case presentations and skill stations. Continuing education hours are available through an Illinois Department of Public Health site code. Course renewal is recommended every four years. More information about the SNEC course can be accessed on the [OSF Children's Hospital of Illinois](#) website.

The following is a list of additional pediatric focused educational courses that the school nurse may consider to further enhance their pediatric clinical skills:

- [Advanced Pediatric Life Support \(APLS\)](#)
- [Emergency Nursing Pediatric Course \(ENPC\)](#)
- [International Trauma Life Support Pediatric Provider Course \(ITLS Pediatric\)](#)
- [Neonatal Resuscitation Program \(NRP\)](#)
- [Pediatric Advanced Life Support \(PALS\)](#)

- [Pediatric Education for Prehospital Professionals \(PEPP\)](#)
- [Pediatric Emergency Assessment, Recognition and Stabilization Course \(PEARS\)](#)
- [Pediatric Disaster Triage: Utilizing the JumpSTART® Method](#)

We hope that you find the Guidelines for the Nurse in the School Setting useful. Access our website at www.osfhealthcare.org/childrens/services/pediatric-readiness for other pediatric emergency care resources.

Illinois EMSC has made every effort to ensure that the information presented in this document is accurate and represents current accepted practice in the United States. However, the recommendations in this document are not intended to indicate an exclusive course of treatment or to be applicable in all circumstances. We recommend that you use this document as a guide for developing local school policies and protocols.

Assessment and Triage

Assessment Is a Lifesaving Tool

Your importance as a healthcare provider for the students in your school cannot be overstated. On a daily basis, you may treat a wide variety of problems in the student population ranging from mild stomach aches or minor lacerations to life-threatening asthma or traumatic brain injury. During the course of a typical school nurse career, you will alleviate pain and prevent suffering on a daily basis, and you will most likely save lives.

When a student experiences emergent illness or injury, your actions can make the difference between permanent disability and full recovery. It is essential to assess the student's condition swiftly and accurately, without overlooking important physical and historical findings. These findings provide the basis for selecting and prioritizing interventions, evaluating the student's response, and determining disposition. A systematic, consistent approach is the key to this process.

Components of a Systematic Assessment

Four major activities

A systematic assessment can be categorized into four major activities, each of which has a specific role in emergency nursing care.

Scene safety assessment

Ensure that it is safe to approach the injured student. Call for backup assistance if necessary.

Across-the-room assessment

Use the Pediatric Assessment Triangle (PAT, Figure 1) for your across-the-room assessment. **Immediately** activate EMS if the situation is obviously emergent.

Primary assessment

Use the C-ABCDE protocol (Table 1) to identify and treat problems that are life-threatening. The primary assessment includes the following components: control hemorrhage/perform CPR, assess airway, breathing, circulation, disability, and expose the injured area briefly for examination.

Secondary assessment

The secondary assessment always follows the primary assessment. Use the FGHI mnemonic (Table 3) to guide your examination of the student. The secondary assessment includes the following components: measure and record full set of vital signs, give comfort measures/pain assessment, obtain the history and perform a head-to-toe or focused physical exam, and provide interventions as applicable. Inspect, auscultate, percuss, and palpate affected areas to identify or investigate additional problems, and perform ongoing reassessment.

These activities can be adapted to virtually any situation you may face at any time, thereby providing a clear, safe, and consistent basis for clinical decision-making and nursing interventions.

The systematic assessment culminates in an accurate triage decision, appropriate interventions, and optimum final disposition.

NOTE | You will perform these four components almost simultaneously during an actual emergency, in far less time than it takes to describe them.

Key Points of a Successful Assessment

Three of the techniques that foster a successful approach to assessment are briefly discussed below. You may find it helpful to keep them in mind as you visualize how to apply these steps in your own practice.

Use developmentally appropriate language

KEY POINT

Remember to use developmentally appropriate language when addressing students, especially for younger children.

It is important to talk with the student throughout the assessment process; explain your actions and provide reassurance. Be sure to use developmentally appropriate language and techniques as you interact and communicate with the student. You must also attune yourself to the ways a child could interpret your meaning. For example, if you say “*I’m going to take your blood pressure,*” a child could interpret this to mean that you will literally take away something related to their blood. To prevent misunderstandings, say “*I’m going to measure your blood pressure,*” or “*I’m going to give your arm a hug,*” for younger students.

NOTE | Language, culture, technology, and environment may affect the way you implement the assessment process. Enlist special resources as needed to help you communicate effectively with all students.

Gather health history information throughout the process

KEY POINT

If possible, gather focused health history information as you perform each step of the assessment.

As you talk to the student and explain what you are doing, ask questions about the injury or illness you are assessing. Use this technique to gather as much focused health history information as possible while you perform the primary assessment and provide interventions, because this will help you to evaluate your findings more accurately. If the student is unable to respond to your questions, query others who were present when the incident occurred.

Treat significant problems as they are encountered

KEY POINT

It is essential to perform any necessary actions as they are encountered (if able) before moving on to the next step of the assessment.

As you progress through the assessment components, it is essential to perform any necessary actions **before** moving on to the next step. Immediately treat life-threatening problems such as uncontrolled hemorrhage, airway obstruction, or inadequate ventilation. These actions will

ensure optimal outcome. Be aware that although triage is discussed here as the end result of a complete assessment, in practice you will activate EMS at the **earliest** sign of an emergent situation.

Scene Safety Assessment

Hazards

Before providing aid, you must ensure the safety of yourself, the student, and others who are present. Maintain a constant awareness of circumstances that could affect your own safety or that of others, even when inside the healthcare office. To determine whether you can safely approach a student after accident or injury, look for the following hazards:

Substances

These include blood or other body fluids, noxious fumes, and toxic chemicals.

Situational dangers

These include an armed perpetrator, hostages, and weapons.

Environmental dangers

These include unstable structures, fires, electrical hazards, or other potential mechanisms of injury.

KEY POINT

Never place yourself in danger. If you cannot control a hazard, do not approach the student.

Never place yourself in danger. If you cannot control hazards, **do not** approach the student—call 911 or your local emergency number to activate backup assistance.

Resources

If the situation does not appear hazardous, you may find it useful to consider briefly whether you have the equipment and resources you need to manage the incident. For example, you might call for

- Additional personnel to help you with interventions or to manage bystanders
- Personal protective equipment (PPE) or specialized equipment that is not in your portable emergency kit, such as an automated external defibrillator (AED), backboard, cold packs, or splints

NOTE

Always be prepared for unexpected emergencies. Carry disposable gloves and a resuscitation mask at all times, and make sure these items are available in key locations for anyone who might need them.


Across-the-Room Assessment

Quickly Evaluate the Student's Condition

KEY POINT

The across-the-room assessment is a quick overall appraisal of the student's condition based on appearance, breathing, and circulation.

FIGURE 1. PEDIATRIC ASSESSMENT TRIANGLE



Reprinted with permission from Tunik et al. (2006) *Teaching resource for instructors in prehospital pediatrics—Basic life support* (2nd ed.).

Across the room assessment is a natural continuation of the scene safety assessment while also focusing your attention on the student. You will briefly evaluate the student for appearance, breathing, and circulation using the pediatric assessment triangle (PAT) as illustrated in Figure 1. The PAT allows you to rapidly decide how quickly you must proceed with further assessments and interventions.

During the subsequent primary assessment, you will perform a hands-on evaluation of the student's appearance, breathing, and circulation, and undertake any necessary interventions. The most crucial action during the primary assessment is to form an overall impression of the student's level of distress.

Appearance

Appearance refers to mental status, muscle tone, and body position. Mental status is based on the student's level of consciousness and interactions with others. Muscle tone and body position can be assessed by noting if the student is sitting or standing upright appropriately, or able to walk with good movement of the extremities.

Breathing

Breathing refers to the presence or absence of visible movement of the chest or abdomen and work of breathing.

- Can you confirm at a glance if the student is breathing?
- Is there evidence that the student is working hard to maintain adequate ventilation (e.g., flaring nostrils, retractions, or difficulty speaking)?

Circulation

Circulation refers to visible skin color, which is an indication of perfusion to vital organs.

- Does the student's skin color appear normal?
- If not, does the skin color appear pallid, dusky, mottled, cyanotic, or flushed?

Assessment Conclusions

Combine the evidence of your observations with your experience and intuition to form a first impression of the student’s level of distress. Compared with baseline, does this student look well, ill, or seriously ill? Take into account any clearly visible signs and symptoms of illness or injury, such as emesis, bleeding, deformities, or expressions of pain.

Next Steps

If the student’s condition is clearly emergent—for example, the student is struggling to breathe, turning dusky or cyanotic, exhibiting seizure activity, or bleeding profusely—activate EMS **immediately**, then approach the student and proceed with the primary assessment and interventions.

Primary Assessment

C-ABCDE Assessment

The ABCDE assessment focuses on airway, breathing, circulation, disability (neurological status), and a brief physical exposure to examine parts of the body directly related to the chief complaint. The American Heart Association has added a first C step to reprioritize care to begin CPR as indicated before beginning the assessment. In addition, trauma experts have advocated for the first C step to include the early initiation of hemorrhage control measures. To assist the school nurse, the acronym C-ABCDE will consistently be utilized throughout this manual. The school nurse should use the first C step to identify life threatening conditions and then begin CPR and/or initiate hemorrhage control measures, as appropriate. An overview of the C-ABCDE assessment is presented in Table 1.

Consider that many factors in addition to illness or injury can contribute to deviations from the norm as you interpret your assessment findings. It is important to be aware of these factors so that you can take them into account. For example, consider the following:

- Certain medications can cause the breathing rate and heart rate to be fast or slow
- A cold ambient temperature can delay capillary refill and affect other skin findings
- Fear, fever, and pain typically increase the respiratory rate and heart rate
- Students with certain chronic conditions may have baseline vital signs that fall outside the normal range for their age

Children’s vital signs vary by age; therefore, subtle abnormalities can be overlooked. During the primary assessment, you are looking primarily for overt discrepancies that are consistent with emergent and urgent conditions. If none are present, you will have time to subsequently assess less obvious signs during the secondary assessment and focused physical examination.

KEY POINT

The goal of the primary assessment is to identify and treat life-threatening emergencies. Activate EMS as soon as the need becomes evident.

The goal of the primary assessment is to identify and treat life-threatening emergencies. As you progress through the assessment, provide interventions as necessary to maintain the airway, breathing, and circulation **before** continuing to the next step. If at any time the student cannot maintain airway patency, adequate ventilation, or adequate perfusion, or there is evidence of

uncontrolled hemorrhage, **immediately** activate EMS. Notify the parent/guardian as soon as you are able to do so.

TABLE 1. PRIMARY C-ABCDE ASSESSMENT

	Assessment	Interventions
C	Control hemorrhage and perform CPR as indicated	In cases where obvious hemorrhage is occurring or if the student is pulseless, the priority of assessment changes. Control the bleeding before proceeding with primary assessment. If no obvious bleeding is observed, continue the ABCDE assessment. If the student is pulseless, begin compressions and follow the CAB sequence.
A	Airway <i>Observe position, sounds, and possible obstruction.</i>	Restrict the motion of the cervical spine if trauma is suspected or mechanism of injury is unknown. Open airway if necessary; use jaw-thrust or chin-lift procedures as appropriate. Position student to maintain airway patency. Look and listen for signs of obstruction. Perform airway-clearing maneuvers as indicated.
B	Breathing <i>Observe rate, depth, pattern, symmetry, sounds, work of breathing, odors, and possible injuries.</i>	If student is breathing spontaneously, allow the student to remain in a position of comfort that promotes the best ventilation. Give oxygen if available and if tolerated by the student. Provide mouth-to-mask ventilation if needed.
C	Circulation <i>Observe heart rate, pulse quality, capillary refill time, skin color, temperature, moisture, and possible bleeding.</i>	Continue CPR as needed. Control uncontrolled external bleeding with direct pressure, pressure dressings or tourniquet. If bleeding occurs in an extremity, elevate the affected limb. If uncontrolled external bleeding is noted at any point during the assessment, controlling the bleeding takes priority over other interventions.
D	Disability (neurological status) <i>Determine level of consciousness using the alert, verbal, painful, unresponsive (AVPU) scale; observe pupillary response.</i>	Provide reassurance, position to maintain comfort, and give oxygen if available.
E	Exposure with environmental control to prevent heat loss <i>Observe and inspect for additional emergent problems.</i>	Treat wounds or other findings as appropriate. Treat as necessary to maintain normothermia. Cover student with blankets after initial exposure. Consider the need for isolation if a rash is noted with potential for communicable disease.

NOTE Always observe standard precautions to prevent exposure to body fluids during a physical examination.

Spinal Motion Restriction

If there is any possibility of injury to the head or spine, provide manual motion restriction of the cervical spine before you begin the airway assessment. Maintain spinal motion restriction until EMS personnel arrive.

Airway

Assess the student's airway for patency by considering the questions below. Remember to restrict the motion of the neck if the student suffered a traumatic injury or if the mechanism of injury is unknown.

- Can the student vocalize?
- Are there secretions or material obstructing the airway?
- Is edema present?

Open the student's mouth and inspect for sources of airway obstruction. Manually open the airway using a head-tilt chin-lift or jaw-thrust if needed. The jaw-thrust is the preferred technique to open the airway if trauma is suspected. Maintain cervical spine motion restriction for any student with suspected trauma or unknown mechanism of injury.

Interpreting your findings

Consider the following possible causes as you interpret your findings:

- Drooling or inability to talk may indicate upper airway edema from infection or anaphylaxis
- Stridor (a crowing sound on inspiration) indicates upper airway obstruction
- Facial injuries can compromise the airway

Breathing

- Assess the respiratory rate (e.g., normal, fast, slow, or apnea), the depth and pattern of breathing (e.g., shallow, gasping, or irregular), and the bilateral symmetry of chest movement
- Listen for adventitious breath sounds such as wheezing or grunting
- Watch for signs that indicate increased work of breathing (e.g., retractions, nasal flaring, or difficulty speaking)
- Assess any unusual breath odors (e.g., fruity, or reminiscent of petroleum products, tobacco, or alcohol)
- Evaluate chest wall integrity, and note any obvious injuries

Interpreting your findings

- Increased work of breathing and audible breath sounds indicate respiratory compromise
- Fast breathing is an early compensatory mechanism for hypoxia; breathing slows as hypoxia worsens
- Decreased, absent, or unequal breath sounds may indicate airway obstruction, a pneumothorax, hemothorax, or atelectasis
- Grunting is a late sign of severe respiratory distress that is worsening into respiratory failure


Circulation

During the circulation assessment, you will

- Assess heart rate and quality of pulses
- Evaluate perfusion by assessing
 - Capillary refill time (normally less than 2 seconds)

ASSESSMENT AND TRIAGE

- Skin color (e.g., normal, pallid, dusky, mottled, cyanotic, or flushed)
- Skin temperature and moisture (e.g., warm, dry, cool, or clammy)
- Look for active bleeding (e.g., none, minor, moderate, profuse, controlled, or uncontrolled)



Heart rate and quality of pulses

Measure heart rate by palpating the central and peripheral pulses. Evaluate the pulse quality and note discrepancies between the central and peripheral pulses. The pulse points are shown in Figure 2. You will not measure the student's blood pressure until the secondary assessment.

Perfusion

Capillary refill time

Assess capillary refill at distal extremities with the limb positioned so that it is level with the heart. Firmly press and release the skin to blanch the underlying capillary bed. Color should return in less than 2 seconds.

Color

Inspect the skin color at the lips and tongue. Note whether the skin color seems normal, pale, flushed, mottled, or cyanotic.

NOTE

To assess skin color in students with dark skin pigmentation, check the nail beds, palms, or mucous membranes.

Temperature

Feel skin temperature at the extremities and compare with skin temperature at the central body regions. If skin is either unusually hot or cold, this is a significant finding.

Moisture and quality

Assess the skin for signs of unusual dryness or clamminess. Pinch the skin gently to check for normal elasticity.

Bleeding

Look for profuse external bleeding. If found, control hemorrhage before resuming assessment.

Interpreting your findings

- Tachycardia is usually the earliest sign of developing shock in children; tachycardia combined with a fast respiratory rate is a strong indicator of compensated shock
- Thready or weak pulses, cool or clammy extremities, and delayed capillary refill time are associated with shock

- A discrepancy between the quality of central and peripheral pulses may be an early sign of decreasing stroke volume
- Hypotension with bradycardia is a late, ominous sign of decompensated shock
- Skin that is inelastic and prone to tenting usually signifies dehydration

Disability

The disability assessment is a brief evaluation of neurological function. First assess the level of consciousness using the alert, verbal, painful, and unresponsive (AVPU) scale (Table 2), and then evaluate pupil size and reactivity.

TABLE 2. AVPU SCALE

	Mnemonic / Topic	Associated Findings
A	Alert	The student is awake and able to speak or interact spontaneously.
V	Verbal	A verbal stimulus elicits some response; for example, the student's eyes may open when you call loudly, or agitation may lessen in response to a command.
P	Painful	The student responds to a painful stimulus by moaning, crying, or withdrawing from pain.
U	Unresponsive	The student shows no response to verbal or painful stimuli.

Interpreting your findings

The level of consciousness is an important indicator of adequate perfusion. A significant decline in responsiveness is an ominous sign in a student with an illness or injury that may cause respiratory compromise or shock.

Exposure

Remove clothing as needed to briefly assess specific factors related to the presenting problem, such as injuries, bleeding, rashes, bites, or stings. Watch for signs of internal hemorrhage i.e. tachycardia. Control ambient temperature if possible or drape the student with coverings as necessary to prevent heat loss. Replace clothing as soon as you are able to do so. Consider need for isolation if communicable disease is suspected.

Secondary Assessment

After completing the primary assessment and addressing all immediate life-threats, you will then move directly to the secondary assessment. The secondary assessment is guided by the FGHI mnemonic, which is presented in Table 3.

Full Set of Vital Signs

Your goal during the secondary assessment is simply to identify any clearly abnormal vital signs that suggest an emergent condition. Establishing baseline vital signs during the secondary assessment is essential for the triage determination (discussed in the following section) and

ongoing reassessment. Baseline vital signs are important elements for the continuity of care if the student is referred for further medical evaluation and treatment.

TABLE 3. SECONDARY ASSESSMENT: FGHI MNEMONIC

	Mnemonic / Assessment	Comments
F	Full set of vital signs	Measure and document baseline vital signs, including blood pressure, heart rate, respiratory rate, and temperature. If possible, measure and document weight and blood glucose level.
G	Give comfort measures/pain assessment	Complete a pain assessment. Perform interventions to alleviate pain, such as applying a cold pack, immobilizing a suspected fracture, or dressing a wound.
H	History Head-to-toe assessment/focused physical examination <i>Inspect, auscultate, palpate, percussion (can be partial or complete)</i>	Obtain history. Perform a head-to-toe examination or specialized focused assessment procedures as indicated, such as abdominal assessment/palpation, or range-of-motion assessments.
I	Injuries Inspect the back Additional Interventions	Assess injuries. If you find injuries at varying stages of healing, consider the possibility of child maltreatment. Inspect the student's posterior surfaces, as these are often overlooked areas. Provide additional interventions according to your findings.

Normal vital signs vary with the student's age (Table 4). Other factors that can affect normal vital signs include chronic medical conditions, physical disabilities, medications, environmental conditions, and the student's emotional state.

TABLE 4. PEDIATRIC VITAL SIGNS BY AGE

Age	RR	HR	BP
Neonate (birth–30 days)	30–60	100–180	60–90
Infant (1–12 months)	24–50	100–160	70–100
Toddler (1–3 years)	24–40	90–150	70–105
Preschooler (3–5 years)	20–30	80–140	75–105
School-aged (5–12 years)	18–30	65–120	80–120
Adolescent (12 years and older)	12–20	60–100	90–128

RR, respiratory rate; HR, heart rate; BP, systolic blood pressure (mm Hg).

Measuring blood pressure

To accurately measure blood pressure (BP), use a cuff that is two-thirds as wide as the student's upper arm from the elbow to the axilla. If you do not have a table of normal blood pressure rates, you can use the following formula to approximate the **lowest** acceptable limit for systolic blood pressure for children older than 1 year:

BP should be $>70 + (2 \times \text{age in years})$

NOTE

Do not rely solely on blood pressure to indicate the severity of the student's condition. Children can maintain normal blood pressure until decompensated shock is imminent.

Temperature and other measurements

Measure and record the student's temperature. If possible, also measure and record the student's weight and blood glucose levels.

Temperature conversion

Use the following formulas or Table 5 to convert temperatures between Fahrenheit and Centigrade:

To convert Centigrade to Fahrenheit: $(1.8 \times ^\circ\text{C}) + 32 = ^\circ\text{F}$

To convert Fahrenheit to Centigrade: $(^\circ\text{F} - 32) \times 0.556 = ^\circ\text{C}$

TABLE 5. TEMPERATURE EQUIVALENTS

$^\circ\text{C}$	$^\circ\text{F}$	$^\circ\text{C}$	$^\circ\text{F}$
34.2	93.6	38.6	101.5
34.6	94.3	39.0	102.2
35.0	95.0	39.4	102.9
35.4	95.7	39.8	103.6
35.8	96.4	40.2	104.4
36.2	97.2	40.6	105.1
36.6	97.9	41.0	105.8
37.0	98.6	41.4	106.5
37.4	99.3	41.8	107.2
37.8	100.0	42.2	108.0
38.2	100.8	42.6	108.7

C= Centigrade; F= Fahrenheit.

Give Comfort Measures/ Pain Assessment

Pain has been referred to as the fifth vital sign because of its significance in assessing pathophysiology. The PQRST assessment tool (Table 6) outlines a detailed, systematic interview that can be used to identify clinically significant aspects of pain.

Severity may be the most challenging aspect of pain to assess because it is largely a subjective determination. Various assessment tools can help students quantify the severity of their pain; it is important to select a tool that is suitable for the student's age, cognitive development, and cultural background. For example, school-aged children with average developmental abilities can

often rate their pain using a simple numerical scale such as “On a scale of 0 to 10, if 0 is no pain at all and 10 is the worst pain you’ve ever felt, tell me how bad the pain is right now.” For younger students (aged approximately 3 years and older), those who are less comfortable with numbers, and those who lack the cognitive skills to describe pain using a number scale, the Wong-Baker FACES Pain Rating Scale may be used (Figure 3). The FACES scale allows a student to point to a cartoon face with an expression that reflects the student’s pain experience. For infants and those with developmental impairment or other impediments to verbal communication, pain may be assessed by observing behavioral cues (e.g., facial grimace) and physiologic parameters (e.g., heart rate).

Comfort measures should be provided for students in pain. These measures can include applying a cold pack, stabilizing suspected fractures, and dressing wounds.

TABLE 6. PQRST PAIN ASSESSMENT

Assessment Points	
P	<ul style="list-style-type: none"> ▪ Problem: How does the student describe the chief complaint? ▪ Provoke: What makes the pain worse? ▪ Palliate: What makes the pain better?
Q	<ul style="list-style-type: none"> ▪ Quality: What is the quality or character of the pain?
R	<ul style="list-style-type: none"> ▪ Radiate: Does the pain or discomfort seem to travel or move?
S	<ul style="list-style-type: none"> ▪ Severity: Using a developmentally appropriate assessment tool (e.g., numerical or FACES^a scale), how does the student rate the severity of pain or discomfort? ▪ Signs: What clinical signs accompany the problem? ▪ Symptoms: What subjective problems does the student report?
T	<ul style="list-style-type: none"> ▪ Timing: When did the pain start? Was the onset sudden or gradual?

^aFACES is the Wong-Baker FACES Pain Rating Scale.

FIGURE 3. WONG-BAKER FACES PAIN RATING SCALE



History and Head-to-Toe/Focused Physical Examination

History

Information gathered from the health history helps you form a care plan. You may already be familiar with the student's history from information supplied by the parent/guardian, primary healthcare provider, and school staff, or from previous interactions with the student. Ensure that you understand the student's current health issues, and obtain specific details relevant to the injury or illness.

NOTE For younger students, obtain the health history from the parent/guardian or from the student's health record.

A useful mnemonic that will help you work through the health history assessment is SAMPLE, a health history tool (Table 7), which can be seen on the following page.



TABLE 7. SAMPLE HISTORY

Mnemonic / Topic		Questions
S	Symptoms <i>Ask the student to describe current symptoms, particularly pain.</i>	<ul style="list-style-type: none"> ▪ What problem brings the student to the health office? ▪ How long has the problem persisted? ▪ If an injury, how and when did it occur? ▪ Does the student have pain, apprehension, or guarding? ▪ What are the location, quality, and duration of pain? ▪ Does repositioning make the pain better or worse? ▪ What strategies make other symptoms better or worse? ▪ What is the student's impression of his or her condition?
A	Allergies	<ul style="list-style-type: none"> ▪ Does the student have any known allergies to food, medications, latex, other materials, or environmental elements such as bee stings?
M	Medications <i>List medications the student takes regularly, including dosage regimen and time of the last dose.</i>	<ul style="list-style-type: none"> ▪ Is the student using any prescription, over-the-counter, home, herbal, or cultural remedies? For what reasons? When was the last dose taken? ▪ Did the student take any medications before coming to the health office? What was the result? ▪ Has the student used any illicit drugs? When?
P	Past health history <i>Note preexisting physical or psychological disabilities, previous trauma, and chronic conditions. Check immunization status, including tetanus prophylaxis.</i>	<ul style="list-style-type: none"> ▪ Does the student have a chronic illness (e.g., asthma, diabetes, hemophilia, or seizure disorder)? ▪ Does the student have special healthcare needs?^a ▪ Does the student rely on a medical device (e.g., oxygen, tracheostomy, nebulizer, central venous line, or gastrostomy tube)? ▪ Are immunizations up to date? ▪ Is isolation necessary (e.g., varicella exposure, or immunosuppression)? ▪ If a student is of child-bearing age or has experienced menarche, when was her last menstrual period? Is she pregnant, or is there a possibility of pregnancy?
L	Last meal <i>Document when and what the student last ate or drank.</i>	<ul style="list-style-type: none"> ▪ Is the student able to eat? ▪ When was the last meal? ▪ Has there been any nausea, vomiting, or diarrhea? ▪ Is bowel and bladder function normal?
E	Events <i>Ask the student to describe the events that led up to the illness or injury.</i>	<ul style="list-style-type: none"> ▪ When did the problem begin? ▪ Were there precipitating factors? ▪ If an injury occurred, were there witnesses? What did they report?

^aYou may need to modify your evaluation of assessment findings for students with special needs, because their baseline findings may differ from accepted averages.

Head-to-toe/focused physical examination

The physical examination may be partial (focusing on the site of illness or injury) or complete (head-to-toe). The choice depends on the circumstances, the student's health status, and applicable protocols.

Isolate, Injuries, and Additional Interventions

Assess the potential for communicable disease and isolate as indicated. Provide appropriate and additional interventions according to your findings.

Triage

Principles of Triage

School nursing practice is unpredictable. There may be times when you have several students and staff or visitors waiting in the health office, some reporting for scheduled interventions, others presenting with unexpected problems. You also must respond to episodes of acute illness or injury arising elsewhere around the facility. Consequently, you must be able to determine quickly who needs immediate intervention and who can wait. You will not always treat your students and other healthcare consumers in the order of arrival. Instead, you must treat them according to the seriousness of the presenting illness or injury, and place those with critical conditions ahead of others. Triage (derived from the French word meaning *to sort*) provides an objective way to do this.

Based on your primary assessment findings, you can classify students into emergent, urgent, or nonurgent categories according to whether the condition has the potential to threaten life, limb, or vision. Specific treatment priorities and dispositions are associated with each triage category. If the student's condition changes during ongoing monitoring and reassessment, the triage category may be changed as appropriate. It is better to overtriage than to undertriage, particularly if your intuition tells you that there could be a significant underlying problem. Experience also can help you decide if your triage determination should be upgraded. For example, if a student who has a history of severe asthma presents with mild dyspnea, you might immediately consider this student's condition emergent and activate EMS transport based on previous events during which the student rapidly developed respiratory distress.

KEY POINT

Do not delegate tasks related to the primary assessment and triage. Only an experienced registered nurse has the expertise to accurately categorize ill and injured students.

Although you may delegate tasks associated with ongoing healthcare, do not delegate tasks related to the primary assessment and triage. Only an experienced registered nurse has the requisite knowledge and expertise to quickly recognize and accurately categorize ill and injured students.

Performing Triage

On the following page, Table 8 describes the three commonly recognized triage categories (emergent, urgent, and nonurgent) and presents examples of illnesses and injuries that fall within each category.

TABLE 8. TRIAGE CATEGORIES, EXAMPLES, AND DISPOSITION

Category	Examples	Disposition
Emergent <i>Student requires immediate medical attention. Condition is acute and potentially threatens life, limb, or vision.</i>	<ul style="list-style-type: none"> ▪ Cardiopulmonary arrest ▪ Altered mental status ▪ Burns, severe or major ▪ Caustic chemical exposure ▪ Childbirth, imminent ▪ Head injury, history of loss of consciousness ▪ Pain, severe or significant location (e.g., chest) ▪ Poisoning or drug overdose ▪ Respiratory distress (severe) or respiratory failure ▪ Shock, any type ▪ Spinal cord injury, suspected ▪ Status epilepticus or first-time seizure ▪ Threatens harm to self or others ▪ Trauma, severe; trauma to limb, no distal pulse 	<ul style="list-style-type: none"> ▪ Monitor^a in health office ▪ Transport to emergency care facility via EMS
Urgent <i>Student requires additional medical intervention within 2 hours. Condition is acute but not severe or life-threatening.</i>	<ul style="list-style-type: none"> ▪ Burns, minor ▪ Deformity or suspected closed fracture without circulatory compromise ▪ Febrile illness (T exceeds 100°F/37.8°C) ▪ GI symptoms (e.g., nausea, vomiting, or diarrhea) ▪ Lacerations requiring sutures without excessive blood loss ▪ Pain, moderate, after abdominal trauma ▪ Seizure, atypical, in student with history of seizures ▪ Wheezing 	<ul style="list-style-type: none"> ▪ Monitor in health office ▪ Transport to emergency care facility via EMS, parent/guardian, or other adult as appropriate
Nonurgent <i>Student may require referral for routine medical care. Minor or nonacute condition.</i>	<ul style="list-style-type: none"> ▪ Essentially well; S/S of mild noncommunicable illness or URI ▪ Headache without fever or abnormal findings ▪ Injury, minor (e.g., abrasions, ecchymoses, muscle sprains, or muscle strains) ▪ Pain, mild (e.g., abdominal or menstrual pain, headache or toothache) 	<ul style="list-style-type: none"> ▪ Monitor in health office ▪ Contact parent/guardian ▪ Return student to class or send home as indicated

^a Monitoring should be performed by the school nurse. EMS, emergency medical services; F, Fahrenheit; C, centigrade; GI, gastrointestinal; S/S, signs/symptoms; T, temperature; URI, upper respiratory infection.

Ongoing Monitoring and Reassessment

Ongoing monitoring and reassessment are an essential part of nursing care. Your initial triage decision allows you to formulate a nursing diagnosis and develop the care plan. However, you may need to adjust the triage category and disposition after evaluating the student's response to interventions and determining whether health status has improved or worsened. This may determine the mode of transport, whether via EMS or by car with the parent/guardian. A student whose condition initially appeared nonurgent may abruptly deteriorate, which requires immediate attention. Therefore, it is crucial that you conduct frequent reevaluations and arrange for ongoing monitoring and observation while you are busy elsewhere. Follow applicable protocols regarding appropriate monitoring using assessment tools as outlined in this document.

Final Disposition

Once your assessment is complete, you will form a triage decision and come to a conclusion about the student's disposition. Disposition typically includes the following options:

- Student returns to class
- Observe the student in the school health office for a period of time
- Contact the parent/guardian to transport the student home, to their primary care provider's office, or to an emergency care facility
- Contact EMS to transport the student to an emergency care facility

Completing the Process

Three Additional Actions

Once the emergency is over, the following three activities must be performed:

- Document the incident and collect data
- Evaluate the incident and follow-up the student's disposition and recovery
- Plan for future incidents and prevent reoccurring incidents

This document provides an overview of the steps needed to complete each of these components, including assessment, triage, and postincident activities. Additional details and examples of how these tools are used in specific situations are the focus of subsequent chapters. Your own experience and common sense integrate the steps into a congruent whole.

Documentation and Data Collection

Document your assessment findings and triage category in the student's health record. List your interventions, the student's response, and record the final disposition. Transfer all relevant information to prehospital care providers such as permission to treat the student, allergies, medications, immunizations, and contact information. This information will be included in their report to emergency department personnel. Collect and analyze data to identify injury and illness patterns so that you can monitor and prevent recurrences.

Follow-Up

If possible, follow up any student healthcare incident with the student's primary healthcare provider, the emergency department physician, the prehospital care professionals, and the parent/guardian. Document the outcome in the student's health record and incident report as appropriate.

Prevention

Prevention should be considered the final component in any healthcare visit. You have an instrumental role in initiating and maintaining measures to minimize the effects of injury or illness and reduce the risk of inaccurate triage decisions. This includes the following actions:

- Establish written triage guidelines or protocols
- Designate experienced registered nurses to perform triage

ASSESSMENT AND TRIAGE

- Maintain knowledge and practice requirements related to triage and decision-making skills
- Know which students have emergency action plans/emergency care plans on file and review the necessary interventions with others who may be responsible for carrying them out
- Develop and maintain individualized healthcare plans and emergency action plans/emergency care plans for students with special needs
- Ensure that crucial information from these guidelines, protocols, and care plans is shared with teachers, aides, coaches, and lunchroom or playground monitors as appropriate

NOTE

Under the Family Educational Rights and Privacy Act, parent/guardian consent is not required before sharing student health records with school personnel **provided** they have a legitimate educational interest in the student or a need to know the information to protect the student's welfare.

Students with Special Needs

Although the pediatric assessment techniques outlined in this chapter are equally applicable to students who have special healthcare needs, you must adjust the techniques to accommodate the student's developmental age rather than their chronological age. Be aware that day-to-day baseline vital signs for a student with a chronic condition may be outside the range of same-age peers who do not have special needs.

As a school nurse, you observe the day-to-day variations among your school's students, and are often the first to notice changes in a student's condition. It is particularly important to become familiar with the baseline status of a special needs student so that you can identify when the student's condition has changed.

Use developmentally appropriate language, gestures, and techniques when communicating with a student who has cognitive impairment or developmental delay. When assessing students who depend on assistive technology or specialized devices, do not allow yourself to be distracted by the equipment. Your focus should always be directed toward the student.

Summary

One of your primary responsibilities as school nurse is to recognize severe illness or injury among the students in your school. It is best to do this using a systematic, consistent assessment schema. Your primary assessment findings give you a basis for determining the urgency of each student's condition through triage. Triage categories allow you to treat students according to their acuity of need. Written triage protocols and policies are important to facilitate this process.

References and Further Information

- American Academy of Pediatrics. (2009.) Emergency preparedness for children with special healthcare needs (updated 2010). <https://pediatrics.aapublications.org/content/125/4/829>
- American Academy of Pediatrics. (2009). Health, mental health and safety guidelines for schools. <http://www.nationalguidelines.org>
- American Academy of Pediatrics. (2021). Children and youth with special needs. <https://www.aap.org/en-us/advocacy-and-policy/aap-health-initiatives/Children-and-Disasters/Pages/CYWSN.aspx>

- American Academy of Pediatrics. (2020). *Pediatric education for prehospital professionals* (4th ed.). Jones & Bartlett Learning.
- Emergency Nurses Association. (2017). *Emergency nursing core curriculum* (7th ed.). Elsevier.
- Emergency Nurses Association. (2009). *Core curriculum for pediatric emergency nursing* (2nd ed.). Jones & Bartlett Learning.
- Emergency Nurses Association. (2020). *Emergency nursing pediatric course provider manual* (5th ed.). Jones & Bartlett Learning.
- Hockenberry, M.J., Wilson, D., Rodgers C. (2017). *Wong's essentials of pediatric nursing* (10th ed.). Mosby.
- National Association of Emergency Medical Technicians (NAEMT). (2016) .*PHTLS: Basic and advanced prehospital trauma life support* (8th ed.). Jones & Bartlett Learning.
- Romig, L.E. (2001). PREP for peds—patient physiology, rescuer responses, equipment, protocols. Size-up & approach tips for pediatric calls. *Journal of Emergency Medical Services*, 26(5), 24–33.
- Tunik, M., Treiber, M., Kim, J., Cooper, A., Foltin, G. (Eds.). (2009). *Teaching resource for instructors in prehospital pediatrics (TRIPP)—Basic life support* (2nd ed.). New York University Langone Medical Center, Department of Pediatrics, Center for Pediatric Emergency Medicine. DOI: [10.1080/10903120802474539](https://doi.org/10.1080/10903120802474539)

School Nurse Protocols

This section contains protocols that have been developed by Illinois EMSC to guide school nurses as they assess and manage students with emergency health issues in the school setting. Please note that Illinois EMSC has made every effort to ensure that the information presented in this document is accurate and represents current accepted practice in the United States. However, the recommendations in this document are not intended to indicate an exclusive course of treatment or to be applicable in all circumstances. We recommend that you use this document as a guide for developing local school policies and protocols.

<u>Systematic Assessment/Immediate Care</u>	23
<u>Assessment Tools</u>	24
<u>Triage</u>	25
<u>Abdominal Pain</u>	26
<u>Anaphylaxis/Allergic Reaction</u>	27
<u>Asthma Episode (Acute)/Reactive Airway Disease</u>	28
<u>Bites and Stings</u>	29
<u>Burns</u>	30
<u>Burn Assessment</u>	31
<u>Chest Pain</u>	32
<u>Chest Trauma</u>	33
<u>Child Maltreatment, Suspected</u>	34
<u>Indications of Child Maltreatment</u>	35
<u>Cold-Related Injuries (Hypothermia/Frostbite)</u>	36
<u>Dental, Oral, and Maxillofacial Emergencies</u>	37
<u>Diabetic Emergencies</u>	38
<u>Disaster/Emergency Response: Mass-Casualty Incidents</u>	39
<u>START/JumpSTART® Algorithms for MCI Triage</u>	40
<u>Drowning/Submersion Injuries</u>	41
<u>Ear Emergencies</u>	42
<u>Eating Disorders</u>	43
<u>Eye Emergencies</u>	44
<u>Foreign Body Airway Obstruction</u>	45
<u>AHA Airway Clearing Maneuvers</u>	46
<u>Headache</u>	47
<u>Head/Spinal Cord Trauma</u>	48
<u>Heat-Related Injuries</u>	49

Hemophilia	50
Hemorrhage Control	51
Tourniquet Application	52
Increased Intracranial Pressure in a Student with VP Shunt	54
Lacerations/Abrasions	55
Musculoskeletal Injuries	56
Interventions for Musculoskeletal Injuries	57
Nose Emergencies	58
Obstetric Emergencies	59
Emergency Neonatal Care	60
Respiratory Distress	61
Seizures	62
Sexual Abuse/Assault and Teen Dating Violence	63
Sickle Cell Disease	64
Substance Abuse	65
Suicide Prevention	66
Syncope/Unconsciousness	67
Throat Emergencies	68
Toxic Exposure (Ingestion/Environmental)	69
Toxidromes	70
Trauma	71
Interventions for Musculoskeletal Trauma	72
Violent Behavior	73
Violent Behavior: Safety Tips	74

Systematic Assessment/Immediate Care

Every nursing encounter begins with a systematic assessment. The **four components** of a comprehensive systematic assessment are listed below. Suggested actions should be performed if indicated and in accordance with applicable protocols and available resources. **Provide any necessary interventions before progressing to the next step of the assessment.** These assessment components will be used for essentially every nursing encounter, and it is important to become completely familiar with each component. It will be useful to keep this information in an easily accessible location for continual reference.

NOTE

Triage determination and EMS activation should occur as soon as the need becomes apparent.

SCENE SAFETY ASSESSMENT

Call for assistance as indicated before proceeding.

ACROSS-THE-ROOM ASSESSMENT

Use the Pediatric Assessment Triangle (PAT)

- **Appearance:** Mental status; muscle tone; and body position
- **Breathing:** Work of breathing (nasal flaring and retractions); abnormal airway sounds
- **Circulation:** Visible skin color

PRIMARY (C-ABCDE) ASSESSMENT/IMMEDIATE INTERVENTIONS

- Standard precautions
- Restrict motion of cervical spine
- Control obvious hemorrhage as applicable ▪ CPR as applicable
- Airway, positioning
- Breathing, O₂, mouth-to-mask
- Circulation, control bleeding ▪ CPR/AED
- Disability, AVPU, pupil check
- Exposure, brief inspection

SECONDARY (FGHI) ASSESSMENT

- Full set of vital signs, temperature, weight, and blood glucose
- Give comfort measures/pain assessment
 - PQRST pain assessment ▫ Numerical pain scale ▫ FACES visual pain scale
- History and Head-to-toe/focused physical examination
 - SAMPLE history: Symptoms, Allergies, Medications, Past health history, Last food/drink, Events
 - Complete or limited physical examination: Inspect, Auscultate, Percuss, Palpate
- Injuries, Inspect the back, and Additional Interventions
 - Assess injuries; Inspect posterior surfaces; perform additional interventions based on findings

TRIAGE (E-U-N)

- **E**mergent: Activate EMS
- **U**rgent: Determine need for EMS
- **N**onurgent: Return to class or send home

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Assessment Tools

These assessment tools will be used for essentially all protocols and nursing encounters. It will be useful to become completely familiar with them and always have them readily available for reference.

SYSTEMATIC ASSESSMENT

NOTE: Perform interventions AS YOU GO. Determine triage/activate EMS at EARLIEST INDICATION of need.

Scene safety assessment

- Call for assistance as indicated

Across-the-room assessment

Use Pediatric Assessment Triangle (PAT)

- Appearance
- Breathing
- Circulation

Primary C-ABCDE assessment

- Standard precautions
- Spinal motion restriction
- Control hemorrhage/CPR (as applicable)
- Airway
- Breathing
- Circulation
- Disability^a
- Exposure

Secondary FGHI assessment

- Full set of vital signs, temperature, weight, and blood glucose
- Give comfort measures/pain assessment (PQRST, FACES)
- History (SAMPLE) and head-to-toe/focused physical examination (inspect, auscultate, percuss, palpate)
- Injuries, inspect back, and provide other interventions

Triage

- Emergent
- Urgent
- Nonurgent

^aDisability assessment

- Assess responsiveness (AVPU):
 - A Alert
 - V Responds to Verbal stimulus
 - P Responds to Painful stimulus
 - U Unresponsive
- Assess pupils
- Assess for transient paresthesia

PEDIATRIC VITAL SIGNS BY AGE

Age	RR	HR	BP
Neonate (birth–30 days)	30–60	100–180	60–90
Infant (1–12 months)	24–50	100–160	70–100
Toddler (1–3 years)	24–40	90–150	70–105
Preschooler (3–5 years)	20–30	80–140	75–105
School-aged (5–12 years)	18–30	65–120	80–120
Adolescent (12 years and older)	12–20	60–100	90–128

RR, respiratory rate; HR, heart rate; BP, systolic blood pressure (mm Hg)

INDICATORS OF CARDIOPULMONARY COMPROMISE IN CHILDREN

- Tachycardia
- Weak, thready, or absent peripheral pulses
- Decreasing consciousness; altered mental status
- Tachypnea/respiratory difficulty
- Central cyanosis and coolness
- Hypotension (late sign)
- Bradycardia (ominous sign)
- No palpable BP (ominous sign)

GLASGOW COMA SCALE (GCS) PARAMETERS

	<2 Years Old (Pediatric GCS)	>2 Years Old (GCS)
Eye opening (E)	(4) Spontaneous	(4) Spontaneous
	(3) To speech	(3) To speech
	(2) To pain	(2) To pain
	(1) None	(1) None
Verbal response (V)	(5) Coos, babbles, appropriate words	(5) Oriented, appropriate words
	(4) Irritable, cries but consolable	(4) Confused
	(3) Cries to pain, inconsolable	(3) Inappropriate words, persistent cry
	(2) Moans to pain	(2) Incomprehensible sounds
	(1) None	(1) None
Motor response (M)	(6) Normal spontaneous movements	(6) Obeys commands
	(5) Withdraws from touch	(5) Localizes to pain
	(4) Withdraws from pain	(4) Withdraws from pain
	(3) Abnormal flexion (decorticate)	(3) Abnormal flexion (decorticate)
	(2) Abnormal extension (decerebrate)	(2) Abnormal extension (decerebrate)
	(1) None	(1) None
Total Glasgow Coma Scale Score = 3–15		

Triage

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

The assessment findings allow you to determine a triage category, which dictates subsequent actions. Key findings associated with each triage category and general interventions are listed below. Specific actions will vary according to the situation. This information should be kept in a readily accessible location because you will use it frequently for many nursing encounters.

EMERGENT

Findings that suggest a potential threat to life or function require immediate medical attention. These include

- Cardiopulmonary arrest
- Altered mental status
- Anaphylaxis
- Burns, severe/major
- Caustic chemical exposure
- Childbirth, imminent
- Head injury with any history of loss of consciousness
- Pain, severe **or** significant location (e.g., chest)
- Poisoning/drug overdose
- Respiratory distress, severe, **or** respiratory failure
- Shock (hypovolemic, anaphylactic, or cardiogenic)
- Spinal cord injury, suspected
- Status epilepticus **or** first-time seizure
- Suicidal behavior
- Trauma, severe, **or** trauma to limb, distal pulse absent
- Uncontrolled hemorrhage
- Violent/homicidal behavior

INTERVENTIONS

- Control hemorrhage
- Activate EMS
- Support C-ABCDE
- Initiate appropriate interventions as per specific protocol or IHP and EAP/ECP
- Directly/continuously observe student
- Contact parent/guardian
- Notify school administrator
- Follow-up

URGENT

Findings that suggest an acute, severe, but **non-life-threatening** condition require additional medical intervention within 2 hours. These include

- Burns, minor
- Deformity/suspected closed fracture **without** circulatory compromise
- Fever exceeding 100°F/37.8°C
- GI symptoms, persistent (nausea, vomiting, or diarrhea)
- Lacerations requiring sutures **without** excessive blood loss
- Pain, moderate, following abdominal trauma
- Seizure, atypical, in a student with a history of seizures
- Wheezing

INTERVENTIONS

- Support C-ABCDE as indicated
- Determine need for EMS
- Observe student closely
- Initiate appropriate interventions as per specific protocol or IHP and EAP/ECP
- Contact parent/guardian to transport student to medical care or home
- Follow-up

NONURGENT

Findings that suggest a nonacute condition that is not severe, but may require referral for routine medical care. These include

- Essentially well with signs/symptoms (S/S) of mild noncommunicable illness or URI
- Headache **without** fever or other abnormal findings
- Injury, minor (abrasions, ecchymoses, or sprains/strains)
- Pain, mild (e.g., abdominal or menstrual pain, headache, or toothache), **without** fever or other abnormal findings

INTERVENTIONS

- Initiate appropriate interventions or administer medications as per specific protocol or IHP and EAP/ECP
- Observe student
- Contact parent/guardian
- Return student to class or send home as indicated
- Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Abdominal Pain

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

KEY ASSESSMENT POINTS FOR ABDOMINAL PAIN

- Focused abdominal examination
- Time of last bowel movement
 - Urinary symptoms
- Events preceding episode, including trauma
- Menstrual history and possibility of pregnancy
- History/pattern of previous occurrences

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Capillary refill time exceeds 2 sec ▪ Change in mental status or level of consciousness (LOC) ▪ Decreased or absent bowel sounds ▪ Abdominal/GU trauma (see <i>Trauma</i> protocol) ▪ Abdominal distention, rigidity, or guarding ▪ Bluish discoloration of flank or periumbilical area (Grey Turner sign) ▪ Severe abdominal pain ▪ Imminent childbirth (see <i>Obstetric Emergencies</i> protocol) ▪ Pregnant with vaginal bleeding <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Place student supine ▪ Give nothing by mouth ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p>Stable vital signs with</p> <ul style="list-style-type: none"> ▪ Moderate abdominal pain or dysuria ▪ Nausea/vomiting or significant diarrhea ▪ Significant pain with fever ▪ Mucus or frank blood in stool; tarry stools ▪ Abnormal vaginal bleeding or discharge without pregnancy <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ Observe student closely ▪ Give nothing by mouth ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<p>Stable vital signs with</p> <ul style="list-style-type: none"> ▪ Mild or intermittent abdominal pain/cramps ▪ Onset related to menses <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ If recurrent abdominal pain (RAP) has been diagnosed, allow student to rest in health office ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Refer to medical care or school support services as indicated for frequent complaints ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Anaphylaxis/Allergic Reaction

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

KEY ASSESSMENT POINTS FOR ANAPHYLAXIS

- **Respiratory assessment**
 - Focused physical examination of skin findings
- History of systemic allergic reaction
- History of food allergy
- Events preceding reaction, such as a bite/sting

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

<p style="text-align: center;">EMERGENT</p> <ul style="list-style-type: none"> ▪ S/S of cardiopulmonary compromise (see <i>Assessment Tools</i>) ▪ Airway compromise ▪ Change in mental status or LOC ▪ Cyanosis at mouth and lips ▪ S/S of severe respiratory distress (wheezing, dyspnea) ▪ Signs of shock/hypotension ▪ History of anaphylaxis ▪ Edema of face, lips, eyes, or tongue ▪ Generalized hives involving large area ▪ Diaphoresis ▪ Complains of (C/o) tightness in throat or chest ▪ C/o apprehension and/or weakness <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Administer prescribed autoinjectable epinephrine if available ▪ Repeat autoinjectable epinephrine in 5 min if no response ▪ Initiate CPR if necessary ▪ For severe respiratory distress, administer prescribed bronchodilator ▪ Consult IHP and EAP/ECP ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p style="text-align: center;">URGENT</p> <ul style="list-style-type: none"> ▪ S/S of mild systemic reaction; e.g., localized hives, abdominal cramps, nausea, or vomiting ▪ Edema of extremities ▪ Persistent coughing ▪ Tingling, itching of face, ears, or nose ▪ History of allergy <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ Consult IHP and EAP/ECP ▪ Observe student closely ▪ Administer prescribed autoinjectable epinephrine if available and activate EMS ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<p style="text-align: center;">NONURGENT</p> <ul style="list-style-type: none"> ▪ Local reaction only ▪ Responsive to medications ▪ Nasal congestion ▪ Persistent sneezing <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Consult IHP and EAP/ECP ▪ Apply cold pack to site ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy
---	--	--

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Asthma Episode (Acute)/Reactive Airway Disease

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

KEY ASSESSMENT POINTS FOR ACUTE ASTHMA EPISODES

- Airway examination to rule out (R/O) obstruction due to infection or foreign body aspiration
- Respiratory assessment
- Skin assessment

IMMEDIATE INTERVENTIONS

Even *before* you determine triage category, perform the following actions as indicated:

- Help student into a position of comfort
- Administer prescribed bronchodilator or other medication as directed
- Perform peak flow assessment if possible

Note: Obtain peak expiratory flow reading **before** administering bronchodilator and again 20 min later (or per EAP/ECP orders)

Note: Use spacer or holding chamber with MDI/nebulizer, if available

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

<p style="text-align: center;">EMERGENT</p> <ul style="list-style-type: none"> ▪ S/S of severe asthma (see <i>Assessment Tools</i>) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Prepare to ventilate if necessary ▪ Activate EMS if S/S are not relieved by medication or if medication is not available ▪ Administer high-flow O₂ if available ▪ Repeat prescribed bronchodilator/ other medications ▪ Directly/continuously observe student ▪ Consult IHP and EAP/ECP ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p style="text-align: center;">URGENT</p> <ul style="list-style-type: none"> ▪ S/S of moderate asthma (see <i>Assessment Tools</i>) ▪ Cannot tolerate normal activity ▪ No improvement within 15–30 min of bronchodilator administration ▪ Bronchodilator unavailable <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ Administer high-flow O₂ if available ▪ Repeat prescribed bronchodilator/other medications ▪ Consult IHP and EAP/ECP ▪ Directly/continuously observe student ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<p style="text-align: center;">NONURGENT</p> <ul style="list-style-type: none"> ▪ S/S of mild asthma (see <i>Assessment Tools</i>) ▪ Symptoms respond to bronchodilator ▪ Student is able to maintain normal level of activity <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Repeat prescribed bronchodilator/other medications ▪ Consult IHP and EAP/ECP ▪ Monitor student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Assess need for parent/guardian–student asthma education ▪ Follow-up as needed or per policy
---	--	---

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student’s clinical presentation and authorized policies.

Bites and Stings

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR BITES AND STINGS

- Time bite/sting occurred
- Location of bite/sting on body
 - Type of bite/sting
 - Number of bites/stings
 - Intensity of pain
- Previous exposure/allergic/anaphylactic reaction to same type of bite/sting
- Wound characteristics (erythema, edema, ecchymosis, drainage, and size/depth)
- Inspection for foreign body (stinger, tooth, or tick)

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ S/S of anaphylaxis or history of anaphylactic reaction (see <i>Anaphylaxis</i> protocol) ▪ S/S of respiratory distress ▪ Hypotension ▪ Cardiac arrest ▪ Loss of consciousness ▪ Known exposure to toxin (see <i>Toxic Exposure</i> protocol) ▪ Severe pain <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Perform CPR as appropriate ▪ Administer IM epinephrine as per EAP/ECP ▪ Directly/continuously observe student ▪ Report animal bites to appropriate local official ▪ Contact poison center as appropriate (800-222-1222)^a ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ S/S of mild systemic reaction with wheezing, progressive pain/edema but normal vital signs (see <i>Anaphylaxis</i> protocol) ▪ Deep puncture wounds ▪ Moderate pain ▪ Lacerations that require sutures (see <i>Lacerations/Abrasions</i> protocol) ▪ Nausea/vomiting ▪ Human bite with broken skin <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ Wash lacerations with soap and water; irrigate as indicated ▪ Observe student closely ▪ Report animal bites to appropriate local official ▪ Refer all incidents involving human bites to ED ▪ Contact Poison Center as appropriate (800-222-1222)^a ▪ Contact parent/guardian to transport to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Mild localized allergic reaction without systemic/respiratory S/S ▪ Mild pain ▪ Mild pruritus ▪ Stinger/tick present <p style="text-align: center;">INTERVENTIONS</p> <p style="text-align: center;">Stinger</p> <ul style="list-style-type: none"> ▪ Remove stinger by scraping with stiff cardboard/credit card; do not squeeze ▪ Apply cold pack ▪ Observe student for 20 min before returning to class <p style="text-align: center;">Tick</p> <ul style="list-style-type: none"> ▪ Grasp tick with fine-point tweezers or gloved fingertips as close to skin as possible ▪ Pull upward with steady, even pressure ▪ Wash bite area and hands with soap and water <p style="text-align: center;">In All Cases</p> <ul style="list-style-type: none"> ▪ Observe student ▪ Contact parent/guardian ▪ Contact Poison Center as appropriate (800-222-1222)^a ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

First Aid Don'ts

- **Do not** apply a tourniquet for bites or stings because it can lead to ischemia. Use of tourniquets is limited to uncontrolled hemorrhage.
- **Do not** incise wound or apply suction; these methods are ineffective and potentially dangerous.

^aTo contact the Poison Center (800-222-1222): identify yourself as a healthcare professional and provide your name and phone number; student's name, age, weight, and vital signs; insect or animal involved (if known); time, route, and duration of exposure; abnormal S/S; first aid and immediate interventions rendered.

Refer student for tetanus booster if it has been 5 years or more since the last vaccination. Tetanus booster is recommended every 10 years.

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Burns

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*; also see *Burn Assessment* information on next page) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR BURN INJURIES

- Scene safety (e.g., live electrical wire)
- Student's age, weight, and general health status
- Burn characteristics (see next page)
- Associated injuries
- Duration of contact with burn source
- Glasgow Coma Scale (GCS) score

IMMEDIATE INTERVENTIONS

Even before you determine triage category, perform the following actions as indicated:

- Remove student from burn source (e.g., heat/electricity)
- Begin irrigation of chemical burns with cool water (clear off any chemical powders/dust before irrigation)
- Remove jewelry, rings, and constricting clothing (if possible)
- Cool thermal burns if less than 20% of total body surface area (% TBSA) is involved
- **Do not remove clothing that has adhered to skin!**
- **NOTE:** Protect student from hypothermia; only immerse superficial or partial-thickness burns less than 5% TBSA

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

EMERGENT

- Apnea or pulseless
- Full-thickness burn (tough, brownish surface)
- Deep partial-thickness burn (charred/white)
- Burns involving face, hands, feet, eyes, ears, or genitalia (see *Eye or Ear Emergencies* protocols)
- Electrical burns
- S/S of inhalation injury (singled nasal hair or carbonaceous sputum) or other complicating injuries
- Altered mental status
- Respiratory distress (see *Respiratory Distress* protocol)
- Suspected child maltreatment (see *Child Maltreatment* protocol **after** providing burn care)
- Severe pain

INTERVENTIONS

- Activate EMS
- Support C-ABCDE

Electrical burns

- Safely remove student from burn source
- Initiate CPR as needed/have AED ready
- Inspect for entrance and exit wounds

Chemical burns

- Continue copious irrigation of chemical burns with tepid running water for at least 20 min
- Consider contacting Poison Center (800-222-1222)
- Send burn agent and SDS to ED with student
- **Do not** apply cold packs

Thermal burns

- Cover with dry, sterile dressings or clean sheet

In all cases

- Directly/continuously observe student
- Contact parent/guardian
- Notify school administrator
- Follow-up

URGENT

- Superficial partial-thickness burn without complicating factors
- Moderate pain
- Erythema/edema, wet/oozing blisters
- Too large to cover with adhesive bandage
- Signs of associated infection

INTERVENTIONS

- Determine need for EMS
- Flush copiously with tepid running water
- If unable to immerse, apply clean, wet, cool cloth
- **Do not apply cold packs**
- Do not break blisters
- Bandage loosely
- Observe student closely
- Contact parent/guardian to transport student to medical care or home
- Follow-up

NONURGENT

- Superficial burn (e.g., sunburn)
- Minor erythema
- Local, mild pain
- Student is alert

INTERVENTIONS

- Immerse area in tepid water 2–5 min
- Apply tepid cloths
- Bandage loosely
- Observe student
- Contact parent/guardian
- Return student to class or send home as indicated
- Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Burn Assessment


The depth, extent, and location of a burn provide a consistent basis for conveying information about a burn injury to EMS and other healthcare providers. The following parameters are used to describe depth:

- A **superficial** burn involves only the epidermis. It is characterized by erythema and local pain.
- A **superficial partial-thickness** burn involves both the epidermis and the corium. This type of burn generally produces erythema and blisters.
- A **deep partial-thickness** burn may appear white and dry, with locally reduced sensitivity to touch and pain.
- A **full-thickness** burn has a tough brownish surface and a hard eschar. The area will be locally insensitive to touch or pain. This type of burn will not heal without intervention.

Extent is described as a percentage of the body surface area (% TBSA). A quick way to determine % TBSA for small or irregular burns is to use the student's hand (including the palm and fingers) as a reference; this is approximately equal to 1% of the student's total body surface area. The Figure on the left illustrates % TBSA by anatomical area for an infant, child, and adult; the Figure on the right illustrates the palm-and-hand method of estimating % TBSA based on the child's hand size.

% TBSA by anatomical area

Palm-and-hand calculation^a



^a Include child's fingers

Chest Pain

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

KEY ASSESSMENT POINTS FOR CHEST PAIN

Consider etiology

- Cardiac
- Respiratory
- Musculoskeletal
- Trauma
- Psychosocial stress

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Severe chest pain ▪ Bradycardia/tachycardia ▪ Cyanosis of lips and nail beds ▪ Altered mental status ▪ Palpitations ▪ Dyspnea ▪ Peripheral pulses weak/thready/absent ▪ Diaphoresis; clammy, cool skin ▪ Restlessness ▪ Hypotension ▪ Nausea ▪ Weakness ▪ Capillary refill exceeds 2 sec <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Have AED readily available ▪ Maintain position of comfort ▪ Directly/continuously observe student ▪ Reassess vital signs every 5 min ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Moderate, persistent chest pain ▪ Anxiety ▪ Stable vital signs ▪ No history of <ul style="list-style-type: none"> • Recent chest trauma • Recent asthma episode • Loss of consciousness <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Maintain position of comfort ▪ Observe student closely ▪ Reassess vital signs ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Mild chest pain ▪ Normal vital signs ▪ No history of <ul style="list-style-type: none"> • Recent chest trauma • Recent asthma episode • Loss of consciousness <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Maintain position of comfort ▪ Monitor closely ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Chest Trauma

SYSTEMATIC ASSESSMENT

**Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.
Provide spinal motion restriction if head/spinal injury is suspected!**

KEY ASSESSMENT POINTS FOR CHEST TRAUMA

- Respiratory assessment
- Chest wall inspection for symmetry, wounds, and/or ecchymoses

IMMEDIATE INTERVENTIONS

Even before you determine triage category, perform the following actions as indicated:

- If student is apneic, perform rescue breathing using mouth-to-mask
- If student is pulseless, initiate CPR; apply AED
- Apply clean nonporous dressing to open chest wounds (if S/S of tension pneumothorax develop, open the dressing)
- Apply direct pressure for profuse hemorrhage

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ S/S of respiratory distress ▪ Open chest wound ▪ Crush injury ▪ Uncontrollable hemorrhage ▪ Muffled heart sounds ▪ Paradoxical or asymmetrical chest wall movement ▪ S/S of shock ▪ S/S of tension pneumothorax or cardiac tamponade: <ul style="list-style-type: none"> • Apprehension • Rapid/shallow respiration • Painful respiration • Jugular vein distension • Cyanosis • Muffled heart sounds • Hypotension (late/ominous sign) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Maintain spinal motion restriction ▪ Cover open chest wounds (occlude on three sides only) ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Closed chest injury without respiratory distress ▪ S/S of closed rib fracture (shallow/painful but unimpaired respiration) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Monitor respiratory/cardiac status ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Stable vital signs ▪ Ecchymoses ▪ No S/S of rib fracture <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Apply cold packs as appropriate ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

NOTE Pneumothorax/cardiac tamponade may develop slowly during a 24–48 hour period following chest trauma; reassessment is crucial. A pneumothorax should be considered emergent; a tension pneumothorax may form if left untreated. Signs/symptoms (S/S) of pneumothorax include tachypnea, tachycardia, unequal chest expansion, and unilateral absent or diminished breath sounds.

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Child Maltreatment, Suspected

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*; also see *Indications of Child Maltreatment* protocol on next page) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR SUSPECTED CHILD MALTREATMENT

- Indications of neglect
- Indications of abuse (see next page)

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT

- Multiple new suspicious injuries (see *Trauma* protocol)
- History of chronic life-threatening illness without appropriate medical treatment

INTERVENTIONS

- Activate EMS
- Support C-ABCDE
- Treat injuries
- Provide emotional support
- Directly/continuously observe student
- **Report suspicions to DCFS: 800-25-ABUSE (22873)**
- Notify crisis response team
- Notify school administrator
- **Document all findings** (see Appendix B for a written report form)
- Follow-up

URGENT

- New injuries **and** history of suspicious injuries (see *Trauma* protocol)
- Appears inadequately fed, clothed, or sheltered
- Inadequate medical care

INTERVENTIONS

- Treat injuries
- Provide emotional support
- Observe student closely
- **Report suspicions to DCFS: 800-25-ABUSE (22873)**
- Notify crisis response team as appropriate
- **Document all findings** (see Appendix B for a written report form)
- Follow-up

NONURGENT

- Student reports abandonment by parent/guardian
- Student reports maltreatment

INTERVENTIONS

- Provide emotional support
- **Report suspicions to DCFS: 800-25-ABUSE (22873)**
- Notify crisis response team as appropriate
- Observe student
- **Document all findings** (see Appendix B for a written report form)
- Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Indications of Child Maltreatment

As a school nurse, you are required by law to report any suspicion of child maltreatment (abuse or neglect) to the appropriate agency in your state. In Illinois, you can reach the Child Abuse Hotline maintained by the Department of Children and Family Services (DCFS) by calling **800-25-ABUSE** (22873). The law provides immunity for mandated reporting.

INDICATORS OF ABUSE

The following findings should increase your suspicion of child abuse:

- The reported history of injury is inconsistent with the physical examination findings.
- Details of the reported history change from one telling to the next.
- There was a prolonged delay between the time of injury and the time medical assistance was sought.
- The child has a history of repeated trauma.
- The parent/guardian responds to questions inappropriately or does not comply with medical advice.

Suspicious injuries include

- Injuries involving soft tissue of the face, neck, abdomen, or similar areas.
- Injuries involving areas that are normally shielded, including the back and chest.
- Fractures of long bones in children younger than 3 years.
- Old scars or injuries in different stages of healing.
- Injuries with an appearance suggesting deliberate infliction, such as human bite marks, cigarette burns, rope marks, or the imprint of a belt or other object.
- Trauma affecting the genital or perianal area.
- Sharply demarcated burns in unusual areas.
- Scald patterns that appear to involve submersion in hot water, such as burns to the hands, feet, or buttocks.

INDICATORS OF NEGLECT

The following findings should increase your suspicion of child neglect:

- Unsafe conditions are evident in the home environment (e.g., weapons within reach, open windows without screens or window guards, or perilously unsanitary conditions).
- The parent/guardian has not provided for medical treatment, refuses to permit medical treatment, or fails to seek necessary and timely medical care for a child who has an acute or chronic life-threatening illness.
- A child younger than 10 years has been left unattended or unsupervised (although some situations permit a parent/guardian to leave a young child alone without endangerment, you cannot make this determination).
- The child appears to be abandoned.
- The parent/guardian appears to be incapacitated due to intoxication, disabling psychiatric problems, debilitating illness, or similar impairment, and cannot adequately care for the child.
- The child appears to be malnourished (seriously underweight, emaciated, or dehydrated), inadequately clothed, or inadequately sheltered.
- The child is found to be intoxicated or under the influence of an illicit substance.

All instances of suspected child maltreatment must be reported to the DCFS for investigation. A sample form for reporting your suspicions in writing appears in Appendix B.

Cold-Related Injuries (Hypothermia/Frostbite)

GENERAL GUIDELINES

- Hypothermia: rewarm slowly
- Frostbite: rewarm quickly
- Do not rub affected area
- Use warm water to rewarm; **never use hot water to rewarm**

Educate students regarding susceptibility to hypothermic injury and need for precautions

FROSTBITE CATEGORIES

Frostnip

- Blanched, white skin that is cold to touch

Superficial frostbite

- Firm, waxy skin with softer tissue underneath
- Blisters develop in 24–48 hours

Deep frostbite

- Mottled or gray-blue skin that is firm to touch
- Severity not apparent until frostbitten area is rewarmed
- Sensory: cold; pruritus or paresthesia
- Necrosis develops over time

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR COLD-RELATED INJURIES

- Skin assessment
- Events preceding episode
- Duration of exposure
- Focused physical examination of extremities

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT

- Altered mental status
- Cyanosis
- Slow/shallow respiration
- Weak, thready pulses, no pulses, or bradycardia
- Hypotension
- Extremities edematous and discolored
- No shivering (severe hypothermia)
- Slurred speech
- Abnormally low body temperature
- S/S of deep frostbite

INTERVENTIONS

- Activate EMS
- Remove student's wet clothing
- Keep student warm and completely covered
- Assess vital signs for **1 full minute**
- If pulseless, initiate CPR
- Directly/continuously observe student
- Contact parent/guardian
- Notify school administrator
- Follow-up

URGENT

- Normal LOC or drowsy
- Shivering (mild hypothermia)
- S/S of superficial frostbite or frostnip

INTERVENTIONS

- Remove wet clothing and keep student warm
- Determine need for EMS
- Observe student closely
- If alert, give sips of warm liquid
- Warm area with warm water
- Reinforce need for precautions when exposed to cold
- Contact parent/guardian to transport student to medical care or home
- Follow-up

NONURGENT

- Alert
- Slight shivering
- Exposed skin feels cold
- No signs of frostbite or frostnip

INTERVENTIONS

- Remove wet clothing and replace with dry coverings
- If student is alert, give sips of warm liquid
- Warm cold skin with warm water
- Observe student
- Reinforce need for precautions when exposed to cold
- Contact parent/guardian
- Return student to class or send home as indicated
- Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Dental, Oral, and Maxillofacial Emergencies

NOTE

Refer student for tetanus booster if it has been 5 years or more since the last vaccination. Tetanus booster is recommended every 10 years.

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

Provide spinal motion restriction if head/spinal injury is suspected!

KEY ASSESSMENT POINTS FOR DENTAL, ORAL, AND MAXILLOFACIAL EMERGENCIES

- Inspect teeth
- Assess ability to open and close mouth
- Assess facial bones, including mandible

IMMEDIATE INTERVENTIONS

Even before you determine triage category, ensure the adequacy of the student’s airway.

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Airway compromise ▪ Change in mental status or LOC ▪ Excessive bleeding ▪ Suspected fracture of mandible or other facial bones (see <i>Trauma</i> protocol as indicated) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Maintain spinal motion restriction ▪ Support C-ABCDE ▪ Treat hemorrhage with direct pressure ▪ Apply cold packs to reduce pain/edema ▪ For suspected mandible fracture only, stabilize jaw by wrapping cravat around the point of the chin and securing it on top of the head; avoiding pressure on the neck ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Displacement of multiple teeth ▪ Avulsion of permanent tooth^a ▪ Major chip/fracture of permanent tooth ▪ Broken orthodontic appliance ▪ Severe toothache <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ If possible, save large tooth chips; cover jagged edge of tooth with gauze ▪ For wire protruding from appliance, gently attempt to bend away from oral tissue; if unsuccessful, cover end with gauze or dental wax (do not remove embedded wire) ▪ Observe student closely ▪ Contact parent/guardian to transport student to dental care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Caries ▪ Exfoliation of primary tooth ▪ Eruption of permanent tooth ▪ Bleeding gums ▪ Minor chip/fracture of tooth <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ For minor tooth chip, have student rinse mouth with warm salt water ▪ Apply cold compress to edematous areas ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

^aAvulsion of permanent tooth

Note: Replantation is most likely to succeed if attempted within 60 minutes.

- Activate EMS or transport immediately to dentist
- Handle the tooth by the crown, not the root
- If the tooth is dirty, gently rinse in milk or water
- Gently replace tooth in socket (do not use force) **only** if student is alert and able to cooperate; instruct student to keep pressure on tooth by biting gently on clean gauze
- If unable to replace tooth in socket, place the tooth in a protective container filled with pH-balanced solution for transport with student to the ED or dental specialist; if this solution is not available, use cold milk or water

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student’s clinical presentation and authorized policies.

Diabetic Emergencies

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR DIABETIC EMERGENCIES

- Respiratory assessment
 - Skin assessment
- Last insulin dose/type of insulin used and delivery route
- Last meal/carbohydrate intake
- Precipitating factors (exercise, change in eating habits/diet, stress, missed insulin dose, or illness)
- Current blood glucose level

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<p>Severe hypoglycemia Glucose less than 60 mg/dL with loss of consciousness or seizures (see <i>Seizures</i> protocol)</p> <p>Severe hyperglycemia/diabetic ketoacidosis (DKA) Glucose exceeds 200 mg/dL with at least one of the following:</p> <ul style="list-style-type: none"> ▪ Moderate to severe dehydration ▪ Abdominal pain/tenderness ▪ Kussmaul respiration and/or fruity breath odor ▪ Tachycardia ▪ Cool extremities ▪ Altered mental status (lethargic to comatose) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Directly/continuously observe student ▪ Place in left lateral recovery position to prevent aspiration ▪ Consult IHP and EAP/ECP Hypoglycemia <ul style="list-style-type: none"> ▪ Administer 0.5 mg or 1 mg glucagon IM/IN/SQ if available per EAP/ECP (place student in recovery position because glucagon may precipitate vomiting) ▪ Severe hyperglycemia Diabetic ketoacidosis <ul style="list-style-type: none"> ▪ Administer insulin per EAP/ECP ▪ Prepare for immediate transport ▪ Give nothing by mouth ▪ In all cases <ul style="list-style-type: none"> ▪ Contact parent/guardian ▪ Notify school administrator ▪ Determine need for diabetes education (parent/guardian, student, and school personnel) ▪ Follow-up 	<ul style="list-style-type: none"> ▪ S/S of moderate hypoglycemia; glucose less than 60 mg/dL, student awake and responsive ▪ S/S of moderate hyperglycemia; glucose 120–200 mg/dL; student awake and responsive with mild abdominal pain/tenderness, nausea, headache, tachycardia, and/or fruity breath odor <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Monitor student closely ▪ Consult IHP and EAP/ECP Hypoglycemia <ul style="list-style-type: none"> ▪ Give glucose tablets/instant glucose equivalent to 15 grams carbohydrates, or 4 oz regular soda or juice (orange/apple), followed by snack/next meal ▪ As indicated, give extra snack (e.g., two peanut butter crackers or one-half sandwich and 8 oz milk) ▪ Hyperglycemia <ul style="list-style-type: none"> ▪ Administer insulin per EAP/ECP ▪ Increase intake of water ▪ Test for ketonuria ▪ In all cases <ul style="list-style-type: none"> ▪ Contact parent/guardian to transport student to medical care or home ▪ Determine need for diabetes education (parent/guardian, student, and school personnel) ▪ Follow-up 	<ul style="list-style-type: none"> ▪ S/S of mild hypoglycemia; glucose 60–80 mg/dL, student awake and alert ▪ S/S of mild hyperglycemia; glucose 120–200 mg/dL, student awake and alert <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Consult IHP and EAP/ECP Mild hypoglycemia <ul style="list-style-type: none"> ▪ Give glucose tablets/instant glucose equivalent to 15 grams carbohydrates, or 4 oz regular soda or juice (orange/apple), followed by snack/next meal ▪ As indicated, give extra snack (e.g., two peanut butter crackers or one-half sandwich and 8 oz milk) ▪ Recheck glucose in 15 min; if no improvement, repeat treatment ▪ If improvement is noted, return student to class or send home as indicated ▪ Instruct student to refrain from tasks requiring intense concentration or exertion for 1 hour ▪ Mild hyperglycemia <ul style="list-style-type: none"> ▪ Administer insulin per EAP/ECP ▪ Increase intake of water ▪ Return student to class or send home as indicated ▪ In all cases <ul style="list-style-type: none"> ▪ Contact parent/guardian as needed or per policy ▪ Determine need for diabetes education (parent/guardian, student, and school personnel) ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Disaster/Emergency Response: Mass-Casualty Incidents

DEFINITION

A mass-casualty incident is a situation in which medical care requirements overwhelm the local emergency response resources

IMMEDIATE ACTIONS

- Assess scene safety
- Activate EMS
- Determine approximate number of casualties
- Activate incident command

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

- Perform triage assessment using START/JumpSTART® algorithms for MCI triage (see next page)
- Assign appropriate triage categories
- Dispatch casualties to designated field treatment area


IMMEDIATE	DELAYED	MINOR	EXPECTANT/ DECEASED
<p>Life-threatening</p> <ul style="list-style-type: none"> ▪ Shock ▪ Respiratory distress/failure ▪ Major burns ▪ Fracture of long bone with circulatory compromise <p>INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Assign team members to multiple casualties ▪ Maintain C-ABCDE ▪ Ensure direct, continuous observation until transfer of care ▪ Notify parent/guardian as specified in emergency response plan ▪ Assist EMS as appropriate ▪ Follow-up 	<p>Care required within 1–2 hours</p> <ul style="list-style-type: none"> ▪ Fracture of long bone without circulatory compromise ▪ Laceration without significant blood loss ▪ Head injury without loss of consciousness <p>INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Maintain C-ABCDE ▪ Initiate appropriate care ▪ Notify parent/guardian as specified in emergency response plan ▪ Assist EMS as appropriate ▪ Provide for counseling ▪ Follow-up 	<p>Nonacute/minor condition</p> <ul style="list-style-type: none"> ▪ Major abrasions/ecchymoses ▪ Muscle sprains/strains ▪ Emotional distress <p>INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Initiate appropriate care ▪ Monitor for changes ▪ Notify parent/guardian as specified in emergency response plan ▪ Provide for counseling ▪ Follow-up as needed or per policy 	<p>Survival unlikely</p> <ul style="list-style-type: none"> ▪ Massive open head trauma ▪ Cardiac arrest <p>INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Do not render care until adequate resources and personnel permit ▪ Follow-up as needed or per policy

EMS TRIAGE CATEGORIZATION FOR MASS-CASUALTY INCIDENTS


Category	Condition
Immediate	Potential threat to life or function that requires immediate intervention
Delayed	Acute condition that does not threaten life or limb; requires care within 1–2 hours
Minor	Nonacute or minor condition
Expectant/ Deceased	Catastrophic injury or condition such that survival is unlikely (in a disaster situation with mass casualties, limited medical resources do not support the resuscitation and care of these patients)

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student’s clinical presentation and authorized policies.

START/JumpSTART® Algorithms for Mass-Casualty Incident Triage



© Lou Romig, MD 2002



Drowning/Submersion Injuries

CAUTION

Ensure scene safety! Never attempt a water rescue unless you are trained to do so.

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR SUBMERSION INJURIES

- Skin assessment
- Duration of exposure
- Inspection for associated injuries

KEY FINDINGS

- Dyspnea, crackles, rhonchi, wheezing, or apnea
- Bradycardia or asystole
- Cyanosis or pallor
- Altered mental status
- Fixed, dilated pupils
- Hypothermia and cool skin

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

The triage category for ALL instances of drowning/submersion is EMERGENT!

INTERVENTIONS

- Activate EMS
- EMS transport to ED for evaluation is REQUIRED in all cases of drowning/submersion!**

If still in water

- Do not remove student without assistance
- Keep student afloat face-up
- Support head and neck in neutral alignment with spine
- Open airway using the jaw-thrust technique and support ventilation
- When adequate assistance is available, place student supine on backboard or other rigid support for removal from water

If out of water

- Support C-ABCDE (use jaw-thrust technique if spinal injury is suspected)
- Assess Glasgow Coma Scale score
- Cover student and maintain warmth to prevent hypothermia

In all cases

- Directly/continuously observe student
- Contact parent/guardian
- Notify school administrator
- Follow-up

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Ear Emergencies

SYSTEMATIC ASSESSMENT

**Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.
Provide spinal motion restriction if head/spinal injury is suspected!**

KEY ASSESSMENT POINTS FOR EAR EMERGENCIES

- Skin assessment for wounds, blisters, erythema, edema, hematomas, or bleeding
- Focused physical examination (external/otoscopic)
- Hearing evaluation

IMMEDIATE INTERVENTIONS

Even before you determine triage category, perform the following actions as indicated:

- Control bleeding
- Apply appropriate dressing

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT

- Change in mental status
- Laceration/avulsion/hematoma of external ear with uncontrollable bleeding (see *Head/Spinal Cord* protocol as indicated)
- Burn or direct thermal injury
- Acute hearing loss

INTERVENTIONS

- Activate EMS
- Maintain spinal motion restriction if applicable
- Support C-ABCDE
- Directly/continuously observe student
- Maintain position of comfort
- Keep student calm
- Contact parent/guardian
- Notify school administrator
- Follow-up

URGENT

- Foreign body in ear
- Mild hematoma
- Abrasions/minor lacerations of external ear (see *Lacerations/Abrasions* protocol)

INTERVENTIONS

- Determine need for EMS
- **Foreign body**
- Do not attempt to remove unless object is visible and can be gripped with forceps or fingers
- If object is a live insect, instill 1–2 drops of mineral oil
- **In all cases**
- Closely monitor student
- Contact parent/guardian to transport student to medical care or home
- Maintain position of comfort
- Observe student
- Follow-up as needed or per policy

NONURGENT

- Mild earache without drainage
- Associated low-grade fever

INTERVENTIONS

- Send student home if pain is persistent **or** accompanied by fever
- Contact parent/guardian
- Return student to class or send home as indicated
- Follow-up as needed or per policy

NOTE

Refer student for tetanus booster if it has been 5 years or more since the last vaccination. Tetanus booster is recommended every 10 years.

There is a high risk for long-term complications with injury to the ear. Always refer the student for evaluation by a Healthcare Practitioner.

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Eating Disorders

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR EATING DISORDERS

- History of food intake and level of exercise
- Skin assessment
- Menstrual history
- Mental status examination or other brief psychosocial profile
- Focused physical examination of weight (have student remove shoes/bulky outer wear)

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

<p style="text-align: center;">EMERGENT</p> <ul style="list-style-type: none"> ▪ Cardiac arrest ▪ Seizure activity (see <i>Seizures</i> protocol) ▪ Hypotension ▪ Bradycardia ▪ Lethargy <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p style="text-align: center;">URGENT</p> <ul style="list-style-type: none"> ▪ Significant weight loss ▪ Tooth enamel erosion ▪ Weakness ▪ Poor skin turgor <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Encourage fluid intake in small quantities ▪ Provide nonthreatening environment ▪ Observe student closely ▪ Discuss the need for medical evaluation and/or counseling with parent/guardian ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<p style="text-align: center;">NONURGENT</p> <ul style="list-style-type: none"> ▪ Normal vital signs ▪ Suspicion or early signs of eating disorder^a <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Provide nonthreatening environment ▪ Observe student ▪ Discuss health consequences of behavior ▪ Contact parent/guardian ▪ Refer to school counselor ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy
--	--	--

^aEarly signs and psychosocial attributes associated with eating disorders

- High achiever/perfectionist
- Low self-esteem/depression
- History of substance abuse
- Intense fear of weight gain
- Evidence of body dysmorphia
- Rigid self-control
- Reduced food intake, self-induced vomiting, and/or use of laxatives/diuretics/emetics
- Vigorous exercising to achieve weight loss rather than for fitness
- Recent history of weight loss/weight fluctuations
- Preference for oversized clothing
- Amenorrhea
- Hypothermia
- Lanugo (downy hair)
- Weakness
- Poor skin turgor
- Esophagitis or oral lesions
- Dental caries or tooth enamel erosion

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Eye Emergencies

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

Provide spinal motion restriction if spinal injury is suspected!

KEY ASSESSMENT POINTS FOR EYE EMERGENCIES

- Events leading up to injury (e.g., chemical exposure or other burn)
- AS TOLERATED, focused physical examination of the eye/vision for
 - Visible wounds, drainage, or foreign body
 - Extraocular movement
 - PERRL (pupils equal, round, and reactive to light)
 - Visual acuity

IMMEDIATE INTERVENTIONS

For chemical burns involving the eye, IMMEDIATELY activate EMS, then ensure scene safety and begin flushing the eye copiously with saline, eyewash solution, or water.

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Change in mental status ▪ Penetrating injury ▪ Chemical/thermal burn ▪ Unequal/irregular pupils ▪ Blunt injury (see <i>Head/Spinal Cord</i> protocol) ▪ Embedded foreign body ▪ Hyphema (haze or blood in iris) ▪ Reduced visual acuity/loss of vision ▪ Bilateral periorbital ecchymoses (raccoon eyes) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Maintain spinal motion restriction as applicable ▪ Support C-ABCDE ▪ Activate EMS ▪ Protect eye from further injury <p style="text-align: center;">Chemical burns</p> <ul style="list-style-type: none"> ▪ Continue irrigation while awaiting EMS ▪ Send copy of SDS to ED <p style="text-align: center;">Penetrating injuries</p> <ul style="list-style-type: none"> ▪ Stabilize object with gauze pads ▪ Tape disposable drinking cup over dressing (do not allow it to contact object) <p style="text-align: center;">Thermal burns</p> <ul style="list-style-type: none"> ▪ Do not inspect eyes ▪ Apply loose, moist dressing <p style="text-align: center;">Radiation burns (From arc welder, sunlight, or sun lamp)</p> <ul style="list-style-type: none"> ▪ Cover with eye patch <p style="text-align: center;">In all cases</p> <ul style="list-style-type: none"> ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Blunt trauma without vision changes ▪ S/S extraocular muscle entrapment ▪ Suspected corneal abrasion ▪ Laceration of lid ▪ Blurry/impaired vision ▪ Diplopia ▪ Eye pain/guarding ▪ Sensation of foreign body that persists for more than 1 hour <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ If no evidence of injury, apply cool compress for 20 min ▪ Instruct student not to move rapidly, bend over, or cough ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Superficial foreign body ▪ Subconjunctival hemorrhage (may result from violent coughing or vomiting) ▪ Minor periorbital lacerations ▪ Minor periorbital ecchymoses <p style="text-align: center;">INTERVENTIONS</p> <p style="text-align: center;">Foreign body</p> <ul style="list-style-type: none"> ▪ If foreign body is visible in sac of lower lid, remove with cotton-tipped applicator ▪ If removal is unsuccessful after two attempts, or if foreign body is located elsewhere, flush eye with saline, eyewash solution, or water <p style="text-align: center;">In all cases</p> <ul style="list-style-type: none"> ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Foreign Body Airway Obstruction

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*; also see *AHA Airway Clearing Maneuvers* protocol on next page) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR FOREIGN BODY AIRWAY OBSTRUCTION

- Evaluation of airway/respiratory status
 - Suddenness of onset
- Events leading up to incident (e.g., witnessed ingestion/aspiration of small object, toy, or food)

Note: Fever or S/S of respiratory illness decreases the likelihood of foreign body etiology.

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<p style="text-align: center;">Severe airway obstruction</p> <ul style="list-style-type: none"> ▪ Choking, silent cough, or ▪ Unable to cough, speak, or make any sound ▪ Apnea ▪ Pallor or cyanosis ▪ Loss of consciousness <p style="text-align: center;">INTERVENTIONS</p> <p style="text-align: center;">Severe obstruction</p> <p>Begin AHA airway clearing maneuvers (see next page for detailed procedure)</p> <ul style="list-style-type: none"> ▪ In infants younger than 1 year, apply five back blows/slaps and five chest thrusts ▪ In children older than 1 year, perform abdominal thrusts (Heimlich maneuver) ▪ Continue until either the object is expelled or the infant or child becomes unresponsive ▪ Begin CPR, checking inside the mouth before each series of rescue breaths to see whether the object is visible and can be removed <p>Note: Do not attempt blind finger sweeps! Foreign body may be pushed further into the airway, exacerbating obstruction</p> <ul style="list-style-type: none"> ▪ Activate EMS if efforts are unsuccessful after 1 minute ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p style="text-align: center;">Mild airway obstruction</p> <ul style="list-style-type: none"> ▪ History of aspiration ▪ Fast breathing ▪ Intermittent wheezing or stridor ▪ Gagging, choking, or coughing ▪ Pink skin color ▪ Mild to moderate dyspnea <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Encourage forceful cough ▪ Do not interfere in any other way ▪ Monitor for worsening distress, ineffective cough, inspiratory wheezing, labored breathing, or tachycardia ▪ If signs of severe obstruction develop, triage as emergent and begin AHA airway clearing maneuvers (see next page) ▪ If student’s efforts clear the obstruction, contact parent/guardian to transport student to medical care or home ▪ Notify school administrator ▪ Provide psychological support ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Witnessed ingestion/aspiration ▪ Student clears obstruction by coughing ▪ No S/S of continued obstruction <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Observe student at frequent intervals throughout day ▪ Contact parent/guardian ▪ Provide psychological support ▪ Return student to class or send home as indicated ▪ NOTE: If student continues to cough the next day, suspect retained bronchial foreign body, bronchitis, or pneumonia ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student’s clinical presentation and authorized policies.

American Heart Association (AHA) Airway Clearing Maneuvers (2020)

Responsive Infant

- Sit or kneel, with the infant in your lap. Hold the infant face down and resting on your forearm, with the head slightly lower than the chest. Support the head and jaw with your hand. Avoid compressing the soft tissue of the infant's throat. Then, rest your forearm on your thigh to provide support.
- With the heel of your other hand deliver up to 5 forceful backslaps between the infant's shoulder blades. Deliver each slap with sufficient force to attempt to dislodge the foreign body.
- Then, place your free hand on the infant's back supporting the head with the palm of your hand. This will cradle the infant between your two forearms as you turn the infant over while carefully supporting the head and neck.
- Keep the infant's head lower than the chest and deliver up to 5 quick downward chest thrusts in the same location you would give compressions, just below the nipple line over the lower half of the breastbone. Do this at the rate of about 1 per second.
- Repeat this sequence of 5 backslaps and 5 chest thrusts until the object is removed or until the infant becomes unresponsive.

Responsive infant



Unresponsive infant



Unresponsive Infant

- Place the infant face up on a hard flat surface. Perform CPR as usual with one difference. Each time you open the airway to give breaths, look for the object.
- To give breaths to an infant, the mouth to mouth and nose technique is preferred. For this technique, you will first open the infant's airway with a head tilt, chin lift. Then, place your mouth over the infant's mouth and nose to create an airtight seal.
- Give one breath, blowing for about 1 second. Watch for the chest to rise as you give the breath. If the chest doesn't rise, repeat the head tilt, chin lift to reopen the airway and try to give a breath that makes the chest rise.
- If you cannot cover the infant's mouth and nose with your mouth, use the mouth to mouth technique instead. The mouth to mouth technique for an infant is the same as an adult.
- If you can see the object, carefully try to remove it. Never perform a blind finger sweep. You should only try to remove an object if you can see it.
- If you're alone, after about 2 minutes, or 5 cycles of CPR, activate the emergency response system.

Responsive child



Responsive Child

- To perform an abdominal thrust, stand or kneel behind the child or adult, and place your arms around the waist and with one hand, locate the navel. Then, make a fist with your other hand and place the thumb side of your fist against the person's abdomen, just above the navel and below the breastbone.
- Grasp your fist with your other hand and press your fist into the abdomen with a quick, forceful upward thrust. Repeat thrusts until the object is expelled from the airway, or the person becomes unresponsive.
- If you can't wrap your arms around the waist because the person is pregnant or overweight, or simply can't stand, wrap your arms around their chest and give chest thrusts instead of abdominal thrusts.

Unresponsive Child / Adult

- If you're not able to relieve choking in an adult or child, or if the person becomes unresponsive, have someone activate the emergency response system (EMS), then lower the person to the ground and begin CPR.
- Perform CPR as usual, but with one difference. Each time you open the airway to give breaths, look for the obstructing object in the back of the throat. If you see an object that can be easily removed, carefully remove it. But, do not perform a blind finger sweep because this may cause the object to become lodged farther back in the airway.

Headache

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR HEADACHES

- Mental status/neurological assessment
- Past health history of headaches or recent head injury
- Events leading up to onset, such as stress, exposure to a known trigger, and/or aura
- Focused physical examination for neck stiffness
- Assessment for visual disturbances (see *Eye Emergencies* protocol)

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

<p style="text-align: center;">EMERGENT</p> <ul style="list-style-type: none"> ▪ Change in mental status ▪ Acute neurological deficit ▪ Seizure activity (see <i>Seizures</i> protocol) ▪ Severe headache (e.g., c/o “worst headache of my life”) ▪ Stiff neck with fever ▪ Recent head injury ▪ Hypertension <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Consult IHP and EAP/ECP ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p style="text-align: center;">URGENT</p> <ul style="list-style-type: none"> ▪ Moderate headache with vomiting ▪ History of aura ▪ Exposure to known trigger ▪ Blurred vision, dizziness, or photophobia ▪ No neurological deficit ▪ History of migraines <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ Provide rest in quiet, darkened room ▪ Consult IHP and EAP/ECP ▪ Administer medication per IHP and EAP/ECP ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<p style="text-align: center;">NONURGENT</p> <ul style="list-style-type: none"> ▪ Generalized mild headache ▪ S/S of upper respiratory infection (URI) ▪ S/S of sinus infection <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Allow student to rest ▪ Consult IHP and EAP/ECP ▪ Administer medication per IHP and EAP/ECP ▪ Observe student ▪ Reassess ▪ Educate about avoiding triggers ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy
---	--	---

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student’s clinical presentation and authorized policies.

Head/Spinal Cord Trauma

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO. Provide spinal motion restriction—DO NOT move student!

KEY ASSESSMENT POINTS FOR HEAD/SPINAL CORD TRAUMA

- Mental status/neurological assessment
 - Events leading up to injury
 - Mechanism of injury
- Assessment for visual disturbances (see *Eye Emergencies* protocol)

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Change in mental status (including drowsiness or lethargy) ▪ Abnormal behavior/loss of normal abilities ▪ Seizure activity (see <i>Seizures</i> protocol) ▪ Loss of/decreased movement or sensation in extremities ▪ Blood/cerebrospinal fluid (CSF) discharge from nose/ears ▪ Significant trauma to head/neck or high-risk mechanism of injury ▪ Evidence of depressed skull fracture ▪ Paresthesia <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE and provide spinal motion restriction ▪ Apply direct pressure to bleeding except over depressed skull injury ▪ Keep student warm ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Brief period of confusion/amenia ▪ Neck pain ▪ Dizziness ▪ Blurred vision/diplopia ▪ Headache ▪ Nausea/vomiting ▪ Laceration requiring sutures (see <i>Lacerations/Abrasions</i> protocol) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Provide spinal motion restriction ▪ Control bleeding with direct pressure ▪ Apply cold packs to swollen areas ▪ Keep student warm ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Alert ▪ Minor abrasions, lacerations, or edema <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Apply cold packs as indicated ▪ Observe student closely ▪ Reassess after 15–30 min ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Advise parent/guardian, teachers, and coaches of injury and emphasize the need for observation during next 24–48 hours; seek medical care if the student exhibits any of the following: <ul style="list-style-type: none"> • Dizziness • Headache • Nausea • Photophobia • Diplopia • Irritability • Poor concentration • Decline in academic ability • Personality changes ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Heat-Related Injuries

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR HEAT-RELATED INJURIES

- Skin assessment (e.g., color, temperature, moisture, and other related skin findings)

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

<p style="text-align: center;">EMERGENT</p> <ul style="list-style-type: none"> ▪ S/S of heat stroke <ul style="list-style-type: none"> • Hyperthermia (temperature exceeding 104°F/40°C) • Confusion/altered mental status • Hot/dry/red skin • Tachycardia/weak peripheral pulses • Syncope • Ataxia ▪ Seizure activity (see <i>Seizures</i> protocol) <p style="text-align: center;">INTERVENTIONS</p> <p style="text-align: center;">Heat stroke is a life-threatening emergency!</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Immediately remove from heat to a cool environment ▪ Immediately initiate cooling measures: <ul style="list-style-type: none"> • Loosen clothing • Apply cool, wet towels to neck, groin, and axillae • Sponge with cool compresses • Fan student ▪ Place in left lateral recovery position in case of vomiting ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p style="text-align: center;">URGENT</p> <ul style="list-style-type: none"> ▪ Oriented ▪ Mild tachycardia ▪ S/S of heat exhaustion <ul style="list-style-type: none"> • Cool, moist, and pale skin • Dilated pupils • Mild to moderate headache • Nausea/vomiting • Muscle cramps • Weakness or dizziness • Normothermic to mildly hyperthermic (temperature less than 101°F/38.3°C) • Diaphoresis <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ Observe continuously ▪ Allow to rest in cool environment ▪ Loosen clothing ▪ Apply cool, wet towels ▪ Fan student ▪ In the absence of vomiting, encourage fluid replacement with water or a diluted electrolyte-replacement drink as permitted by applicable protocols ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<p style="text-align: center;">NONURGENT</p> <ul style="list-style-type: none"> ▪ Mild cramping of calves, thighs, or shoulders ▪ Normothermic ▪ Awake and alert <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Allow to rest in cool environment ▪ Loosen clothing ▪ Observe student ▪ Encourage fluid replacement with water or a diluted electrolyte-replacement drink as permitted by applicable protocols ▪ Educate student about maintaining adequate hydration during higher-risk activities ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy
--	---	---

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Hemophilia

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

KEY ASSESSMENT POINTS FOR HEMOPHILIA

- History relating to recent surgery, medical procedures, dental extractions, or injuries
- Focused physical examination for external bleeding, lacerations, ecchymoses, abrasions, and/or hematomas

IMMEDIATE INTERVENTIONS

Even *before* you determine triage category, attempt to control external bleeding by applying firm pressure for 10 min.

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ S/S of internal bleeding: <ul style="list-style-type: none"> • Headache • Dizziness • Visual disturbances • Neurological deficit • Altered mental status/pupillary changes ▪ Signs of shock ▪ Profuse, uncontrollable hemorrhage ▪ Severe abdominal pain <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ For shock, place in Trendelenburg position ▪ Consult IHP and EAP/ECP ▪ Directly/continuously observe student ▪ Frequently reassess vital signs and AVPU ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Acute joint swelling/pain ▪ Abdominal discomfort, tenderness on palpation, or nausea ▪ External bleeding that does not diminish with prolonged pressure ▪ Indications of intramuscular hematoma (most commonly felt in knees, ankles, and elbows): tingling, pain, limited range of motion (ROM), edema, and/or increased warmth/tenderness <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Consult IHP and EAP/ECP ▪ Reassess vital signs ▪ Apply cold pack to swollen joint ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Minor lacerations, ecchymoses, or abrasions ▪ Bleeding responds to pressure <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Apply firm, direct pressure ▪ Apply cold pack as indicated ▪ Consult IHP and EAP/ECP ▪ Observe student ▪ Contact parent/guardian ▪ Return to class when bleeding stops or send home as indicated ▪ Educate student, parent/guardian, and school personnel about playground/sport safety ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Hemorrhage Control

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*); and perform interventions **AS YOU GO**.

Controlling the bleeding is the first priority!

KEY ASSESSMENT POINTS FOR HEMORRHAGE CONTROL

- Scene Safety!
- Assess for uncontrolled bleeding
- Activate EMS
- Control hemorrhage and then assess C-ABCDE

IMMEDIATE INTERVENTIONS

- Elevate limb while applying direct pressure to hemorrhage site with gloved hand or fingers
 - Apply a tourniquet when these measures fail

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

<p style="text-align: center;">EMERGENT</p> <ul style="list-style-type: none"> ▪ Bleeding requiring a tourniquet such as limb amputations, partial limb amputations, or injuries with arterial bleeding ▪ Acute change from baseline mental status/LOC ▪ Irregular pulse/bradycardia ▪ Hypotension/Hypovolemic Shock <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Apply direct pressure ▪ Apply a tourniquet for bleeding not controlled by direct pressure ▪ Activate EMS ▪ Maintain spinal motion restriction as appropriate ▪ Support C-ABCDE ▪ Keep the student warm ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p style="text-align: center;">URGENT</p> <ul style="list-style-type: none"> ▪ Bleeding from any injury controlled by direct pressure ▪ Large lacerations or avulsions ▪ Tachycardia/bradycardia <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Direct pressure with dressing ▪ Support C-ABCDE as indicated ▪ Keep the student warm ▪ Determine need for EMS ▪ Contact parent/guardian to transport student to medical care or home ▪ Closely observe student ▪ Follow-up 	<p style="text-align: center;">NONURGENT</p> <ul style="list-style-type: none"> ▪ Minor abrasions ▪ Superficial lacerations ▪ No associated injuries <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Perform wound care ▪ Apply dressing ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy
---	---	---

Note: If the incident poses a safety risk (e.g., mass-casualty incident or active shooter), follow lockdown policies or other emergency procedures for any internal or external communication.

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Hemorrhage Control, continued

TOURNIQUET APPLICATION

Indications

- Significant hemorrhage in a limb, that cannot be controlled by direct pressure and elevation

Contraindications

- None when used in the mitigation of a life threatening hemorrhage

Equipment needed

- Appropriate PPE
- Approved arterial tourniquet

Procedure (for commercial tourniquet device)

1. Follow standard precautions to prevent exposure to body fluids
2. Wrap band of tourniquet around the limb, pass the band through the buckle or other fastener device (depending on the model tourniquet) and position tourniquet 2 to 3 inches above the bleeding site. Be sure to apply tourniquet directly to skin.
3. Pull the band tightly and fasten it back on itself all the way around the limb. Be sure to obtain good Velcro contact and do not pass band over the rod clips. Note: the band should be tight enough so that the tips of three fingers cannot be passed between band and skin.
4. Twist rod until bleeding has stopped.
5. Place rod between clips and check bleeding and distal pulse
6. If bleeding is still active, consider further tightening of tourniquet or the application of a second tourniquet above the first.
7. Secure rod and band with time strap and document the time of application directly on the tourniquet.

Reference

North American Rescue (<https://www.narescue.com/pre-hospital-care-ems-products/massive-hemorrhage/combat-application-tourniquet-c-a-t>)

If a commercial tourniquet is unavailable

1. Create a 2–4 inch wide band of soft cloth. A belt, scarf, blood pressure cuff, or triangle bandage may be utilized. Avoid using items that are less than 2 inches wide (such as shoe laces) to prevent further tissue damage. The length required will depend on the size of the student and girth of the extremity.
2. Apply the band around the extremity 2 inches above the wound or amputation site, and tie a half knot (a half knot is the same as the first step in tying a shoe lace).
3. Place a rigid stick or rod on top of the half knot (to create a windlass), and then use the ends of the band to tie a full knot over the stick. This will tie the band to the stick, holding it in place.
4. Rotate/twist the stick until the tourniquet is tight around the limb and/or the bleeding stops.
5. Secure the stick in place so that the tourniquet does not unwind.
6. You may still have a distal pulse so check for signs of circulation after the tourniquet has been applied. The tourniquet must be tightened enough to stop the bleeding.
7. Frequently recheck the wound or stump for signs of bleeding and tighten the tourniquet as needed.
8. Document the time of tourniquet application directly on the tourniquet.

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Hemorrhage Control, continued



No matter how rapid the arrival of professional emergency responders, bystanders will always be first on the scene. A person who is bleeding can die from blood loss within five minutes, therefore it is important to quickly stop the blood loss.

"Stop the Bleed" is a nationwide campaign to empower individuals to act quickly and save lives.

*Remember to be aware of your surroundings and move yourself and the injured person to safety, if necessary.

Call 911.

Bystanders can take simple steps to keep the injured alive until appropriate medical care is available. Here are three actions that you can take to help save a life:

COMPRESS

Expose to find where the bleeding is coming from and apply **firm, steady, pressure** to the bleeding site with bandages or clothing



TOURNIQUET

If the bleeding doesn't stop, **Place** a tourniquet 2-3 inches closer to the torso from the bleeding. (The tourniquet may be applied & secured over clothing).

Pull the strap through the buckle, **Twist** the rod tightly, **Clip** and secure the rod with the clasp or the Velcro strap.

* One type of tourniquet is depicted.

COMPRESS AGAIN


If the bleeding still doesn't stop, **Place** a second tourniquet closer to the torso from first tourniquet.

Pull the strap through the buckle, **Twist** the rod tightly, **Clip** and secure the rod with the clasp or the Velcro strap


* One type of tourniquet is depicted.



Non-Commercial Tourniquets



DTB1F1306



Source: Integrated Publishing

(<http://www.tpub.com/seabee/6-1.htm>)

Source: U.S. Department of Homeland Security

"Stop the Bleed" campaign

(<https://www.dhs.gov/stopthebleed>)

Increased Intracranial Pressure in a Student with a VP Shunt

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

KEY ASSESSMENT POINTS FOR STUDENTS WITH VP SHUNTS

- Facial symmetry
 - Gag reflex
 - Pupil size/reactivity
- Extraocular eye movements
- Neurological function
 - Glasgow Coma Scale (GCS) score
 - Symmetry of function/strength, posture, gait, balance, and spontaneous movement

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Change in mental status ▪ Lethargy ▪ Acute neurological deficit ▪ Inability to look up/roll eyes upward ▪ New onset eye deviation ▪ Seizure (see <i>Seizures</i> protocol) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Consult IHP and EAP/ECP ▪ Directly/continuously observe student ▪ Contact physician for instructions ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Early S/S of shunt dysfunction: <ul style="list-style-type: none"> • Headache • Irritability • Vomiting • Decreased appetite • Change in personality • Loss of existing skills or abilities • Swelling/erythema along shunt path • Seizures • Loss of balance ▪ S/S of shunt tract infection <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Place student supine ▪ Elevate head ▪ Allow student to rest ▪ Reduce environmental stimuli ▪ Consult IHP and EAP/ECP ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Mild headache without other S/S of shunt dysfunction <p>Normal neurological assessment and GCS score</p> <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Consult IHP and EAP/ECP ▪ Allow student to rest, and then reassess ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Reassess every 2 hours if student remains at school ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Lacerations/Abrasions

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR LACERATIONS AND ABRASIONS

- Inspection of wound
- Neurovascular assessment distal to injury:
 - 6 Ps mnemonic: Pain, pulselessness, pallor, paresthesia, paralysis, and poikilothermia
 - Capillary refill
 - Edema
 - Skin temperature

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Absent distal pulses ▪ Significant blood loss ▪ Crush injury ▪ Amputation (see <i>Trauma</i> protocol) ▪ Penetrating wound ▪ Capillary refill exceeds 2 sec ▪ Altered mental status ▪ S/S of respiratory distress <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Control hemorrhage ▪ Activate EMS ▪ Support C-ABCDE ▪ Elevate/immobilize extremity ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Stable vital signs ▪ Pulses are present distal to injury ▪ Significantly contaminated lacerations ▪ Facial lacerations ▪ Puncture wounds of foot ▪ Wounds requiring sutures ▪ Controllable bleeding <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Control bleeding with direct pressure ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Stable vital signs ▪ Superficial abrasion, scrape, or wound ▪ Small splinter or foreign body <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Remove splinter or foreign body ▪ Cleanse wounds using aseptic technique (see below) ▪ Bandage wounds ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

General wound care/aseptic cleansing

- Clean wounds thoroughly with soap
- Rub abrasions gently with 4x4 gauze to remove debris and crusts
- Rinse copiously with water
- Bandage abrasions loosely using non-adherent gauze to allow air circulation
- Apply butterfly bandage to lacerations after bleeding has been controlled

Due to high risk of infection, all deep puncture wounds of the foot must be referred to a physician.

How to control bleeding

Controlling bleeding is a priority that initially involves direct pressure. Activate EMS for severe injuries that result in hemorrhage.

- Apply dressing and direct manual pressure to any hemorrhage
- Elevate area if possible
- Do not remove dressing; reinforce with additional gauze if needed
- Apply tourniquet or hemostatic dressing for hemorrhage not controlled by the above measures, if available and allowed by policy/protocol (see *Hemorrhage Control* protocol for further information).

Note

Refer student for tetanus booster if it has been 5 years or more since the last vaccination. Tetanus booster is recommended every 10 years.

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Musculoskeletal Injuries

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*; also see *Interventions for Musculoskeletal Injuries* protocol on next page) and perform interventions AS YOU GO.

Provide spinal motion restriction if head/spinal injury is suspected!

KEY ASSESSMENT POINTS FOR MUSCULOSKELETAL INJURIES

Focused physical examination of affected area:

- Bilateral symmetry
- Range of motion (ROM)/strength
- Visual inspection for abnormalities

Neurovascular status distal to injury:

- Pain, pulselessness, pallor, paresthesia, paralysis, and poikilothermia (6 Ps)
- Edema
- Capillary refill time
- Skin temperature

IMMEDIATE INTERVENTIONS

Even before you determine triage category, immobilize and support the affected area proximal and distal to the injury!

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Open fracture ▪ Amputation (see <i>Trauma and Hemorrhage Control</i> protocols) ▪ Degloving injury ▪ Neurovascular compromise ▪ Severe edema/deformity at joint or extremity ▪ Joint deviation ▪ Suspected femoral fracture <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Instruct student to avoid weight-bearing/movement of injured area ▪ Immobilize and position suspected fractures/dislocations (see next page) ▪ Reassess neurovascular status distal to injury every 5–10 min ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Stable vital signs ▪ Moderate deformity/discoloration without open wound ▪ Moderate edema at joint or extremity ▪ Moderate pain/guarding ▪ Normal neurovascular findings <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Immobilize and position suspected fracture/dislocation (see next page) ▪ Elevate extremity ▪ Apply cold packs ▪ Observe student closely ▪ Reassess neurovascular status distal to injury every 5–10 min ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Normal vital signs ▪ No deformity ▪ Mild soft tissue edema ▪ Mild pain/point tenderness ▪ Able to bear weight ▪ Normal neurovascular findings <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Apply cold pack ▪ Elevate area ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Interventions for Musculoskeletal Injuries

MANUAL CERVICAL SPINE MOTION RESTRICTION

- Position student supine
- Place both hands along lateral aspect of student's head
- Position the head so that the neck is in neutral alignment with the spine
- Continue to support the head to maintain neutral cervical alignment



Correct method of simultaneous cervical spine motion restriction during airway opening in a child with multiple injuries.

IMMOBILIZATION/POSITIONING OF FRACTURES AND DISLOCATIONS

Area/Injury	Treatment
Upper extremity	<ul style="list-style-type: none"> ▪ Apply sling/triangular bandage ▪ Swathe if additional immobilization is indicated
Clavicular injury/shoulder dislocation	<ul style="list-style-type: none"> ▪ Apply sling/triangular bandage ▪ Swathe if additional immobilization is indicated
Angulation with unimpaired circulation	<ul style="list-style-type: none"> ▪ Immobilize as presented ▪ Do not move extremity
Angulation with absent distal pulse and/or cyanosis	<ul style="list-style-type: none"> ▪ Return extremity to proper physiological position ▪ Apply gentle traction until pulse is restored ▪ Splint or immobilize injured area and joints proximal and distal to injury ▪ Reassess pulses every 5–10 minutes

SLING-AND-SWATHE IMMOBILIZATION




Place the arm across the chest and position as shown. Bring the bandage over the arm and behind the neck.



Adjust the length as necessary and tie the ends. The arm should be well supported and relieve pressure on the shoulder.



Place the knot so that it lies over the shoulder rather than against the cervical spine. Placing a pad under the knot will enhance comfort.



Secure the sling at the elbow with a safety pin or knot to create a pocket in which the elbow rests securely. Reassess neurovascular integrity.



If further immobilization is needed to secure the extremity and a second bandage is available, swathe the arm as permitted by applicable protocols. Lay the second bandage flat, and then fold it several times lengthwise. Use the folded bandage to swathe the injured arm against the chest wall, which immobilizes the injured extremity.

Nose Emergencies

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

Provide spinal motion restriction if head/spinal injury is suspected!

KEY ASSESSMENT POINTS FOR EMERGENCIES INVOLVING THE NOSE

- Mechanism of injury/events preceding episode
- Airway status
- Physical examination/inspection for abnormalities, bleeding, and drainage

IMMEDIATE INTERVENTIONS

Even before you determine triage category, begin to control bleeding with pressure.

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<p>▪ Suspected nasal fracture with potential head/neck injury (see <i>Head/Spinal Cord</i> protocol)</p> <p>▪ Change in mental status</p> <p>▪ Airway compromise</p> <p>▪ Cerebrospinal fluid (CSF) drainage</p> <p>INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Maintain spinal motion restriction as applicable ▪ Support C-ABCDE ▪ Directly/continuously observe student ▪ Keep student calm ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p>▪ Suspected nasal fracture, but no possibility of head/neck injury</p> <p>▪ Tenderness on palpation</p> <p>▪ Epistaxis not controlled after 10 min</p> <p>▪ Moderate periorbital edema</p> <p>INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS <p>Prolonged epistaxis</p> <ul style="list-style-type: none"> ▪ Have student blow their nose to remove clots ▪ Pinch nostrils closed and apply pressure for 10 min while student leans forward ▪ If epistaxis continues, consider activating EMS <p>Suspected nasal fracture</p> <ul style="list-style-type: none"> ▪ Apply cold packs ▪ See <i>Lacerations/Abrasions</i> protocol for treatment of associated wounds <p>In all cases</p> <ul style="list-style-type: none"> ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<p>▪ Foreign body</p> <p>▪ Controllable epistaxis</p> <p>▪ S/S of acute sinusitis:</p> <ul style="list-style-type: none"> • Pain/pressure over sinus areas • Throbbing • Headache, malaise, fever • Mucopurulent secretions • Mild periorbital edema <p>INTERVENTIONS</p> <p>Epistaxis</p> <ul style="list-style-type: none"> ▪ Have student blow their nose to remove clots ▪ Pinch nostrils closed and apply pressure for 10 min <p>Foreign body</p> <ul style="list-style-type: none"> ▪ Have student blow nose while occluding unobstructed nostril ▪ Attempt removal only if object is visible and can be grasped with forceps or fingers ▪ During extraction, occlude nostril superior to object so that it cannot be pushed further in ▪ If object cannot be removed, reclassify as urgent <p>In all cases</p> <ul style="list-style-type: none"> ▪ Observe student ▪ Contact parent/guardian for referral to primary care physician ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Obstetric Emergencies

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*; also see *Emergency Neonatal Care* protocol on next page) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR OBSTETRIC EMERGENCIES

- Obstetric history:**
- Anticipated due date
 - Recent drug use
 - Possibility of multiple births
 - Prenatal care (name/phone number of obstetrician)
 - Delivery hospital
 - Color of amniotic fluid (clear, red, green, or yellow-tinged)
 - Progression of labor
 - Bloody show/expulsion of mucous plug
 - Timing/strength of contractions
 - Inspection for crowning
 - Reported urge to move bowels

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ S/S of shock ▪ Seizure activity (see <i>Seizures</i> protocol) ▪ S/S of preeclampsia: <ul style="list-style-type: none"> • BP equals or exceeds 140/90 or • BP elevated by 30 mm Hg (systolic) or 15 mm Hg (diastolic) above known baseline ▪ Crowning ▪ Breech presentation ▪ Prolapsed umbilical cord ▪ Abruptio placentae ▪ Placenta previa/vaginal bleeding ▪ Multigravida ▪ Premature labor ▪ Contractions less than 10 min apart <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ For signs of shock, if delivery is not imminent, place in left lateral recovery position ▪ If delivery is imminent, prepare for emergency delivery (see next page for emergency neonatal care) ▪ Directly/continuously observe student ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Pregnancy-induced hypertension ▪ History of trauma ▪ Active labor, but amniotic sac intact ▪ Contractions more than 10 min apart <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ Observe student closely ▪ Contact parent/guardian to transport student for medical care ▪ Monitor closely ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Variable contractions ▪ Amniotic sac intact ▪ Vomiting with stable vital signs <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Observe student ▪ Contact parent/guardian to transport student for medical care ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Emergency Neonatal Care

IMMEDIATE INTERVENTIONS

- Suction the infant’s mouth first, then the nose
- Dry the infant with a towel
- If the neonate does not need resuscitation, delay clamping the umbilical cord for 30 seconds
- Wrap the infant in a clean, warmed, dry towel or blanket; cover head
- Stimulate breathing by rubbing infant’s back or flicking feet
- Resuscitate if necessary according to current AHA guidelines
- Calculate APGAR score (see below)

NOTE:

Do not interrupt resuscitation procedures to calculate APGAR score!

- Directly/continuously observe student and infant
- Contact significant others per student’s request

APGAR SCORING CRITERIA

Unless resuscitation measures are needed, assess the newborn’s APGAR score 1 minute after birth and again 5 minutes after birth. The following Table summarizes APGAR categories and scoring.

APGAR EVALUATION OF THE NEONATE

	Sign	0	1	2
A	Skin color	Blue	Pink, blue extremities	All pink
P	Heart rate	Absent	<100 bpm	>100 bpm
G	Reflex irritability ^a	No response	Weak cry and grimace	Vigorous cry
A	Muscle tone	Flaccid, limp	Some flexion	Active motion
R	Respiratory effort	Absent	Slow, irregular	Good vigorous cry

^a In response to nasal or oral stimulation; bpm, beats per minute

Respiratory Distress

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR RESPIRATORY DISTRESS

- Breathing assessment

IMMEDIATE INTERVENTIONS

Even before you determine triage category, perform the following actions as indicated:

- Loosen restrictive clothing
- Help student into position of comfort
- Maintain airway patency

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ S/S of severe respiratory distress or failure ▪ Apnea or dyspnea ▪ Grunting or drooling ▪ S/S of impending respiratory failure <ul style="list-style-type: none"> • Cyanosis • Tachycardia • Shallow respiration • Altered mental status/restlessness • Hypotension ▪ S/S of airway obstruction (see <i>Foreign Body</i> protocol) ▪ Severe asthma episode (see <i>Asthma</i> protocol) ▪ S/S of epiglottitis; dysphagia, drooling, high fever, stridor, or tripod positioning 	<ul style="list-style-type: none"> ▪ S/S of moderate respiratory distress ▪ Fever, chills ▪ Persistent or barking cough ▪ Stridor or wheezing ▪ Nasal flaring ▪ Retractions ▪ Pleural pain ▪ Mild to moderate asthma (see <i>Asthma Episode</i> protocol) 	<ul style="list-style-type: none"> ▪ Hyperventilation ▪ S/S of mild upper respiratory infection (URI) <ul style="list-style-type: none"> • Cough • Nasal congestion • Sore throat • Hoarseness • Low-grade fever
<p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Maintain position of comfort ▪ Administer high-flow O₂ as tolerated if available <p style="text-align: center;">Epiglottitis</p> <ul style="list-style-type: none"> ▪ Do not inspect hypopharynx ▪ Keep student calm <p style="text-align: center;">Respiratory failure</p> <ul style="list-style-type: none"> ▪ Assist ventilation via mouth-to-mask ▪ Anticipate need for CPR <p style="text-align: center;">In all cases</p> <ul style="list-style-type: none"> ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Maintain position of comfort ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<p style="text-align: center;">INTERVENTIONS</p> <p>For hyperventilation</p> <ul style="list-style-type: none"> ▪ Encourage student to relax <p style="text-align: center;">In all cases</p> <ul style="list-style-type: none"> ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated (send home for fever with temperature exceeding 100°F/37.8°C or per policy) ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and on authorized policies.

Seizures

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

- Provide spinal motion restriction if head/spinal injury is suspected!
 - Open airway as necessary using the jaw-thrust maneuver.

KEY ASSESSMENT POINTS FOR SEIZURES

- Past health history; particularly note
 - Epilepsy
 - Syncope
 - Diabetes
- Events preceding episode, particularly head injury
- Indicators of drug overdose, meningitis, or hypoglycemia (see *Substance Abuse and Diabetic Emergencies* protocols)

IMMEDIATE INTERVENTIONS

During an active seizure, perform the following actions *before* you proceed with triage:

- Do not put anything in student’s mouth—do not restrict movement in any way
- Provide privacy
- Protect student from injury

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ First-time seizure/no known history of seizures ▪ History of seizures and medication noncompliance with no recent seizures ▪ Seizure/series of seizures that persist for more than 5 min ▪ Associated respiratory compromise ▪ Associated head injury or trauma <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Consult IHP and EAP/ECP ▪ Directly/continuously observe student ▪ Provide psychological support ▪ Document time, characteristics, and duration of seizure ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Atypical seizure in student with history of seizures <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Consult IHP and EAP/ECP ▪ Check glucose if possible ▪ Allow to rest in left lateral recovery position ▪ Provide psychological support ▪ Observe student closely ▪ Document characteristics and duration of seizure ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Typical seizure in student with baseline history of frequent seizures <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Consult IHP and EAP/ECP ▪ Check blood glucose if possible ▪ Allow to rest in left lateral recovery position during postictal phase ▪ Provide psychological support ▪ Observe student ▪ Document characteristics and duration of seizure ▪ Contact parent/guardian as per policy ▪ Return student to class or send home as indicated ▪ For persistent drowsiness, notify parent/guardian to transport student home ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student’s clinical presentation and authorized policies.

Sexual Abuse/Assault and Teen Dating Violence

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR SEXUAL ABUSE OR ASSAULT

- Psychosocial history
- Menstrual status/possibility of pregnancy
- Long-term sexual abuse must be reported to appropriate authorities but may not require EMS transport
- Focused physical examination for acute injuries
- Limit physical contact if assault just occurred to preserve forensic evidence
- Encourage student to bring all clothing worn during the assault to the ED

IMMEDIATE INTERVENTIONS

Even before you determine triage category, perform the following actions as indicated:

- Treat any injuries as indicated
- Provide a safe, nonthreatening environment
- Ask open-ended questions
- Support student to seek professional help

Note

Any case of suspected sexual assault requires ED treatment and local law enforcement notification. The student should disclose the full account of the assault ONLY to a Sexual Assault Nurse Examiner RN or physician performing the forensic exam. Only specially trained professionals should interview victims of sexual assault. Advocacy centers are located throughout Illinois. These centers conduct interviews and provide a variety of victim services; some offer forensic exams.

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent! In all cases, report suspicions to DCFS at 800-25-ABUSE (22873) and/or local law enforcement (see Appendix B for a report form). For students ≥13 years, consent to inform parent must be granted by the student.

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Thoughts of death or suicide ▪ Severe/life-threatening injuries (see <i>Trauma</i> protocol) ▪ Suspected sexual assault <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Directly/continuously monitor student ▪ Refer to school counselor as appropriate ▪ Contact parent/guardian as per policy ▪ Preserve evidence and send to ED with student ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Ecchymoses/injuries, not life-threatening ▪ Alcohol/drug use ▪ Current, previous, or potential pregnancy <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Observe student closely ▪ Provide support ▪ Refer to school counselor ▪ Contact parent/guardian to transport student to medical care or home as per policy ▪ Follow-up 	<ul style="list-style-type: none"> ▪ History of truancy ▪ Sudden change in dress or makeup ▪ Difficulty making decisions ▪ Abrupt changes in mood or personality ▪ Combative, possessive, or jealous behavior ▪ Withdrawal/self-isolation <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Refer to school counselor ▪ Observe student's behavior with others ▪ Document findings ▪ Provide support ▪ Contact parent/guardian as per policy ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Sickle Cell Disease

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

KEY ASSESSMENT POINTS FOR SICKLE CELL DISEASE

- Past health history including recent illness
 - Events leading up to episode, including exposure to temperature extremes
- Focused physical examination/palpation for localized pain, edema, or other abnormalities
- Last food/drink (likelihood of dehydration)

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Fever (T exceeds 100°F/37.8°C) ▪ S/S of infection or sepsis ▪ Severe, continuous pain in extremities, back, chest, or abdomen ▪ Seizure activity (see <i>Seizures</i> protocol) ▪ Change in mental status ▪ Dyspnea ▪ Chest pain ▪ Priapism/penile pain ▪ Severe splenomegaly and S/S of shock ▪ Pallor or lethargy with other abnormal findings ▪ S/S of impending respiratory collapse: <ul style="list-style-type: none"> · Cyanosis · Tachycardia · Shallow respiration · Altered mental status/restlessness · Hypotension · Decreased breath sounds ▪ Respiratory distress (acute chest syndrome) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Refer to IHP and EAP/ECP <li style="padding-left: 20px;">Respiratory distress/shock ▪ Administer high-flow O₂ if available ▪ Help to maintain position of comfort (shock position as indicated) <li style="padding-left: 20px;">In all cases ▪ Directly/continuously observe student ▪ Reassess vital signs every 5 min ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Fever (T exceeds 100°F/37.8°C) ▪ Mild to moderate pain ▪ Severe edema/tenderness of affected areas <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Refer to IHP and EAP/ECP ▪ Provide oral hydration (4–8 oz/hour) ▪ Allow to rest ▪ Elevate affected extremity ▪ Observe student closely ▪ Contact parent/guardian to transport student to medical care or home ▪ Educate student about need for good hydration and protection from temperature extremes ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Minor localized pain <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Refer to IHP and EAP/ECP ▪ Provide oral hydration (4–8 oz/hour) ▪ Allow to rest ▪ Reassess pain ▪ Contact parent/guardian ▪ Observe student ▪ Return student to class or send home as indicated ▪ Reassess every 2 hours if student remains at school ▪ Educate student about need for good hydration and protection from temperature extremes ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Substance Abuse

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR SUBSTANCE ABUSE

- Respiratory assessment
- Orientation to person/place/time
- Ability to recall event/injuries incurred
- Type/amount of substance involved (request medication/drug container if available)
- Time/route of exposure (dermal, ocular, inhalation, or ingestion)
- Underlying health problems (**SAMPLE** history)
- Other students involved

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Respiratory/cardiac arrest ▪ Unconscious/cannot be roused ▪ Change in mental status ▪ Labored/shallow breathing ▪ Hallucinations, violent behavior, or extreme agitation ▪ S/S of trauma or injury ▪ Seizure activity (see <i>Seizures</i> protocol) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Initiate CPR as appropriate ▪ Place student in left lateral recovery position ▪ Directly/continuously observe student ▪ Monitor airway/respiratory status ▪ See <i>Trauma</i> protocol as appropriate ▪ Contact the Poison Center as indicated^a ▪ Contact parent/guardian ▪ Notify school administrator ▪ Initiate counseling/support measures per school policy ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Somnolence, emesis, or unusual behavior ▪ Unsteady gait ▪ Memory problems ▪ Mild agitation/restlessness ▪ Suspicious odors ▪ Needle marks/drug residue on skin, nose, or clothes ▪ Previous loss of consciousness ▪ Complicating health problems <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ Observe student continuously ▪ Give nothing by mouth ▪ Remain with student ▪ See <i>Trauma</i> protocol as appropriate ▪ Contact the Poison Center as indicated^a ▪ Contact parent/guardian to transport student to medical care or home ▪ Notify school administrator ▪ Initiate counseling/support measures per school policy ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Coherent and oriented ▪ Stable vital signs ▪ No loss of consciousness <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Observe student ▪ Remain with student ▪ Contact the Poison Center as indicated^a ▪ Notify school administrator ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Initiate counseling/support measures per school policy ▪ Follow-up as needed or per policy

Note

^aWhen calling the Poison Center (800-222-1222), identify yourself as a healthcare professional and provide the following information: your name and phone number; student's name, age, weight, and vital signs; substance involved (if known); amount, time, route, and duration of exposure; abnormal signs and symptoms (S/S); first aid and immediate interventions rendered.

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Suicide Prevention

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

KEY ASSESSMENT POINTS FOR POTENTIAL SUICIDE

Focused Psychosocial Examination/Risk Factor Assessment

Precipitating events

- Parents recently divorced
 - Multiple life stressors
- Recent breakup with girlfriend or boyfriend
 - Unplanned pregnancy

Preparatory actions

- Acquiring the means
- Putting affairs in order
 - Suicide talk
- Giving away prized possessions
- Precautions against discovery

Current Symptoms

- Hopelessness/powerlessness
- Depressed mood
- Suicidal ideation
- Abrupt change in personality

History

- Previous suicide attempts
- Affective disorders or conduct disorder
- Family history of mental illness, suicidal behavior, or affective disorders
- Alcoholism or substance abuse
- Chronic health condition

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

EMERGENT

- Thoughts of death
- Suicide plan and preparations
- Suicide attempt
- Previous suicide attempt

INTERVENTIONS

- Activate EMS
- Support C-ABCDE as indicated
- Remove personal effects
- **Do not leave student alone under any circumstances!**
- Listen to student carefully
- Take conversation seriously
- Notify crisis response team
- Contact parent/guardian
- Notify school administrator
- Follow-up

URGENT

- Suicidal ideation
- Depression
- Withdrawal
- Self-blame
- Self-reproach

INTERVENTIONS

- Determine need for EMS
- **Remain with student at all times**
- Listen to student
- Notify crisis response team
- Contact parent/guardian to transport student to medical care or home
- Follow-up

NONURGENT

- Frequent physical complaints
- Sad affect

INTERVENTIONS

- Observe student regularly
- Refer to school counselor
- Notify crisis response team of your concerns and findings
- Contact parent/guardian
- Return student to class or send home as indicated
- Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Syncope/Unconsciousness

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

Provide spinal motion restriction if head/spinal injury is suspected!

KEY ASSESSMENT POINTS FOR SYNCOPE

- Mental status/neurological assessment
 - Psychosocial history
- Past health history/current menstrual status
- Medications taken
- Events leading up to episode, including activities and weather conditions
- Last food/drink taken

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Acute change from baseline mental status ▪ Irregular pulse ▪ Acute neurological deficit ▪ Head injury/headache with altered mental status or vomiting ▪ Severe headache with altered mental status ▪ Head injury/history of anemia, hemophilia, or other coagulopathy ▪ Associated seizure activity (see <i>Seizures</i> protocol) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Provide spinal motion restriction as applicable ▪ Support C-ABCDE ▪ Directly/continuously observe student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Possible ventriculoperitoneal (VP) shunt dysfunction ▪ S/S of moderate hypoglycemia ▪ Signs of dehydration ▪ Severe headache without altered mental status ▪ Persistent or severe dizziness ▪ Exercise-induced syncope (possible cardiac etiology) ▪ Possible medication reaction without emergent findings ▪ History of substance abuse or eating disorder <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE as indicated ▪ Determine need for EMS ▪ Contact parent/guardian to transport student to medical care or home ▪ Closely observe student ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Hyperventilation ▪ Vasovagal reaction to anxiety/pain or other known trigger ▪ Exposure to ambient heat ▪ Evidence of carotid sinus reaction (e.g., subsequent to neck hold by classmate) ▪ No associated injuries <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Place student supine ▪ Allow student to wake spontaneously ▪ For hyperventilation, encourage student to relax ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

See the following protocols as appropriate:

- *Diabetic Emergencies*
- *Head/Spinal Cord Trauma*
- *Heat-Related Injuries*
- *Hemophilia*
- *Increased intracranial pressure (ICP) in a Student With a VP Shunt*
- *Seizures*
- *Trauma*

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Throat Emergencies

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*) and perform interventions **AS YOU GO**.

KEY ASSESSMENT POINTS FOR THROAT EMERGENCIES

- Across-the-room assessment (note tripod positioning)
 - Respiratory assessment
- Events preceding illness/suddenness of onset
- Focused physical examination with inspection of pharynx and palpation of lymph nodes

IMMEDIATE INTERVENTIONS

Even *before* you determine triage category, perform the following actions as indicated:

- Loosen restrictive clothing
- Help student into position of comfort
- Maintain airway patency

DETERMINE TRIAGE CATEGORY AND ADDITIONAL INTERVENTIONS

Determine triage category and activate EMS **AS SOON AS** the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Airway compromise ▪ Change in mental status ▪ S/S of epiglottitis <ul style="list-style-type: none"> • Sudden onset • Stridor, drooling, or dysphagia • High fever ▪ Anaphylactic reaction (see <i>Anaphylaxis</i> protocol) ▪ S/S of retropharyngeal abscess <ul style="list-style-type: none"> • Fever • Stiff, painful neck • Asymmetrical edema of posterior pharyngeal wall • Dyspnea <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Keep student calm ▪ Administer high-flow O₂ as tolerated if available <p style="text-align: center;">Epiglottitis</p> <ul style="list-style-type: none"> ▪ Do not inspect hypopharynx ▪ Maintain position of comfort <p style="text-align: center;">In all cases</p> <ul style="list-style-type: none"> ▪ Directly/continuously observe student ▪ Reassess vital signs every 5 min ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ S/S of peritonsillar abscess <ul style="list-style-type: none"> • Severe pain • Fever • Muffled voice (e.g., “hot potato voice”) <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Determine need for EMS ▪ Maintain position of comfort ▪ Contact parent/guardian to transport student to medical care or home ▪ Observe student closely ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Swollen, tender lymph nodes ▪ S/S of tonsillitis/pharyngitis <ul style="list-style-type: none"> • Tonsillar exudate • Erythema • Deviation of tonsils toward midline <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Observe student ▪ Contact parent/guardian ▪ Return student to class or send home as indicated (send home for fever exceeding 100°F/37.8°C) ▪ Follow-up as needed or per policy

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student’s clinical presentation and authorized policies.

Toxic Exposure (Ingestion/Environmental)

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*; also see *Toxidromes* on next page) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR TOXIC EXPOSURES

- Assess/ensure scene safety (notify school administrator as indicated to activate hazmat protocols)
 - Respiratory assessment
 - Skin assessment
- LOC/neurological assessment, including PERRL
 - Type/amount of substance (get container if available)
 - Time/route of exposure (dermal, ocular, inhalation, or ingestion)
- Location where exposure occurred
- Subsequent S/S; respiratory status, LOC, and emesis
- Ability to recall event, including injuries incurred
- Underlying health problems (SAMPLE history)
- Focused physical examination, including injury and/or odors
- Other students involved

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

Refer to Toxidromes table (see next page)

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Respiratory arrest ▪ Seizures (see <i>Seizures</i> protocol) ▪ Loss of consciousness ▪ Dyspnea or severe respiratory distress ▪ Signs of shock/hypotension <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Activate EMS ▪ Support C-ABCDE ▪ Initiate CPR as necessary ▪ Contact the Poison Center^a ▪ Refer for medical care as recommended by the Poison Center ▪ Send safety data sheet (SDS) and substance (if possible) to ED with student ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Unusual behavior ▪ History of emesis ▪ Minor abnormal findings <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS ▪ Contact the Poison Center^a ▪ Refer for medical care as recommended by the Poison Center ▪ Consult SDS ▪ Contact parent/guardian to transport student to medical care or home ▪ Notify school administrator ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Asymptomatic ▪ Stable vital signs <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Observe student ▪ Consult SDS ▪ Contact the Poison Center^a ▪ Refer for medical care as recommended by the Poison Center ▪ Provide supportive care as indicated ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Notify school administrator ▪ Follow-up as needed or per policy

NOTE

^a**When calling the Poison Center (800-222-1222), identify yourself as a healthcare professional and provide the following information: your name and phone number; student's name, age, weight, and vital signs; substance involved (if known); amount, time, route, and duration of exposure; abnormal signs and symptoms (S/S); first aid and immediate interventions rendered.**

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Toxidromes

If assessment findings suggest a toxic ingestion and you do not know what substance is involved, it may be helpful to look for signs of identifiable toxic syndromes (toxidromes). Toxidromes involve a recognizable group of signs and symptoms that tend to occur consistently with particular toxins. The following Table describes the four major toxidromes and provides mnemonics that can help you remember the associated findings. Examples of substances that can cause each toxidrome are included.

TOXIDROMES

Toxidrome	Clinical Findings	Causative Substances
Opioid	<ul style="list-style-type: none"> ▪ Constricted pupils ▪ CNS depression ▪ Respiratory depression 	<ul style="list-style-type: none"> ▪ Heroin ▪ Codeine ▪ Fentanyl ▪ Methadone
Sympathomimetic	<ul style="list-style-type: none"> ▪ Hypertension ▪ Tachycardia ▪ Hyperthermia ▪ Diaphoresis ▪ Dilated pupils 	<ul style="list-style-type: none"> ▪ Epinephrine ▪ OTC diet aids containing caffeine ▪ Amphetamines ▪ Oral decongestants (e.g., pseudoephedrine) ▪ Bronchodilators
Anticholinergic <i>Listed phrases may help you recall clinical findings</i>	<ul style="list-style-type: none"> ▪ Hyperthermia (<i>hot as a hare</i>) ▪ Flushed skin (<i>red as a beet</i>) ▪ Hypertension, dry skin (<i>dry as a bone</i>) ▪ Delirium (<i>mad as a hatter</i>) ▪ Dilated pupils (<i>blind as a bat</i>) ▪ Urinary retention (<i>full as a flask</i>) ▪ Tachycardia ▪ Absent bowel sounds 	<ul style="list-style-type: none"> ▪ Antihistamines ▪ GI antispasmodics ▪ Certain toxic plants (e.g., jimson weed, deadly nightshade, atropine) ▪ Tricyclic antidepressants
Cholinergic	<p>SLUDGE</p> <ul style="list-style-type: none"> ▪ Salivation ▪ Lacrimination ▪ Urination ▪ Diarrhea ▪ Gastrointestinal distress ▪ Emesis <p><i>Other symptoms</i></p> <ul style="list-style-type: none"> ▪ Miosis, muscle fasciculations ▪ Bradycardia, bronchorrhea ▪ Weakness 	<ul style="list-style-type: none"> ▪ Organophosphate and carbamate insecticides ▪ Physostigmine

CNS, central nervous system; OTC, over-the-counter; GI, gastrointestinal

Trauma

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*; also see *Interventions for Musculoskeletal Trauma* protocol on next page) and perform interventions AS YOU GO. Ensure scene safety before approaching. Provide spinal motion restriction if head/spinal injury is suspected!

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

EMERGENT	URGENT	NONURGENT
<ul style="list-style-type: none"> ▪ Profuse hemorrhage ▪ Slow or fast RR, other S/S of respiratory distress/failure ▪ Capillary refill exceeds 2 seconds, other S/S of shock ▪ Currently/previously unconscious ▪ Penetrating wound or significant blunt trauma to head, chest, or abdomen ▪ Open chest wound ▪ Suspected pelvic/femoral fracture ▪ Suspected spinal injury with paresthesia ▪ Amputation/crush injury ▪ S/S of intra-abdominal injury: <ul style="list-style-type: none"> • Hematuria • Grey Turner sign (bluish discoloration of flank/periumbilical area) • Abdominal asymmetry/distention • Tenderness/guarding/pain on gentle palpation <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Control hemorrhage ▪ Activate EMS ▪ Support C-ABCDE ▪ Maintain spinal motion restriction ▪ Place supine or in shock position ▪ Keep student warm ▪ Give nothing by mouth ▪ Stabilize impaled object with dressings—do not remove ▪ Splint/elevate suspected fracture ▪ Control hemorrhage ▪ Directly/continuously observe student ▪ Frequently reassess vital signs/AVPU ▪ Contact parent/guardian ▪ Notify school administrator ▪ Follow-up <p style="text-align: center;">Amputation management</p> <ul style="list-style-type: none"> ▪ Wrap part in gauze slightly moistened with sterile saline ▪ Place in plastic bag, seal securely, and label ▪ Place bag on cold packs (not ice) for transport ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Stable vital signs ▪ No loss of consciousness ▪ Deformity/suspected closed fracture without neurovascular compromise ▪ Controllable bleeding ▪ Significant mechanism of injury without other significant abnormalities <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Support C-ABCDE ▪ Reassess vital signs ▪ Determine need for EMS ▪ Control bleeding ▪ Proceed with detailed physical examination ▪ Splint/immobilize suspected fractures, elevate extremity, and apply cold packs <p style="text-align: center;">Blunt abdominal injury</p> <ul style="list-style-type: none"> ▪ Observe closely for 15 min for S/S of intra-abdominal injury <ul style="list-style-type: none"> • Hematuria • Grey Turner sign (bluish discoloration of flank/periumbilical area) • Abdominal asymmetry/distention • Tenderness/guarding/pain on gentle palpation <p>▪ Reassess in 1 hour or if S/S recur</p> <p style="text-align: center;">In all cases</p> <ul style="list-style-type: none"> ▪ Contact parent/guardian to transport student to medical care or home ▪ Follow-up 	<ul style="list-style-type: none"> ▪ Stable vital signs ▪ No deformity or suspicion of fracture ▪ Minor abrasions or lacerations ▪ Mild muscle strain or sprain <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Proceed with detailed physical examination ▪ Refer to appropriate protocol^a <p style="text-align: center;">Blunt abdominal injury</p> <ul style="list-style-type: none"> ▪ Observe closely for 15 min for S/S of intra-abdominal injury <ul style="list-style-type: none"> • Hematuria • Grey Turner sign (bluish discoloration of flank/periumbilical area) • Abdominal asymmetry/distention • Tenderness/guarding/pain on gentle palpation <p>▪ Reassess in 1 hour or if S/S recur</p> <p style="text-align: center;">In all cases</p> <ul style="list-style-type: none"> ▪ Contact parent/guardian ▪ Return student to class or send home as indicated ▪ Follow-up as needed or per policy

^aSee the following protocols as appropriate: ▪ *Abdominal Pain* ▪ *Burns* ▪ *Chest Trauma* ▪ *Head/Spinal Cord Trauma* ▪ *Hemorrhage Control* ▪ *Lacerations/Abrasions* ▪ *Musculoskeletal Injury*

Refer student for tetanus booster if it has been 5 years or more since the last vaccination. Tetanus booster is recommended every 10 years.

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student's clinical presentation and authorized policies.

Interventions for Musculoskeletal Trauma

MANUAL CERVICAL SPINE MOTION RESTRICTION

- Position student supine
- Place both hands along lateral aspect of student's head
- Position the head so that the neck is in neutral alignment with the spine
- Continue to support the head to maintain neutral cervical alignment



Correct method of simultaneous cervical spine motion restriction during airway opening in a child with multiple injuries.

IMMOBILIZATION/POSITIONING OF FRACTURES AND DISLOCATIONS

Area/Injury	Treatment
Upper extremity	<ul style="list-style-type: none"> ▪ Apply sling/triangular bandage ▪ Swathe if additional immobilization is indicated
Clavicular injury/shoulder dislocation	<ul style="list-style-type: none"> ▪ Apply sling/triangular bandage ▪ Swathe if additional immobilization is indicated
Angulation with unimpaired circulation	<ul style="list-style-type: none"> ▪ Immobilize as presented ▪ Do not move extremity
Angulation with absent distal pulse and/or cyanosis	<ul style="list-style-type: none"> ▪ Activate EMS ▪ Splint or immobilize injured area, including joints that are proximal and distal to the injury ▪ Reassess pulses every 5–10 minutes

SLING-AND-SWATHE IMMOBILIZATION




Place the arm across the chest and position as shown. Bring the bandage over the arm and behind the neck.



Adjust the length as necessary and tie the ends. The arm should be well supported and relieve pressure on the shoulder.



Place the knot so that it lies over the shoulder rather than against the cervical spine. Placing a pad under the knot will enhance comfort.



Secure the sling at the elbow with a safety pin or knot to create a pocket in which the elbow rests securely. Reassess neurovascular integrity.



If further immobilization is needed to secure the extremity and a second bandage is available, swathe the arm as permitted by applicable protocols. Lay the second bandage flat, and then fold it several times lengthwise. Use the folded bandage to swathe the injured arm against the chest wall, which immobilizes the injured area.

Violent Behavior

SYSTEMATIC ASSESSMENT

Begin the four components of assessment (see *Systematic Assessment/Immediate Care and Assessment Tools*; also see *Violent Behavior: Safety Tips* protocol on next page) and perform interventions AS YOU GO.

KEY ASSESSMENT POINTS FOR VIOLENT INCIDENTS

- Assess/ensure scene safety before approaching (notify police liaison/security)
- Perform across-the-room assessment followed by mental status examination (below)
- Remember to use communication techniques that are appropriate for the student’s developmental level and ability

- **Appearance:** General, grooming, posture
- **Mood:** Cooperative, frightened, irritable
 - **Speech:** Soft/loud, fast, slurred
- **Behavior:** Fidgeting, pacing, eye contact
- **Memory:** Recent memory/immediate recall
 - **Orientation:** Realistically oriented to person/place/time
- **Thought process:** Mental activity, evidence of delusions/hallucinations

- **Thought content:** What the student says, suicidal ideation, hopelessness
- **Insight:** Recognizes responsibilities or blames others for problems
- **Judgment:** Decision-making ability (e.g., superficial or impulsive)
- **Perception:** Awareness of self and thoughts (e.g., guilt or indecisiveness)

DETERMINE TRIAGE CATEGORY AND APPROPRIATE INTERVENTIONS

Determine triage category and activate EMS AS SOON AS the need becomes apparent!

<p style="text-align: center;">EMERGENT</p> <ul style="list-style-type: none"> ▪ Danger to self/others ▪ Brandishing weapon ▪ Physical cues indicate escalation ▪ Drug or alcohol intoxication ▪ Physical restraint necessary ▪ History of violence <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Initiate lockdown procedures ▪ Activate EMS/security ▪ Never intervene alone ▪ See next page for safety tips ▪ Contact parent/guardian ▪ Notify school administrator ▪ Monitor behavioral progress ▪ Follow-up 	<p style="text-align: center;">URGENT</p> <ul style="list-style-type: none"> ▪ Moderately agitated but not violent ▪ Expressing verbal anger without physical aggression <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Determine need for EMS/security ▪ Speak in low, measured tones ▪ Explain that you know something is bothering the student and you will help them control their behavior ▪ Walk with the student to diffuse agitation ▪ Repeat/restate what the student says: <ul style="list-style-type: none"> • “You’re feeling angry.” • “I’m concerned for you. I’m going to help you control yourself.” • “What do you need? What do you need to do?” • “When you felt like this before, what helped you?” ▪ Notify crisis response team ▪ Contact parent/guardian to transport student to medical care or home ▪ Monitor behavioral progress ▪ Follow-up 	<p style="text-align: center;">NONURGENT</p> <ul style="list-style-type: none"> ▪ Mildly anxious or frightened ▪ Previously angry but now calm <p style="text-align: center;">INTERVENTIONS</p> <ul style="list-style-type: none"> ▪ Take student to a quiet area with backup support available ▪ Provide reassurance as needed ▪ Speak in low, measured tones ▪ Contact parent/guardian ▪ Allow student to return to class if student is calm, ready, and able to identify ways to deal with feelings; or send home as indicated ▪ Monitor behavioral progress ▪ Follow-up as needed or per policy
---	---	--

The Illinois Emergency Medical Services for Children School Nurse Committee has exercised extreme caution that all information presented is accurate and in accordance with professional standards in effect at the time of publication. The information does not serve as a substitute for the professional advice of a physician/advanced practice nurse, does not dictate an exclusive course of treatment, and should not be construed as excluding other acceptable methods of treatment. It is recommended that care must be based on the student’s clinical presentation and authorized policies.

Violent Behavior: Safety Tips

Emergent situation

- Do not invade student's personal space (stand back at least 5 ft)
- Stand at a 45° angle to student; do not stand directly in front of student
- Maintain an open posture
- Maintain a clear exit route
- Be prepared to move quickly
- Do not make any abrupt moves unless necessary
- Be sure student has no weapons before approaching
- Give student brief, clear, assertive directions before taking any action
- Establish yourself as a concerned professional
- Proceed without hesitation
- Enlist adequate, trained assistance (at least six people, if possible) before attempting physical restraint

Urgent situation

- Speak in low, measured tones
- Explain that you know something is bothering the student and you will help them control their behavior
- Walk with the student to diffuse agitation
- Repeat/restate what the student says:
 - "You're feeling angry."
 - "I'm concerned for you. I'm going to help you control yourself."
 - "What do you need? What do you need to do?"
 - "When you felt like this before, what helped you?"

