



turning knowledge into practice

**Methodology:
U.S. News & World Report
Best Children's Hospitals 2020-21**

Murrey G. Olmsted

Rebecca Powell

Joe Murphy

Denise Bell

Marshica Stanley

Rebekah Torcasso Sanchez

June 23, 2020



To Whom It May Concern:

U.S. News & World Report's "Best Children's Hospitals" study is the sole and exclusive property of U.S. News & World Report, L.P., which owns all rights, including but not limited to copyright, in and to the attached data and material. Any party wishing to cite, reference, publish or otherwise disclose the information contained herein may do so only with the prior written consent of U.S. News. Any U.S. News-approved reference or citation must identify the source as "U.S. News & World Report's Best Children's Hospitals" and must include the following credit line: "Copyright © 2020 U.S. News & World Report, L.P. Data reprinted with permission from U.S. News." For permission to cite or use, contact permissions@usnews.com. For custom reprints, please contact Wright's Media at 877-652-5295 or usnews@wrightsmedia.com.

Executive Summary

Pediatrics has been an element of Best Hospitals ever since 1990, when U.S. News & World Report published the first annual “America’s Best Hospitals” rankings, as they were then called. The initial evaluations, in 12 specialties, comprised short lists of centers that were identified through a survey of physician specialists as providing the best care for the most challenging patients.

For the first time, patients and families had a tool to help narrow their search for a hospital particularly skilled in performing difficult procedures, treating serious conditions and otherwise demonstrating an especially high level of care. While that core mission has not changed, U.S. News broadened its scope in 2015 by adding ratings of some 4,600 hospitals in relatively commonplace procedures and conditions such as heart bypass surgery, knee and hip joint replacement and COPD.

By 1993 hard data had been incorporated into most Best Hospitals specialty rankings, but until 2007 the pediatric rankings continued to rely entirely on an annual survey of pediatric specialists because hard data comparable to the MedPAR files for Medicare recipients was unavailable. Pediatric-specific data were critical. Benchmarking and data generated from adult patients, to the extent that such information existed at all, could not be applied to children. Structuring coordinated care for congenital conditions such as spina bifida and cystic fibrosis, determining drug dosages and minimizing vulnerability to infection are a few of many factors that make pediatric patients unique.

Lacking robust pediatric data bases, U.S. News elected to collect data directly from children’s hospitals through a comprehensive clinical and operational survey. The first rankings to incorporate data from such a survey, developed by RTI International*, were published in 2007 in the form of the top 30 children’s centers in General Pediatrics. Specialty rankings were not included.

In the years that followed, data collection was broadened and deepened. The current methodology continues to include reputational survey results (expert opinion) as well as supplemental information from resources such as the National Cancer Institute. Best Children’s Hospitals now ranks the top 50 centers in 10 specialties: Cancer, Cardiology & Heart Surgery, Diabetes & Endocrinology, Gastroenterology & GI Surgery, Neonatology, Nephrology, Neurology & Neurosurgery, Orthopedics, Pulmonology & Lung Surgery and Urology.

To provide parents with information about more centers and demonstrate transparency, pediatric centers below the line – that is, those that are not nationally ranked – are now displayed.

* RTI International is the trade name of Research Triangle Institute.

Pediatric centers that provided sufficient data to receive an overall U.S. News Score but fell short of the top 50 are displayed with their calculated metrics but without rank or score.

Each of the 193 facilities surveyed for the 2020-21 Best Children’s Hospitals rankings is either a freestanding children’s hospital or a “hospital within a hospital” – a large, essentially autonomous multidisciplinary pediatric department within a major medical center. Most are members of the Children’s Hospital Association (CHA).[†]

RTI International[‡] collects and analyzes the data for the “Best Children’s Hospitals” rankings. The methodology reflects *clinical outcomes*, such as patient survival, infection rates and complications; the level and quality of *hospital resources* directly related to patient care, such as staffing, technology and special services; *delivery of healthcare*, such as programs that prevent infections and adherence to best practices; and *expert opinion* among pediatric specialists.

In the 2020-21 rankings, 88 of the 193 surveyed hospitals were ranked among the top 50 in at least one specialty. The 2020-21 Best Children’s Hospitals Honor Roll recognizes the 10 hospitals with the highest rankings across all specialties.

[†] The National Association for Children’s Hospitals and Related Institutions (NACHRI) was renamed the Children’s Hospital Association in 2012. See <http://www.childrenshospitals.net> for details.

[‡] RTI International is the trade name of Research Triangle Institute.

Table of Contents

Executive Summary	1
I. Introduction	1
II. Eligibility	3
A. General Eligibility	3
B. Specialty-Specific Eligibility	3
III. Pediatric Hospital Survey	4
IV. Structure	5
A. Structural Measures	6
Accredited by FACT (Cancer).....	6
Active Fellowship Program (All Specialties)	6
Adoption of Health Information Technology (All Specialties)	8
Adult Congenital Heart Program (Cardiology & Heart Surgery)	9
Advanced Clinical Services Offered (All Specialties)	9
Advanced Technologies (All Specialties).....	17
Bone Marrow Transplant Services (Cancer)	20
Clinical Support Services Offered (All Specialties).....	20
Commitment to Clinical Research (All Specialties)	22
Commitment to Quality Improvement (All Specialties).....	25
Congenital Heart Program (Cardiology & Heart Surgery)	27
ECMO Availability (Neonatology)	27
Enlists Families in Structuring Care (All Specialties).....	27
Has Fulltime Subspecialists Available (All Specialties)	28
Heart Transplant Program (Cardiology & Heart Surgery)	34
Help for Families (All Specialties)	34
Liver Transplant Program (Gastroenterology & GI Surgery)	35
Lung Transplant Program (Pulmonology & Lung Surgery).....	35
Neonatal Transport (Neonatology)	36
Nurse Staffing (All Specialties).....	36
Percent of Dialysis Patients Who Had Transplants (Nephrology)	37
Provides Advanced Palliative Care Program (Cancer).....	37
Recognized as Nurse Magnet Hospital (All Specialties)	38
Specialized Clinics and Programs (Cancer, Diabetes & Endocrinology, Gastroenterology & GI Surgery, Neonatology, Neurology & Neurosurgery, Orthopedics, Urology)	38
Success in Helping Patients Manage Their Asthma (Pulmonology & Lung Surgery).....	40
Success in Managing Neuromuscular Weakness Disorder (Pulmonology & Lung Surgery)	41
Tracking Growth Metrics for Treated Patients (Neonatology)	41
Volume of Patients (All Specialties)	41
B. Normalization.....	52
C. Weighting	52
V. Process	55

A.	Commitment to Best Practices	55
B.	Ability to Prevent Infections.....	85
	All-Specialty Infection-Preventing Measures.....	86
	Specialty-Specific Infection-Preventing Measures	87
C.	Expert Opinion with Pediatric Specialists	90
	2020 Survey Approach.....	91
	Log Transformation.....	96
D.	Normalization and Weighting.....	97
VI.	Outcomes	98
A.	Outcome Measures.....	98
	Cancer.....	98
	Cardiology & Heart Surgery.....	100
	Diabetes & Endocrinology	104
	Gastroenterology & GI Surgery.....	107
	Neonatology	109
	Nephrology.....	110
	Neurology & Neurosurgery	112
	Orthopedics.....	113
	Pulmonology & Lung Surgery.....	114
	Urology.....	116
B.	Normalization and Weighting.....	119
VII.	Calculation of the U.S. News Score	121
VIII.	Pediatric Honor Roll.....	122
IX.	2020-21 Changes	123
X.	Future Improvements	125
XI.	Contact Information	126
XII.	References	126

List of Tables

Table 1. Specialty-Specific Eligibility Requirements	4
Table 2. Active Fellowship Programs by Specialty.....	7
Table 3. Advanced Clinical Services Offered by Specialty.....	10
Table 4. Advanced Technologies by Specialty	17
Table 5. Clinical Support Services by Specialty.....	21
Table 6. Subspecialists by Specialty.....	29
Table 7. Volume Measures by Specialty.....	42
Table 8. Relative Weights of Individual Structural Measures by Specialty.....	53
Table 9. Commitment to Best Practices by Specialty	55
Table 9. Commitment to Best Practices by Specialty, continued	56
Table 9. Commitment to Best Practices by Specialty, continued	57
Table 9. Commitment to Best Practices by Specialty, continued	58
Table 9. Commitment to Best Practices by Specialty, continued	59
Table 10. Core Infection-Preventing Measures, All Specialties (31 points).....	86
Table 11. Expert Opinion Weight by Survey Year	90
Table 12. Population Counts by Best Hospitals Specialty, Proximity Members and Nonmembers ...	92
Table 13. Member Survey Response Rates (%) by Region and Specialty, 2020	94
Table 14. Nonmember Survey Response Rates (%) by Region and Specialty, 2020	95
Table 15. Weight of Individual Process Measures (All Specialties Except Pediatric Cardiology & Heart Surgery).....	97
Table 16. Relative Weights of Outcomes Measures by Specialty	120
Table 17. Component Weighting	121

List of Figures

Figure 1. Impact of Log Transformation on Expert Opinion.....	96
Figure 2. Example funnel plot of AMR and patient volume with 1 and 2 standard error lines used for point boundaries	102
Figure 3. Example funnel plot of optimal A1C values for patients 13-17 years of age on private insurance with Type 1 diabetes and patient volume with 1 standard error lines used for point boundaries	105
Figure 4. Example funnel plot unplanned hospital admissions for urologic issue within 30 days of inpatient urological surgery and surgical volume with 1 standard error lines used for point boundaries	117

List of Appendixes

Appendix A 2020-21 Best Children's Hospitals Working Groups and Members.....	A-1
Appendix B Glossary of Terms.....	B-1
Appendix C 2020-21 Sample Physician Questionnaire	C-1
Appendix D 2020-21 Best Children's Hospitals Rankings by Specialty.....	D-1
Appendix E 2020-21 Best Children's Hospitals Honor Roll	E-1

I. Introduction

Rankings in pediatrics were included in the initial “America’s Best Hospitals” rankings in 1990. Until 2007, however, the pediatric rankings relied entirely on reputational surveys of board-certified pediatricians and adolescent-medicine specialists.

The reason was that quantitative pediatric measures in barely existed. A large, rich database, comparable to the Centers for Medicare & Medicaid Services MedPAR (Medicare Provider Analysis and Review) files that determine mortality in 12 adult specialties, was unavailable^{**}. Reliable structural measures also were absent. Available data sources generally reported volume, advanced technologies and patient services for the hospital as a whole and did not break out pediatric-specific information.

Continuing to rank children’s hospitals solely on expert opinion for an indeterminate period while performance data were codified and the means of collecting and verifying them settled was felt to be unacceptable. U.S. News asked RTI International, U.S. News contractor for the adult Best Hospitals rankings, to develop a rigorous methodology for ranking hospitals in pediatrics that would incorporate data obtained directly from the hospitals.

The resulting methodology and initial version of a direct hospital survey (referenced in this report as the Pediatric Hospital Survey) produced General Pediatrics rankings of 30 hospitals, published in the September 3, 2007, issue of U.S. News & World Report as “Best Children’s Hospitals.” The issue was separate from the issue with the adult rankings, to highlight the change and minimize possible confusion.

The Pediatric Hospital Survey and the reputational Physician Survey were expanded in 2008, permitting pediatric hospitals to be ranked in six pediatric specialties as well as in General Pediatrics.^{††} In 2009, General Pediatrics was dropped and the specialties expanded to 10 that still define the specialty universe:

- Cancer
- Cardiology & Heart Surgery
- Diabetes & Endocrinology
- Gastroenterology & GI Surgery
- Neonatology
- Nephrology
- Neurology & Neurosurgery
- Orthopedics
- Pulmonology & Lung Surgery
- Urology

^{**} A relatively small number of children do receive care under Medicare under narrow eligibility definitions because of legislatively mandated coverage.

^{††} Previous methodology reports are available online at www.rti.org/besthospitals.

Like their adult counterpart, the Best Children's Hospitals rankings reflect the interrelationship between structure, process and outcomes, the three components of the Donabedian paradigm.¹⁻⁵ Individual measures, their weights and approach to scoring are quite different in the pediatric rankings, however.

The Donabedian components represent the following healthcare concepts:

- *Structure* refers to hospital resources directly related to patient care. Examples include the ratio of nurses to patients, specialized clinics and programs, and certification by recognized external organizations.
- *Process* encompasses overall rendering of diagnosis, treatment, prevention and patient education. Process is represented in part by an expert opinion score based on the annual survey of board-certified physicians cited above. Starting with the 2012-13 rankings, the pediatric methodology has incorporated compliance with best practices and activities to prevent infections and other patient safety issues.
- *Outcomes* obviously include survival but can also include functional success (as in children with cystic fibrosis) and incidence of adverse events (such as bloodstream infections and transplant-organ failure).

Each major component of the Best Children's Hospitals ranking score—structure, outcomes and process—is worth exactly one-third of the overall score other than in Cardiology & Heart Surgery, in which outcomes weight in the overall score has been increased in the 2017-18 rankings to 38.3 percent and the process weight lowered to 28.3 percent. (Details are provided in Table 15.) The specific measures, their weights and the scoring algorithm all are quite different from their adult counterparts.

Section II of this report outlines the general eligibility requirements for consideration in the pediatric rankings. As in previous years, most structure and outcomes data for the 2020-21 rankings were obtained directly from children's hospitals through the Pediatric Hospital Survey (**Section III**). Data for three measures were supplied by external organizations: Nurse Magnet recognition (American Nurses Credentialing Center), accreditation by FACT for BMT and tissue transplant (Foundation for the Accreditation of Cellular Therapy) and commitment to best practices in treating patients with seizure disorders (National Association of Epilepsy Centers).

The specific mission of the Best Children's Hospitals rankings is to identify hospitals that provide the highest quality of care for children with the most serious or complicated medical conditions, using the most robust and sensitive measures available to represent the three

Donabedian components. *Sections IV, V, and VI* describe the data and the construction of each component.

The methodology also incorporates nominations of hospitals from board-certified pediatric specialists in each of the 10 specialties through the Pediatric Physician Survey, as described in *Section V*.

II. Eligibility

A. General Eligibility

To be considered for the 2020-21 pediatric rankings, hospitals had to provide extensive data about their services and capabilities through the 2020-21 Pediatric Hospital Survey (<https://usnewspediatricsurvey.rti.org/PaperSurvey.aspx>).

Historically, initial eligibility for the rankings has been determined by membership in the Children’s Hospital Association (CHA) or by nomination by teams of expert advisers. U.S. News and RTI have supplemented the universe by adding hospitals that have expressed interest in public reporting, after consideration of the size and scope of each hospital’s pediatric program.

Of the 193 hospitals that qualified for 2020-21 inclusion, 118, or 61.1 percent, submitted sufficient data through the Pediatric Hospital Survey to be considered for ranking in at least one specialty. Each facility met the description of one of three groups: a freestanding children’s hospital, a “hospital within a hospital” (as described above, a pediatric service that functions autonomously within a larger medical center) or a specialty hospital (such as one that exclusively treats cancer or orthopedic patients).

As of the 2016-17 rankings, participating hospitals had to agree that their evaluated data would be publicly displayed even if they did not rank among the top 50 in a specialty. In previous years, only results for the top 50 hospitals in a specialty were publicly available. This requirement provides families with additional important information and demonstrates commitment to transparency by the hospitals and the Best Children’s Hospital rankings process.

B. Specialty-Specific Eligibility

To be eligible for ranking within a given specialty, hospitals had to satisfy two additional requirements:

- In specialties other than Neonatology, a hospital had to verify in the Pediatric Hospital Survey that services in the specialty were in fact available. In Neonatology, a

hospital also had to have a Level IV neonatal intensive care unit (NICU). Validation of this NICU status could be met based either on state determination of Level IV status or satisfaction of Level IV eligibility requirements as specified by the American Academy of Pediatrics guidelines.^{‡‡}

- A full-time equivalent (FTE) of at least 1.0 attending physicians in certain specialty-related medical fields was required. The physician categories are shown in **Table 1**. Text and table references (e.g. “B2a”) indicate the related section and question in the Pediatric Hospital Survey.

Table 1. Specialty-Specific Eligibility Requirements

Specialty	Must have at least 1.0 FTE attending staff in the following categories:
Cancer	Pediatric hematologist/oncologist (B2a)*
Cardiology & Heart Surgery	Pediatric cardiothoracic surgeon (E2a) and Pediatric cardiac intensivist (from training in cardiology, pediatric critical care or anesthesiology) or Other pediatric cardiac specialist (pediatric cardiac interventionalist, pediatric cardiac electrophysiologist, pediatric anesthesiologist with specialty cardiac training, or pediatric advanced imaging specialist (cardiologist or radiologist)) (E2b, E2c, E2d, E2e, E2f, E2g, or E2h)
Diabetes & Endocrinology	Pediatric endocrinologist (C2a)
Gastroenterology & GI Surgery	Pediatric gastroenterologist (D2a)
Neonatology	Pediatric neonatologist (F2a)
Nephrology	Pediatric nephrologist (G2a)
Neurology & Neurosurgery	Pediatric neurologist (H2a) or Pediatric neurosurgeon (H2b)
Orthopedics	Pediatric orthopedic surgeon (I2a)
Pulmonary	Pediatric pulmonologist (J2a) or Pediatric sleep medicine physician (J2b)
Urology	Pediatric urologist (K2a)

* Parenthetical references indicate related survey questions

III. Pediatric Hospital Survey

As part of the process of creating the initial pediatric rankings, RTI convened advisory panels to inform the hospital survey. These working groups have been retained to help the survey

^{‡‡} AAP guidelines, Pediatrics, 2012, 130:587-597.

evolve by providing new findings and perspectives that can be incorporated before the survey is finalized and sent to hospitals.

Panel members do not serve fixed terms. Members who drop out through normal attrition are replaced by RTI through a request to the pediatric hospital community for candidates with broad expertise in both general and specialty pediatric medical care and familiarity with current research on hospital quality. The 2020-21 panels included pediatric physicians, nurses, hospital quality experts, health information systems/coding experts and other healthcare professionals. A group of infection-control experts and a group of radiologists worked with the 10 specialty panels to address specialty-specific infection-control and prevention issues. The names and institutions of all individual working group members are provided, with their permission, in *Appendix A*.

Through conference calls, ad hoc phone discussions and emails during the summer and fall of 2019, working group members proposed, reviewed and discussed revisions to the previous version, including potential new measures. Smaller subgroups of members in each working group were responsible for reviewing the revised codes to ensure that the selected codes were relevant and appropriate. The RTI project team then created a draft set of measures and a survey instrument.

A draft of the survey was provided as a Microsoft Word document to hospitals at the beginning of December 2019 on an FYI basis, to give them as much time as possible to collect and organize data before the official start date. They received the finalized survey in early January 2020 via a dedicated web page. Survey responses were accepted until mid- March.

Some measures were ultimately excluded after data were submitted because the results failed to demonstrate meaningful variability. The remaining items are described in detail below, with references to the corresponding survey question numbers in parentheses.

The Pediatric Hospital Survey will continue to be updated and modified in subsequent years to reflect the quality of care provided by U.S. pediatric facilities and the evolving discipline of quality improvement.

IV. Structure

The structural component is represented by volume, technology, clinical services and other characteristic features of a high-quality pediatric hospital. In the Best Hospitals adult specialty rankings, most structural measures and their associated data are extracted from the American Hospital Association (AHA) annual survey. Because the AHA survey focuses primarily on overall hospital and system measures, the pediatric data from the survey lack specificity. Structural data were therefore collected through the Pediatric Hospital Survey.

All measures used in the rankings are described in the following sections. The print version of the rankings displays a subset of the online measures.

A. Structural Measures

The structural measures used in the rankings represent fundamental elements of high-quality, hospital-based pediatric care. Descriptions of the measures and the specialties to which they are applied are listed alphabetically. Text and table references such as (A6a) indicate the related section and question in the Pediatric Hospital Survey. Each measure's relative weight within a specialty is provided in *Section IV.B. Normalization and Weighting*.

Accredited by FACT (Cancer)

Accreditation indicates that as of March 1, 2020, a hospital met standards set by the Foundation for Accreditation of Cellular Therapy for transplanting cells to treat pediatric cancer, an indication of a high degree of care in handling and using cellular tissue. Programs could be certified as providing adult or pediatric services and as offering two types of transplant services: autologous and allogeneic. For the Cancer specialty, a hospital was awarded 1 point if it was accredited by FACT as a pediatric or adult service provider for either autologous or allogeneic transplants (B19a). Currently accredited facilities are listed at <http://accredited.factwebsite.org/>.

Active Fellowship Program (All Specialties)

Participation in fellowship training programs represents a commitment by hospitals to provide high-quality care in a specialty area and assure that their programs meet standards of quality. Hospitals that offer fellowship programs accredited by the Accreditation Council for Graduate Medical Education (or other advanced fellowship programs) were awarded 1 point for each fellowship program that had at least one active fellow in the program in the past academic year. Each specialty has one or more programs that are considered flagship programs in their specialty. Hospitals that have at least one active fellow in these programs are awarded 2 points for each program. *Table 2* indicates fellowships credited and the number of points for each specialty.

Table 2. Active Fellowship Programs by Specialty

Fellowship Program*	Cancer	Cardiology & Heart Surgery	Diabetes & Endocrinology	Gastroenterology & GI Surgery	Neonatology	Nephrology	Neurology & Neurosurgery	Orthopedics	Pulmonology & Lung Surgery	Urology
Child neurology (A6a)					1		2			
Congenital cardiac surgery (A6b)		2			1					
Neonatal-perinatal medicine (A6c)					2					
Pediatric Neurosurgery (A6d)	1						2			
Pediatric cardiology (A6e)		2			1					
Pediatric endocrinology (A6f)			2							
Pediatric gastroenterology (A6g)				2	1					
Pediatric hematology-oncology (A6h)	2									
Pediatric nephrology (A6i)						2				
Pediatric neuroradiology (A6j)	1									
Pediatric pulmonology (A6k)					1				2	
Pediatric urology (A6l)	1									2
Pediatric surgery (A6m)	1	1								
Pediatric infectious diseases (A6n)	1	1	1	1	1	1	1	1	1	1
Orthopedic surgery of the spine, including POSNA fellows (with training in pediatrics) (A6o)								2		
Pediatric critical care medicine (A6p)	1	1	1	1	1	1	1	1	1	1
Pediatric advanced transplant hepatology (A6q)				1						
Pediatric rheumatology (A6r)			1					1		
Pediatric physical medicine and rehabilitation (A6s)		1	1	1		1	1	1	1	1
Pediatric radiology (A6t)	1	1	1	1	1	1	1	1	1	1

(continued)

Table 2. Active Fellowship Programs by Specialty (continued)

Fellowship Program*	Cancer	Cardiology & Heart Surgery	Diabetes & Endocrinology	Gastroenterology & GI Surgery	Neonatology	Nephrology	Neurology & Neurosurgery	Orthopedics	Pulmonology & Lung Surgery	Urology
Pediatric interventional radiology (A6u)	1					1	1			
Advanced motility training program (D34)				1						
Advanced nutritional training program (D35)				1						
Advanced hepatology training program (D36)				1						
Pediatric orthopedics (I6.1)								2		
Total Elements	10	9	7	10	10	7	9	9	6	6

* Parenthetical references indicate related survey questions

Adoption of Health Information Technology (All Specialties)

In each specialty, hospitals received up to 14 points for incorporating and using a computerized physician order entry (CPOE) system and electronic medical records (EMRs).

Hospitals received up to 6 points for CPOE: 1 point for implementing a CPOE system (A20), 1 point for documenting 95% or more of inpatient medication orders (A21a), 1 point for identifying medication orders if an allergy to the medication is documented (A21b), 1 point for including alerts for dosing errors for high-risk medications (A21c) and up to 2 points for providing details on two or more current projects using CPOE that focus on dosing errors for high-risk medications (A21.1).

Hospitals received up to 8 points for EMR: 1 point if the EMR identifies and reports potential adverse events for patients (A23), 1 point if they could exchange patient information with organizations that have the same EMR system or 2 points if they could exchange patient information with organizations that have a different EMR system (A23.2). Hospitals that have an EMR with certain patient engagement features received up to 5 points, 1 for each of the following: online access to medical notes or records (A23.3a), ability to request a revision to medical notes or records online (A23.3b), ability to schedule visits online (A23.3c), ability to send and receive electronic messages with medical providers (A23.3d), and patients have online access to radiology reports (A23.3e).

Adult Congenital Heart Program (Cardiology & Heart Surgery)

In Cardiology & Heart Surgery, hospitals received up to 11 points for having an adult congenital heart program. Hospitals received 1 point for providing an organized adult congenital heart program (E16). Hospitals received up to 2 additional points based on the status of accreditation with the Adult Congenital Heart Association to become an Adult Congenital Heart Association Comprehensive Care Center (E20): 2 points if the program is fully accredited, or 1 point if the program has partially completed the accreditation process. These programs are often provided by pediatric heart centers, which frequently have the most expertise in inherited and congenital heart disorders.

Up to 6 additional points were awarded if the adult congenital heart program provided the following: a formal program to transition patients from the pediatric to adult congenital heart program (E17a); joint participation by adult and pediatric cardiologists (E17b); participation by cardiothoracic surgeons (E17c), cardiothoracic interventionalists (E17d) and cardiothoracic electrophysiologists, who have specialty expertise in the care of adults with congenital heart disease (E17e); and specialty care for high-risk obstetrics patients with congenital heart disease (E17f).

Hospitals received 1 point for 1 to 49 cardiac surgical encounters^{§§} on patients age 18 and above in the past four calendar years or 2 points for 50 or more surgical encounters in the past four calendar years (E19).

Advanced Clinical Services Offered (All Specialties)

Hospitals frequently offer clinical services and organize teams or programs to address special needs of specific groups of patients. These services or programs may be organized around a particular diagnosis, need or age group. The structure of the services or programs ensures that a range of resources is available. Specialized skills of a multidisciplinary staff improve overall quality of care, and presumably outcomes. The clinical services recognized in each specialty are described in **Table 3**. Up to eight points were awarded for having a pediatric trauma center in Neurology & Neurosurgery, Orthopedics, and Pulmonology & Lung Surgery. The trauma center measure recognizes the enhanced resources and staff available to hospitals that provide this service, which benefit other inpatient specialty care. One point was awarded for the additional services listed for each specialty.

^{§§} Specific adult cardiac surgical operations included are listed in Table 7 of the STS Congenital Heart Surgery Database for the past four reporting years.

Table 3. Advanced Clinical Services Offered by Specialty

Cancer (27 points)		
Service	Description*	Points
Sedation services	Provides sedation/anesthesia by pediatric specialists for radiation therapy, lumbar punctures and bone marrow biopsies. (B7)	1
Support staff/programs	<p>Offers the following programs and supporting staff (B11, B11.1, and B11.2):</p> <ul style="list-style-type: none"> • Complementary and alternative medicine or holistic health program • Pediatric cancer child-life specialists • Psychosocial support program • Social work support • School programs for hospitalized patients • Neuropsychological evaluation focused on school re-entry issues • APHON chemotherapy/biotherapy course and safe handling procedures • Adolescent and young adult support program • Fertility preservation program • Cancer genetics/hereditary program • Sibling targeted support services • Bereavement services for families • Developmentally appropriate procedural preparation support for invasive medical procedures • On-site inpatient pediatric rehabilitation unit with individualized dedicated cancer rehabilitation programming • Having direct clinical care RNs with a national oncology certification (certified pediatric hematology-oncology nurse (CPHON) or certified pediatric oncology nurse (CPON)) or certified bone marrow transplant nurse (BMTCN) <ul style="list-style-type: none"> ○ 1 point for 25% or more of direct clinical care RNs with these certificates ○ 2 points for 50% or more of direct clinical care RNs with these certificates • Having 50% or more of chemotherapy patients with a formal initial psychosocial assessment before or within 4 weeks of therapy 	17

* Parenthetical references indicate related survey questions

(continued)

Table 3. Advanced Clinical Services Offered by Specialty (continued)

Cancer, continued (27 points)		
Service	Description*	Points
Chemotherapy support services	Offers the following: <ul style="list-style-type: none"> • Dedicated pediatric chemotherapy pharmacy (B15a) • Pediatric oncology pharmacist (B15b) • Pharmacists assigned to participate in daily inpatient rounds with the pediatric cancer treatment team (B15c) • Formal annual chemotherapy training (e.g., order writing, dispensing, administration) (B15d) • Formal chemotherapy safety program with standardized procedures and event tracking (including order misses/near-misses) (B15e) • Designated pediatric oncology faculty leader for the chemotherapy safety program (B15f) 	6
Chemotherapy orders	1 point for orders written using word processing or spreadsheet software; 2 points for CPOE; 3 points for CPOE with plan-driven orders and formal multiple co-signatures/review required (B16)	3
Cardiology & Heart Surgery (15 points)		
Service	Description*	Points
ECMO	ECMO program designated as center of excellence by the Extracorporeal Life Support Organization (ELSO) (A9)	1
Echocardiography laboratory	Offers certified echocardiography laboratory certified by Intersocietal Commission for the Accreditation of Echocardiography Laboratories (ICAEL) or the American Institute of Ultrasound in Medicine (AIUM) (E5) in: <ul style="list-style-type: none"> • Transthoracic echocardiographic testing • Transesophageal echocardiographic testing • Fetal echocardiographic testing 	3
Cardiovascular services	Offers these diagnostic and treatment services (E6a-i, E6k): <ul style="list-style-type: none"> • Dedicated pediatric cardiac surgical operating room • Cardiac intensive care unit • Remote monitoring capability • Cardiac diagnostic catheterization laboratory • Cardiac interventional catheterization laboratory • Electrophysiology laboratory • Ventricular assist program • 24/7 ECMO • Cardiovascular genetics clinic • Pediatric cardiac anesthesia services 	10
Circulatory support	Provided ventricular assist devices (other than ECMO) for one or more patients in the past 4 years (E26)	1

* Parenthetical references indicate related survey questions

(continued)

Table 3. Advanced Clinical Services Offered by Specialty (continued)

Diabetes & Endocrinology (18 points)		
Service	Description*	Points
Diabetes & Endocrinology support staff	Having at least 1 of the following staff with Certified Diabetes Educator certification provide diabetes education to patients: <ul style="list-style-type: none"> • Nurses, pharmacists, social workers, psychologists (C5a and C5c) • Dietitians (C5b) • Certified exercise physiologist or Physical therapist (C5d) 	2
	Having at least 1.0 FTE of the following staff dedicated to pediatric endocrinology patients: <ul style="list-style-type: none"> • Social workers (C6a) • Psychologists (C6b) • Genetic counselors (C7a) • Psychiatrists (C7b) • Pharmacists (C7c) 	6
Diabetes patient services	Provides the following services onsite (C9): <ul style="list-style-type: none"> • Written educational protocol used to evaluate and prepare patients for use of an insulin pump • Certified pump educators to provide insulin pump training to patients and their families • Written education program used to evaluate and prepare patients for use of continuous glucose monitors (CGMs) • Certified CGM trainers to provide CGM training to patients and their families • Written educational program for families of new-onset diabetes patients • Formal diabetes educational program for school nurses through a yearly school nurse education conference • A specified RN or CDE responsible for advising and supporting schools in setting up safe programs for managing diabetes 	7
Support services	Offered the following programs or services in the last calendar year: <ul style="list-style-type: none"> • Hosted or was actively involved in organizing diabetes-specific support group for parents and families (C12) • Took a leadership role in organizing or supporting family-support groups for special populations other than diabetes (e.g., Turner syndrome) (C60) • A Family Advisory Board that includes families of non-diabetes Endocrinology patients (C61) 	3

* Parenthetical references indicate related survey questions

(continued)

Table 3. Advanced Clinical Services Offered by Specialty (continued)

Gastroenterology & GI Surgery (9 points)		
Service	Description*	Points
Gastro-intestinal (GI) specialists	Has following specialists available for consultation 7 days a week (D8): <ul style="list-style-type: none"> • Pediatric gastroenterology/liver-specialized pathologists • Interventional radiologists with pediatric gastroenterology experience • Pediatric anesthesiologists for endoscopy sedation/anesthesia 	3
GI support groups	Provides access to the following support groups (D12): <ul style="list-style-type: none"> • Inflammatory bowel disease • Celiac disease • Liver disease or transplant • Eosinophilic esophagitis • Chronic intestinal failure • Congenital colorectal malformation 	6
Neonatology (11 points)		
Service	Description*	Points
NICU support staff	NICU-dedicated staff in these units: <ul style="list-style-type: none"> • NICU-dedicated respiratory therapy team who attends weekday work rounds daily with clinical team (F7a) • NICU-designated dietician who attends weekday work rounds daily with clinical team (F7b) • NICU-dedicated social workers (F11) 	3
Pediatric Subspecialists	Has following subspecialists on-call and available to be on-site 24/7 (F11.2): <ul style="list-style-type: none"> • Gastroenterologist • Nephrologist • Pediatric Ophthalmologist • Pulmonologist • Urologist • Echocardiographer • Geneticist • Infectious Disease Specialist 	8

* Parenthetical references indicate related survey questions

(continued)

Table 3. Advanced Clinical Services Offered by Specialty (continued)

Nephrology (8 points)		
Service	Description*	Points
Maintenance dialysis staff	Has at least 1 FTE of clinical nurses dedicated to maintenance dialysis (G5a)	1
	Has at least 0.5 FTE of the following staff dedicated to maintenance dialysis: <ul style="list-style-type: none"> • Social workers (G5b) • Dieticians (G5c) • Psychologists/Psychiatrists (G5d) 	3
Dialysis treatment	Provides following dialysis options for acute kidney insufficiency (G7): <ul style="list-style-type: none"> • Hemodialysis • Peritoneal dialysis • Continuous renal replacement therapy 	3
Kidney transplant	United Network for Organ Sharing (UNOS)-recognized kidney transplant program (G28)	1
Neurology & Neurosurgery (21 points)		
Service	Description*	Points
Pediatric trauma center	8 points for Level 1 pediatric trauma center or 4 points for Level 2 pediatric trauma center certified by American College of Surgeons or state licensing board (A19)	8
Neurology & neurosurgery support services and technology	Offers the following: <ul style="list-style-type: none"> • Ketogenic diet evaluation and management program (H5c) • Neuroendovascular interventionalists (H5d) • Neuroanesthesia program (H5e) • A dedicated multidisciplinary neurocritical care team (H24) • Multidisciplinary neonatal neurocritical care (H24.1) • Inpatient pediatric rehabilitation program with pediatric physiatrist (H13) • Inpatient pediatric rehabilitation program certified by Commission on Accreditation of Rehabilitation Facilities (H13.1) • Inpatient pediatric rehabilitation program that participates in and submits data to the Universal Data System for Medical Rehabilitation (UDSMR) (H13.2) • Neuropsychological testing by pediatric neuropsychologists (H14) 	9
Epilepsy treatment	Offers the following: <ul style="list-style-type: none"> • Electroencephalography (EEG) lab accredited by ABRET (H7) • In-house EEG technologists available 24/7 on 300 or more days per year to place electrodes (H7.1) • In-house EEG technologists available to review EEG continuously 24/7 on 300 or more days per year (H7.2) • Epilepsy monitoring unit with emergency management of seizures protocols (H30) 	4

* Parenthetical references indicate related survey questions

(continued)

Table 3. Advanced Clinical Services Offered by Specialty (continued)

Orthopedics (16 points)		
Service	Description*	Points
Pediatric trauma center	8 points for Level 1 pediatric trauma center or 4 points for Level 2 pediatric trauma center certified by American College of Surgeons or state licensing board (A19)	8
Advanced care services	Comprehensive pediatric orthopedic program with: <ul style="list-style-type: none"> • Designated inpatient unit for pediatric orthopedic patients (I7) • Dedicated pediatric imaging center located in outpatient clinics (not separate facility) (I8) • Imaging center staffed by a pediatric radiologist with qualification in pediatric radiology by American Board of Radiology (I9) • Multidisciplinary musculoskeletal oncology program (I16) • Gait laboratory accredited by the Commission for Motion Laboratory Accreditation (CMLA) (I19.1) • Providing seating services or wheelchair clinics for at least 1 patient with neuromuscular disorders (I43 & I44) 	6
Gait Laboratory	Provides an Advanced Motion Analyses Laboratory to patients (I19): <ul style="list-style-type: none"> • 2 points if the lab is on-site • 1 point if the lab is available through a formal contractual relationship with another hospital/institution 	2
Pulmonology & Lung Surgery (38 points)		
Service	Description*	Points
Pediatric trauma center	8 points for Level 1 pediatric trauma center or 4 points for Level 2 pediatric trauma center certified by American College of Surgeons or state licensing board (A19)	8
Asthma care specialists	Access to at least 0.5 FTE staff with clinical responsibilities (J5): <ul style="list-style-type: none"> • Respiratory therapists • Social workers • Dieticians • Physical therapists • Psychiatrists or psychologists • Genetic Counselor • Child Life Specialist • Pharmacist • Speech and language pathologists 	9

* Parenthetical references indicate related survey questions

(continued)

Table 3. Advanced Clinical Services Offered by Specialty (continued)

Pulmonology & Lung Surgery, continued (35 points)		
Description*	Description*	Points
Dedicated staff	<p>Following cystic fibrosis center staff who attend clinic or participate in patient care conferences (J17):</p> <ul style="list-style-type: none"> • Gastroenterologist • Endocrinologist • Psychiatrists or psychologists <p>Following staff who support patients with bronchopulmonary dysplasia (J28):</p> <ul style="list-style-type: none"> • Gastroenterologist • Cardiologist • Neurologist or Neurodevelopmental Specialist • Social worker • Occupational Therapist <p>Following staff who support patients with neuromuscular weakness disorders (J32):</p> <ul style="list-style-type: none"> • Pulmonologist • Physiatrist • Orthopedist • Cardiologist • Neurologist • Physical therapist • Psychiatrist or psychologist • Dietician • Social worker • Pediatric radiologist 	18
Support services	<p>Offers following:</p> <ul style="list-style-type: none"> • Cystic fibrosis center accredited by Cystic Fibrosis Foundation (J16) • Program accredited by the Primary Ciliary Dyskinesia (PCD) foundation (J31) • Sleep center accredited by American Academy of Sleep Medicine (J35) 	3
Urology (3 points)		
Service	Description*	Points
Treatment options	<p>Offers the following treatment modalities (K11):</p> <ul style="list-style-type: none"> • Stone treatment, including shock wave lithotripsy • Laparoscopic orchiopexy/orchidectomy for intra-abdominal testes • Laparoscopic surgery, including cyst ablation, pyeloplasty, nephrectomy, partial nephrectomy, heminephrectomy, ureteral reimplantation, or ureteroureterostomy 	3

* Parenthetical references indicate related survey questions

Advanced Technologies (All Specialties)

To receive credit, hospitals must provide access to key diagnostic and treatment technologies. For the technologies listed in A10, hospitals had to offer services onsite. For other technologies, hospitals could offer the services onsite or through the hospital's health system, a local community network or indirectly, through a contractual arrangement or joint venture with another community provider. Data are from the Pediatric Hospital Survey. The values for this measure were based on specialty-specific mixes of technology, as listed in *Table 4*. Definitions can be found in the glossary in *Appendix B*.

Table 4. Advanced Technologies by Specialty

Specialty	Technologies*
Cancer (17 technologies)	<ul style="list-style-type: none"> • Positron emission tomography (PET)/magnetic resonance imaging (MRI) or PET/computerized tomography (CT) scanning offered onsite (A10a or A10b) • Intraoperative magnetic resonance imaging (ioMRI) offered onsite (A10c) • 3-Tesla magnetic resonance imaging (3T MRI) offered onsite (A10d) • Image-guided radiation therapy offered onsite (A10e) • Intensity-modulated radiation therapy offered onsite (A10f) • Fast shunt MRI for hydrocephalus offered onsite (A10h) • Nuclear medicine SPECT/CT (A10j) • Contrast-enhanced ultrasound (A10k) • Linear accelerator or other linear particle accelerator, Gamma Knife, CyberKnife, or other shaped-beam stereotactic radiation therapies (A11) • Therapeutic meta-iodo-benzyl-guanidine with I-131 radionuclide (B8a) • Functional magnetic resonance (B8b) • Brachytherapy (B8c) • Stereotactic radiosurgery (B8d) • Intra-arterial chemotherapy or embolization for solid tumors (B8e) • Radiofrequency ablation and/or cryoablation (B8f) • Proton Beam Therapy with Pencil-Beam Scanning (PBS) capability (B8g) • Pediatric interventional radiology equipment and room (B9)

* Parenthetical references indicate related survey questions

(continued)

Table 4. Advanced Technologies, by Specialty (continued)

Specialty	Technologies*
Cardiology & Heart Surgery (8)	<ul style="list-style-type: none"> • Dedicated interventional radiology team offered onsite (A10i) • Nuclear medicine SPECT/CT (A10j) • Intraoperative transesophageal echocardiographic testing (E6j) • Cardiac CT angiography (E7a) • Cardiac MRI with functional cardiac imaging (E7b) • Stress echo testing (E7c) • Quantitative Pulmonary Perfusion Scan (E7d) • Transcatheter arrhythmia ablation methodologies (three-dimensional mapping, cryoablation or radiofrequency ablation) (E14a-c)
Diabetes & Endocrinology (11)	<ul style="list-style-type: none"> • PET/MRI or PET/CT scanning offered onsite (A10a or A10b) • Dedicated interventional radiology team offered onsite (A10i) • Nuclear medicine SPECT/CT (A10j) • Diagnostic radioisotope scan (C51a) • Therapeutic radioiodine treatment for Graves' disease (C51b) • Therapeutic radioiodine treatment for thyroid cancer (C51c) • Ultrasound guided fine needle biopsy or aspiration of thyroid nodule (C51d) • Thyroidectomy (C51e) • Intraoperative PTH assay (C51f) • Intravenous bisphosphonate therapy (C51g) • Endocrine testing and infusion studies (with endocrinology providers on site) (C55)
Gastroenterology & GI Surgery (19)	<ul style="list-style-type: none"> • PET/MRI or PET/CT scanning offered onsite (A10a or A10b) • 24/7 in house availability of ultrasound for emergency cases (A10g) • Dedicated interventional radiology team offered onsite (A10i) • Nuclear medicine SPECT/CT (A10j) • Contrast-enhanced ultrasound (A10k) • Magnetic resonance cholangiopancreatography (D7a) • Magnetic resonance enterography (D7b) • Ultrasound elastography (USE) for assessing liver fibrosis (D7c) • Magnetic resonance elastography (MRE) for assessing liver fibrosis (D7d) • Ultrasound contrast for liver lesion characterization (D7e) • Capsule endoscopy (D11a) • Endoscopic band ligation/sclerotherapy (D11b) • Esophageal impedance or resolution esophageal manometry (D11c) • Endoscopic retrograde cholangiopancreatography (D11d) • Antroduodenal and full colonic motility studies (D11e) • Esophageal dilation, either bougie or pneumatic (D11f) • Alternative Hemostasis Therapies (Electrocautery, Hemo-Clip application, and Argon plasma coagulation) (D11g) • Deep enteroscopy-single or double balloon (D11h) • Endoscopic ultrasound (D11i)

* Parenthetical references indicate related survey questions

(continued)

Table 4. Advanced Technologies, by Specialty (continued)

Specialty	Technologies*
Neonatology (10)	<ul style="list-style-type: none"> • PET/MRI or PET/CT scanning offered onsite (A10a or A10b) • Fast shunt MRI for hydrocephalus offered onsite (A10h) • Dedicated interventional radiology team offered onsite (A10i) • Continuous video electroencephalography (EEG) monitoring and reading with telemetry capability and with interpretation and consult by a pediatric neurologist (F12a) • Less than 24-hour turnaround time for HSV PCR for cerebrospinal fluid (F12b) • Formal mechanism to order and send samples for genomic analysis and then provide interpretation and counseling of the results within 10-days (F12c) • Less than 24-hour turnaround time for comprehensive respiratory viral molecular testing (F12d) • Less than 24-hour turnaround time for amino acid analysis (F12e) • Less than 24-hour turnaround time for urine organic acid (F12f) • Rapid blood culture (within 24 hours) identification system for differentiation of MSSA from MRSA (F12g)
Nephrology (1)	<ul style="list-style-type: none"> • PET/MRI or PET/CT scanning offered onsite (A10a or A10b)
Neurology & Neurosurgery (13)	<ul style="list-style-type: none"> • PET/MRI or PET/CT scanning offered onsite (A10a or A10b) • 3T MRI offered onsite (A10d) • Fast shunt MRI for hydrocephalus offered onsite (A10h) • Dedicated interventional radiology team offered onsite (A10i) • Nuclear medicine SPECT/CT (A10j) • Neurophysiological intraoperative monitoring (H5a) • EEG source localization (H5b) • Functional MRI (H5f) • Availability of 24/7 EEG monitoring in pediatric intensive care unit (PICU)/neonatal intensive care unit (NICU) (H5g) • Nuclear medicine brain SPECT or brain PET scanning (H5h) • Functional mapping capability and/or diffusion tensor imaging (DTI) evaluation (H5i) • Stereotactic EEG capabilities (H5j) • Transcranial magnetic stimulation (H5k)
Orthopedics (3)	<ul style="list-style-type: none"> • PET/MRI or PET/CT scanning offered onsite (A10a or A10b) • Dedicated interventional radiology team offered onsite (A10i) • Nuclear medicine SPECT/CT (A10j)
Pulmonology & Lung Surgery (2)	<ul style="list-style-type: none"> • PET/MRI or PET/CT scanning offered onsite (A10a or A10b) • Dedicated interventional radiology team offered onsite (A10i)

* Parenthetical references indicate related survey questions

Table 4. Advanced Technologies, by Specialty (continued)

Specialty	Technologies*
Urology (8)	<ul style="list-style-type: none"> • PET/MRI or PET/CT scanning offered onsite (A10a or A10b) • Dedicated interventional radiology team offered onsite (A10i) • Nuclear medicine SPECT/CT (A10j) • Contrast-enhanced ultrasound (A10k) • Dedicated laparoscopic skills lab for faculty and trainees (K7a) • Video pediatric urodynamic fluoroscopy (K7b) • Magnetic resonance urography (MRU) (K7c) • Nuclear Medicine functional assessment of upper urinary tract function and scarring (K7d)

* Parenthetical references indicate related survey questions

Bone Marrow Transplant Services (Cancer)

In Cancer, hospitals could receive up to 15 points for having a stem cell transplant program. Stem cell transplants are critical in treating a variety of cancers:

- Hospitals received 1 point for having a stem cell transplant unit with specially trained pediatric nurses and physicians (B17).
- Hospitals received up to 4 points for offering various stem cell transplant services: autologous stem cell transplantation (B18a), allogeneic matched (related or unrelated) transplantation (B18b), haploidentical (half-matched) transplantation (B18c), and cellular therapy infusions (B18d).
- Hospitals received up to 8 points based on transplant volume (B18). For each of the four types of transplantation listed above, hospitals received points as follows for all transplant except for haploidentical (half-matched) transplantation: 1 point for conducting from 2 to 10 transplants in the past 3 years and 2 points for conducting 11 or more transplants in the past 3 years. For haploidentical (half-matched) transplantation: 1 point for conducting from 2 to 5 transplants in the past 3 years and 2 points for conducting 6 or more transplants in the past 3 years.
- Hospitals received up to 2 points for transplant center recognition by the National Marrow Donor Program (B19b) and for membership in the Pediatric Blood and Marrow Transplant Consortium (B19c).

Clinical Support Services Offered (All Specialties)

Many hospitals provide access to medical and surgical clinical support services through the hospital's health system, a local community network or a contractual arrangement or joint venture with another provider in the community. On- and offsite services received equal credit. Up to 11

services are included in the clinical support services, depending on specialty. Data came from the Pediatric Hospital Survey. For eligible hospitals, specialty-specific mixes of medical and surgical services are used in computing the points for this measure. *Table 5* presents the complete list of medical and surgical services considered for each specialty in 2020-21. Definitions can be found in the glossary in *Appendix B*.

Table 5. Clinical Support Services by Specialty

Clinical Support Service*	Cancer	Cardiology & Heart Surgery	Diabetes & Endocrinology	Gastroenterology & GI Surgery	Neonatology	Nephrology	Neurology & Neurosurgery	Orthopedics	Pulmonology & Lung Surgery	Urology
Neonatal intensive care unit (A7a)	●	●	●	●		●	●	●	●	●
Pediatric intensive care unit (A7b)	●	●	●	●		●	●	●	●	●
Patient care rooms with protective environment (A7c)	●									
Genetic testing/counseling (A7d)	●		●	●	●					
Palliative care program (A7e)	●	●	●	●	●	●	●	●	●	●
Rehabilitation program and consultation service (A7f)	●	●	●	●	●	●	●	●	●	●
Maternal fetal medicine or fetal treatment program (A7g)					●		●			
Vascular tumor program (A35)	●	●	●	●	●	●	●	●	●	●
Rapid-response team available onsite 24/7 (A8a)	●	●	●	●	●	●	●	●	●	●
Pediatric anesthesia program available onsite 24 hours a day (A8b)	●	●	●	●	●	●	●	●	●	●
Pediatric pain management program available onsite 24/7 (A8c)	●	●	●	●	●	●	●	●	●	●
Multidisciplinary pediatric acute pain/sedation service available onsite 24/7 hours a day (A8d)	●	●		●	●	●	●	●	●	●
Total Elements	11	9	9	10	9	9	10	9	9	9

* Parenthetical references indicate related survey questions

Commitment to Clinical Research (All Specialties)

Networks, clinical trials and other research activities advance the ability of the field to treat pediatric patients and also enhance care by making new or novel treatments available at centers that participate in such research.

Cancer (12 points). Hospitals received up to 12 total points for participating in clinical research activities such as clinical trials or other translational research activities. Hospitals received up to 4 points for participating in cancer research networks such as the Children’s Oncology Group (B24a), National Cancer Institute (NCI) Phase 1/Pilot Consortium (B24b), NCI-Designated Cancer Center (B24c), or another cancer-related organized clinical research network (B24d). Hospitals received 1 point each (2 points total) for enrolling at least one patient in a Phase I or Phase II clinical trial (translational research) during the past two years (B25). Hospitals received up to 5 points for engaging in clinical trials in these specific areas: leukemia/lymphoma only (B26a), solid tumors only (B26b), CNS tumors only (B26c), transplants only (B26d), and trials that are not disease-specific (B26e). Hospitals could receive an additional 1 point by demonstrating the depth of their involvement in any of the clinical trials (B26.1).

Cardiology & Heart Surgery (4 points). Hospitals received points for participating in externally audited, national quality-improvement research networks. Hospitals received up to 2 points for participating and contributing data organizations. Hospitals were awarded 1 point for participating in 1, 2 or 3 of the following organizations, or 2 points for participating in 4 or more of the following organizations:

- Society of Thoracic Surgeons (E29a)
- Congenital Heart Surgeons’ Society (E29b)
- National Pediatric Cardiology Quality Improvement Collaborative (E29c)
- Congenital Cardiac Anesthesia Society database (E29d)
- National Cardiovascular Disease Registry—improving pediatric and adult congenital treatment or the Congenital Cardiac Catheterization Project (C3PO) (E29e)
- ELSO registry (E29f)
- Pediatric Cardiac Critical Care Consortium or Virtual Pediatric ICU System (E29g)
- Pediatric Heart Transplant Study (E29h)
- Cardiac Neurodevelopmental Outcome Collaborative (CNOC) (E29i)
- Pediatric Acute Care Cardiology Collaborative (PAC3) (E29j)

- Pedimacs Registry (FDA Database for Ventricular Assist Devices) “and/or” ACTION (E29k)
- ACC QNET program (E29l)
- Congenital Cardiac Catheterization Project (C3PO) (E29m)
- Other externally audited national quality-improvement initiatives (E29.1)

Hospitals received up to 2 additional points based on the number of types of investigative studies they participate in (E30). Hospitals were awarded 1 point for participating in 1 or 2 of the following types of studies or 2 points for participating in 3 or more of the following types of studies: single institution retrospective studies, multi-institutional retrospective studies, basic science studies with extramural funding, prospective clinical trials or studies with industry funding, or prospective clinical trials or studies with competitive extramural funding.

Diabetes & Endocrinology (3 points). Hospitals received up to 3 points based on the number of trials that give patients access to novel, unlabeled medications, diagnostic/monitoring devices or treatment options in the following areas (C67). Hospitals received 1 point for participating in 1 to 8 studies, 2 points for participating in 9 to 30 studies, or 3 points for participating in 31 or more studies in the past year.

Gastroenterology & GI Surgery (7 points). Hospitals received up to 5 points for participating in externally audited, national quality-improvement research networks. Hospitals received 1 point each for participating in prospective research activities: randomized clinical trials (D15a), observational studies (D15b), clinical databases on patient care (D15c), or non-randomized clinical trials (D15d). Hospitals received up to 3 additional points for having IRB-approved studies being led by the Pediatric Gastroenterology & GI Surgery program (D16): 1 point for 1 to 5 studies, 2 points for 6 to 9 studies, or 3 points for 10 or more studies.

Neonatology (4 points). Hospitals received up to 4 total points for participating in externally audited, national NICU treatment and quality-improvement research networks. Hospitals received up to 3 points for participation in the following organizations:

- Vermont Oxford Network, Children’s Hospitals Neonatal Consortium or Child Health Corporation of America database (F24a)
- ELSO data exchange network/registry (F24b)
- Other clinical research or data exchange program (F24c).

Hospitals received 1 additional point for participating in clinical research activities, registered on clinicaltrials.gov, that allow patients access to novel medications or experimental treatment options (F25).

Nephrology (9 points). Hospitals received points for participating in externally audited national quality-improvement research networks. Hospitals received 1 point for participating in specialty-specific clinical research activities that allow patients access to novel medications or experimental treatment options (G39). Hospitals received up to 8 additional points for participation in the following research collaboratives:

- Midwest Pediatric Nephrology Consortium or Pediatric Nephrology Research Consortium (G40a)
- International Pediatric Dialysis Network (G40b)
- North American Pediatric Renal Trials and Collaborative Studies (G40c)
- Nephrotic Syndrome Study Network (G40d)
- Cure GN (G40e)
- Chronic Kidney Disease in Children cohort study (G40f)
- Standardizing Care to Improve Outcomes in Pediatric Endstage Renal Disease (SCOPE) collaborative (G40g).
- Neonatal Kidney Collaborative (G40h)

Neurology & Neurosurgery (7 points). Hospitals received 1 point for belonging to the Pediatric Brain Tumor Consortium, Children's Oncology Group, or the Pediatric Neuro-Oncology Consortium (H21) and up to 6 additional points for actively enrolling or studying patients in unique, IRB-approved pediatric clinical studies, trials, registries, or databases (H6). Hospitals receive up to 3 points for each NIH funded and non-NIH funded activities as follows: 1 point for 1-9 studies, 2 points for 10-19 studies, or 3 points for 20 or more studies.

Orthopedics (1 point). Hospitals received 1 point for participating in 1 or more IRB-approved trials, studies or databases, such as prospective randomized clinical trials, prospective observational studies or prospective clinical database on patient care (I38).

Pulmonology & Lung Surgery (3 points). Hospitals received 1 point for participating in 1 or more IRB-approved trials, studies or databases, such as prospective randomized clinical trials, prospective observational studies or prospective clinical database on patient care (J51). Hospitals received 2 points for being a member of 4 of the following research networks, or 1 point for being a

member of 1-3 of the following research networks: Children's Interstitial Lung Disease Foundation (J52a); Therapeutics Development Network of the CF Foundation (J52b); certified site for the Severe Asthma Research Program, the Inner City Asthma Consortium (ICAC), American Lung Association (ALA) Airways Clinical Research (ACRC), or the Precision Interventions for Severe Asthma (PrecISE) (J52c); and PCD Foundation Clinical and Research Centers Network (J52d).

Urology (3 points). Hospitals received up to 3 total points for participating in the following prospective research activities: randomized clinical trials (K18a), observational studies (K18b), or clinical databases on patient care (K18c).

Commitment to Quality Improvement (All Specialties)

Hospitals received points in all specialties for participation in quality-improvement activities. Such activities promote internal review and improvement programs and procedures that often lead to improvements in care. The number of points varies by specialty from 15 to 18 points. In all specialties, hospitals could receive up to 7 points for participating in the following quality improvement activities:

- 1 point for publicly reporting performance data on one or more quality metrics (A16 and A16.1);
- Hospitals received up to 2 points for sponsoring quality improvement activities that provide credit to physicians for maintenance of certification (MOC) Part IV:
 - 1 point for being approved by the ABMS as a multispecialty portfolio program (MSPP) sponsor (A17a), or for being approved by the ABP as a pediatric portfolio sponsor (A17b);
 - 1 point for sponsoring one or more projects that are approved by the ABP (A17c).
- 1 point for participating in an external review process for measuring patient/parent satisfaction (A18 and A18.1);
- 1 point for bedside care staff (e.g., nurses, physician assistants, nurse practitioners) participating in quality and safety initiatives (A40);
- Up to 2 points for having a physician serve as a designated Chief Quality/Safety Officer (A41):
 - 2 points for at least .50 FTE
 - 1 point for at least .25 FTE but less than .50 FTE.

In all specialties, hospitals received up to 7 additional points for implementing specialty-specific quality measures (B23/B23.1, C53/C53.1, D25/D25.1, E28/E28.1, F27/F27.1, G11/G11.1, H23/H23.1, I11/I11.1, J45/J45.1, K5/K5.1). These include 1 point each for developing and implementing a written plan for program review and quality improvement, determining appropriate performance-based metrics for clinical quality, tracking and reporting patient data and other supporting information to leadership at least quarterly, presenting results of clinical quality performance metrics to clinical staff at least quarterly, engaging in one or more quality improvement initiatives specific to each specialty, demonstrating how the improvement initiative improved the quality of care, and reporting quality improvement/performance metrics to hospital leadership at least quarterly.

In all specialties except for Gastroenterology & GI Surgery, hospitals received 1 point for participating in national quality and safety collaboratives such as the American College of Surgeons National Surgical Quality Improvement Program (NSQIP) or Children's Hospital Solutions for Patient Safety learning network (A30a-d, A30.1).

In Cancer, hospitals received an additional 2 points (17 points total): 1 point for participating in pediatric cancer-related organized quality improvement network (B23.2) and 1 point for having a pediatric cancer quality committee with an identified medical leader/director that meets at least monthly (B23.3)

In Diabetes & Endocrinology, hospitals received an additional 1 point (16 points total) for supporting development of a physician-led innovation to improve health care delivery for Pediatric Endocrinology patients (C66).

In Gastroenterology & GI Surgery, hospitals received up to 2 additional points (16 points total) for participating in national quality and safety collaboratives or formal initiatives. Hospitals received 1 point for participating in national quality and safety collaboratives such as the Children's Hospital Solutions for Patient Safety learning network (A30b-d, A30.1). Hospitals received 1 point for participating in any formal, multicenter (3 or more institutions) initiatives targeted to GI or liver disorders (D14 and D14.1).

In Neonatology, hospitals received up to 3 additional points (18 points total) if the quality initiatives included having a specified quality-improvement or safety leader and including a parent or family member. Hospitals received 1 point for having a safety leader with less than 0.5 FTE devoted to quality improvement or safety and 2 points for 0.5 FTE or more (F28). Hospitals received 1 point for having a parent/family member of a former NICU patient involved in one or more initiatives as an integral member of the QI/safety team (F28.1).

Congenital Heart Program (Cardiology & Heart Surgery)

In Cardiology & Heart Surgery, hospitals received up to 23 points for having a congenital heart program. Hospitals were rewarded for tracking and reporting data for their congenital heart surgery program and for the volume and type of congenital heart surgeries offered:

- Hospitals received 1 point for having at least one congenital heart surgeon who performed 75 or more congenital heart procedures in the past calendar year or 2 points for having two or more surgeons (E39).
- Hospitals could receive up to 8 points based on the mechanism for determining and reporting volume and outcomes measures. For each of the past four reporting years, hospitals received 2 points each year for reporting to the Society of Thoracic Surgeons (STS) Congenital Heart Surgery Database or 1 point for reporting to another organization (E18).
- Hospitals received 1 point for treating 1 to 4 patients with a Berlin Heart or other ventricular assist device or 2 points for treating 5 or more patients (E26).
- Hospitals received up to 8 points based on the number of cardiac surgical procedures performed in the operating room in the four reporting years: 1 point for 100-249 surgeries/year or 2 points for 250 or more surgeries/year (E38).
- Hospitals received up to 3 points based on the number of neonatal cardiac operations: 1 point for 1-44 operations, 2 points for 45-89 operations, or 3 points for 90 or more operations (E20.1).

ECMO Availability (Neonatology)

In Neonatology, hospitals received up to 2 points for extracorporeal membrane oxygenation (ECMO) services. ECMO technology involves a pump that circulates blood through an artificial lung back into the bloodstream of a very ill neonate, essentially providing heart-lung bypass support outside the child's body.

Hospitals received 1 point for having an ECMO program designated by ELSO as a Center for Excellence (A9).

Hospitals received 1 point for having a specialized multidisciplinary ECMO team with neonatologists managing or co-managing the patient (F14d).

Enlists Families in Structuring Care (All Specialties)

This measure reflects the extent to which a hospital involves parents and families in care. It includes a core set of measures that applied to all pediatric specialties and was worth up to 7 points

in all specialties except Neonatology, in which 8 points were possible. Hospitals received 1 point for having a parent advisory committee that meets one to three times a year or 2 points for having a committee that meets four or more times a year (A14.1).

Hospitals received up to 4 additional points for meeting the following requirements: At least one parent or family member is an active member of the strategic or facility committee (A15a); at least one parent or family member is an active member of one or more standing committees (e.g., quality improvement, patient safety, ethics) (A15b); parents or family members are regularly involved in clinical decision-making through such ways as family-centered rounds, care conferences or other participatory programs (A15c); and parents or family members can participate in family-centered rounds (A15d).

Hospitals received 1 additional point for describing the impact of patients' family members on advisory committees (A15.1).

In Neonatology, hospitals could receive 1 additional point (for a total of 8 points) for having a NICU-specific parent advisory committee that meets at least quarterly (F9).

Has Fulltime Subspecialists Available (All Specialties)

This measure evaluates the presence of a variety of physician specialists, surgeons and dedicated full-time medical staff who are critical to the delivery of appropriate care by pediatric hospitals. *Table 6* identifies the relevant specialists, surgeons and other medical staff for each pediatric specialty. Hospitals received 1 point for each appropriate specialist or surgeon and 1 point for having at least 1.0 FTE of the other medical staff relevant to the specialty.

Table 6. Subspecialists by Specialty

Cancer* (17 points)	Points
Having at least one of each of the following physician specialists: <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist (A4c) • Radiologist specializing in pediatric interventional radiology (A4d) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist or rehabilitation specialist (A4g) • Radiologist specializing in pediatric neuroradiology (A4h) 	7
Having at least one of each of the following pediatric surgeons: <ul style="list-style-type: none"> • Pediatric head and neck surgeon (A5a) • Pediatric general surgeon (A5c) • Pediatric neurosurgeon (A5d) • Pediatric ophthalmology surgeon (A5e) • Pediatric orthopedic surgeon (A5f) • Pediatric urology surgeon (A5g) • Pediatric vascular surgeon (A5j) 	7
Having at least 1.0 FTE of the following other medical staff: <ul style="list-style-type: none"> • Pediatric hematologists/oncologists (B2a) • Other attending on-staff physicians with specific involvement in pediatric cancer program (B2b) • Nurse practitioner and/or physician assistant (B3a and B3b) 	3
Cardiology & Heart Surgery* (22 points)	Points
Having at least one of each of the following physician specialists: <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist specializing in diagnostic radiology (A4c) • Radiologist specializing in pediatric interventional radiology (A4d) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist (A4g) 	6
Having at least one pediatric vascular surgeon (A5j)	1
At least 2.0 FTE of the following staff: <ul style="list-style-type: none"> • Pediatric cardiothoracic surgeon (E2a) • Pediatric cardiac intensivists (cardiologists, pediatric critical care or anesthesiologists) or pediatric radiologists (E2b, E2c, E2d, or E2h) • Pediatric cardiac interventionalists (E2e) 	3
At least 1.0 FTE of the following staff: <ul style="list-style-type: none"> • Pediatric cardiac electrophysiologist (E2f) • Anesthesiologist with pediatric training/experience (E2g) • Cardiologist with subspecialty certification in adults with congenital heart disease (E2i) • Clinical nurse, advanced registered nurse practitioner, advanced practice registered nurse, or physician assistant (E4a, E4b, and E4c) 	4

* Parenthetical references indicate related survey questions

(continued)

Table 6. Subspecialists by Specialty (continued)

Cardiology & Heart Surgery, continued* (21 points)	Points
<p>Up to 2 points for 24/7 in-house coverage of the cardiac ICU:</p> <ul style="list-style-type: none"> • 2 points if staffed with pediatric intensivists; pediatric cardiology, pediatric cardiac intensive care or pediatric cardiac surgery trainees; or non-physician advanced practice providers (APPs) dedicated to cardiac intensive care management (E3.1) • 1 point if staffed with other medical staff (E3.1 and E3.2) 	2
<p>Up to 3 points for the type of 24-hour in-house coverage provided every day to the <u>cardiac-specific ICU (E3)</u>:</p> <ul style="list-style-type: none"> • 3 points for having a dedicated Cardiac ICU (CICU) • 2 points for having a dedicated section of a Pediatric ICU (PICU) and/or Neonatal ICU (NICU) • 1 point for having a blended Pediatric ICU (PICU) and/or Neonatal ICU (NICU) without a dedicated section 	3
<p>Having eligible RNs working in the CICU (or dedicated beds in the PICU) meet the following thresholds:</p> <ul style="list-style-type: none"> • Less than 20% with less than 2 years of cardiac critical care experience (E4d) • At least 80% with a BSN or higher degree (E4e) • At least 10% with CCRN certification for critical care nursing from the AACN (E4f) 	3
Diabetes & Endocrinology* (14 points)	Points
<p>Having at least one of each of the following physician specialists:</p> <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist specializing in diagnostic radiology (A4c) • Radiologist specializing in pediatric interventional radiology (A4d) • Pediatric rheumatologist (A4e) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist (A4g) 	7
<p>Having at least one of each of the following pediatric surgeons:</p> <ul style="list-style-type: none"> • Pediatric head and neck surgeon (A5a) • Pediatric general surgeon (A5c) • Pediatric neurosurgeon (A5d) 	3
<p>At least 1.0 FTE of the following staff:</p> <ul style="list-style-type: none"> • Pediatric endocrinologist (C2a) • Nurse practitioner and/or physician assistant (C3) • Bachelor’s-level registered nurse dedicated to outpatient care (C4a) • Master’s or doctorate-level registered nurse dedicated to outpatient care (C4b, C4c) 	4

* Parenthetical references indicate related survey questions

(continued)

Table 6. Subspecialists by Specialty (continued)

Gastroenterology & GI Surgery* (11 points)	Points
Having at least one of each of the following physician specialists: <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist specializing in diagnostic radiology (A4c) • Radiologist specializing in pediatric interventional radiology (A4d) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist (A4g) 	6
Having at least one pediatric general surgeon (A5c)	1
Having at least 1.0 FTE of the following other medical staff: <ul style="list-style-type: none"> • Pediatric gastroenterologist (D2a) • Pediatric surgeon available 7 days a week (D2.2) • Nurse practitioner and/or physician assistant (D3) • Dedicated social workers (D3.1a), dedicated psychologists (D3.1b), dedicated dietitians (D3.1c), or patient care coordinators (D3.1d) 	4
Neonatology* (18 points)	Points
Having at least one of each of the following physician specialists: <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist specializing in diagnostic radiology (A4c) • Radiologist specializing in pediatric interventional radiology (A4d) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist (A4g) • Radiologist specializing in pediatric neuroradiology (A4h) 	7
Having at least one of each of the following pediatric surgeons: <ul style="list-style-type: none"> • Pediatric head and neck surgeon (A5a) • Pediatric cardiothoracic surgeon (A5b) • Pediatric general surgeon (A5c) • Pediatric neurosurgeon (A5d) • Pediatric ophthalmology surgeon (A5e) • Pediatric orthopedic surgeon (A5f) • Pediatric urology surgeon (A5g) • Pediatric plastic surgeon (A5h) 	8
Having at least 1.0 FTE of the following other medical staff: <ul style="list-style-type: none"> • Pediatric neonatologist (F2a) • Clinical registered nurse (F4a) 	2
Having at least 1 physician extender (F3)	1

* Parenthetical references indicate related survey questions

(continued)

Table 6. Subspecialists by Specialty (continued)

Nephrology* (9 points)	Points
Having at least one of each of the following physician specialists: <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist specializing in diagnostic radiology (A4c) • Radiologist specializing in pediatric interventional radiology (A4d) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist (A4g) 	6
Having at least one pediatric general surgeon (A5c)	1
Having at least 1.0 FTE of the following other medical staff: <ul style="list-style-type: none"> • Pediatric nephrologist (G2a) • Nurse practitioner and/or physician assistant (G3) 	2
Neurology & Neurosurgery* (15 points)	Points
Having at least one of each of the following physician specialists: <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist specializing in diagnostic radiology (A4c) • Radiologist specializing in pediatric interventional radiology (A4d) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist (A4g) • Radiologist specializing in pediatric neuroradiology (A4h) 	7
Having at least one of each of the following pediatric surgeons: <ul style="list-style-type: none"> • Pediatric general surgeon (A5c) • Pediatric neurosurgeon (A5d) • Pediatric vascular surgeon (A5j) 	3
Having at least 1.0 FTE of the following other medical staff: <ul style="list-style-type: none"> • Pediatric neurologist (H2a) • Pediatric neurosurgeon (H2b) • Nurse practitioner and/or physician assistant (H3) 	3
Having at least 1.0 FTE of nurses with advanced neurologic certification (H4)	1
Having at least 1.0 FTE of staff who support Ketogenic Diet planning and implementation with patients (H5.1)	1
Orthopedics* (22 points)	Points
Having at least one of each of the following physician specialists: <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist specializing in diagnostic radiology (A4c) • Pediatric radiologist specializing in interventional radiology (A4d) • Pediatric rheumatologist (A4e) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist (A4g) 	7

* Parenthetical references indicate related survey questions

(continued)

Table 6. Subspecialists by Specialty (continued)

Orthopedics*, continued (22 points)	Points
Having at least one of each of the following pediatric surgeons: <ul style="list-style-type: none"> • Pediatric general surgeon (A5c) • Pediatric orthopedic surgeon (A5f) • Pediatric plastic surgeon (A5h) • Pediatric hand surgeon (A5i) • Pediatric vascular surgeon (A5j) • Pediatric orthopedic surgery resident (I6.1b) 	6
Having at least one of each of the following specialists: <ul style="list-style-type: none"> • Hand surgery (I6a) • Spinal surgery (I6b) • Musculoskeletal oncologist (I6c) • Orthopaedic sports surgeons (I6d) • Hip preservation specialist (I6e) • Musculoskeletal radiologist (I6f) 	6
Having at least 1.0 FTE of the following other medical staff: <ul style="list-style-type: none"> • Pediatric orthopedic surgeon (I2a) • Nurse practitioner and/or physician assistant (I3) • Dedicated clinical registered nurses or medical assistants (I4) 	3
Pulmonology & Lung Surgery* (12 points)	Points
Having at least one of each of the following physician specialists: <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist specializing in diagnostic radiology (A4c) • Radiologist specializing in pediatric interventional radiology (A4d) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist (A4g) 	6
Having at least one of each of the following pediatric surgeons: <ul style="list-style-type: none"> • Pediatric general surgeon (A5c) • Pediatric vascular surgeon (A5j) 	2
Having at least 1.0 FTE of the following other medical staff: <ul style="list-style-type: none"> • Pediatric pulmonologist (J2a) • Pediatric sleep medicine physician (J2b) • Nurse practitioner and/or physician assistant (J3) • Clinical registered nurse (J4) 	4

* Parenthetical references indicate related survey questions

(continued)

Table 6. Subspecialists by Specialty (continued)

Urology* (13 points)	Points
Having at least one of each of the following physician specialists: <ul style="list-style-type: none"> • Pediatric anesthesiologist (A4a) • Pediatric critical care specialist (A4b) • Pediatric radiologist specializing in diagnostic radiology (A4c) • Pediatric radiologist specializing in interventional radiology (A4d) • Pediatric infectious disease specialist (A4f) • Pediatric physiatrist (A4g) 	6
Having at least one of each of the following pediatric surgeons: <ul style="list-style-type: none"> • Pediatric general surgeon (A5c) • Pediatric urology surgeon (A5g) • Pediatric plastic surgeon (A5h) 	3
Having at least 1.0 FTE of the following other medical staff: <ul style="list-style-type: none"> • Pediatric urologist (K2a) • Nurse practitioner and/or physician assistant (K3) • Clinical registered nurse (K4) 	3
Having an in-house ultrasound technologist to support prompt imaging and diagnosis of suspected testicular torsion (K20)	1

* Parenthetical references indicate related survey questions

Heart Transplant Program (Cardiology & Heart Surgery)

In Cardiology & Heart Surgery, hospitals received up to 11 points for having a heart transplant program. Hospitals received 1 point for having an onsite heart or heart-lung transplant program recognized by the United Network for Organ Sharing (UNOS) (E21). Hospitals received up to 3 points based on the number of unique patients who received heart transplants in the past 4 years combined (E22): 1 point for 1-7 transplants, 2 points for 8-15 transplants and 3 points for 16 or more transplants. Three additional points were awarded based on the number of patients < 1 year of age who received heart transplants (E22.1): 1 point for 1-4 patients, 2 points for 5-9 patients, and 3 points for 10 or more patients. Hospitals also received 1 point for having performed cardiac transplantation in a recipient with high ($\geq 10\%$) panel reactive antibody (PRA) (E25a), 1 point for having a written protocol for the management of recipients with high ($\geq 10\%$) PRA (E25b), 1 point for having performed an ABO-incompatible heart transplant (E25c), and 1 point for having a written protocol for the management of ABO incompatible recipients (E25d).

Help for Families (All Specialties)

The Patient and Family Services measure evaluates access to medical specialists and services. A core set of sub-measures for all specialties is worth up to 8 points, which includes providing direct access to certified child life specialists (A12a), family-support specialists (A12b), pediatric

psychologists or psychiatrists (A12c), a family resource center (A13a), sleep rooms for parents or siblings (A13b), a school intervention program (A13c), and a Ronald McDonald House (or other residential facility) (A13d). Hospitals could also receive 1 additional point for having direct access to interpreter services^{***} either through having access to in-person interpreters, interpreters through electronic means, or both (A12.1).

In Neonatology, hospitals could receive up to 8 additional points (for a total of 16 points). Hospitals received points for offering the following patient and family services: NICU-specific family psychosocial support program (F8a), 24/7 parental visitation (F8b), sibling visitation (F8c), NICU-specific parent-to-parent support groups (F8d), designated psychologists or psychiatrists available for referrals and consultations with parents (F8e), Child Life support team available to NICU families (F8f), NICU-dedicated multidisciplinary developmental care team (F8g), and complex discharge coordinator (F8h).

In Nephrology, hospitals could receive up to 6 additional points (for a total of 14 points). Hospitals received 1 point for offering summer camp for kidney transplant patients (G33b). Hospitals received up to an additional 4 points for offering the following programs to support patients in a pediatric maintenance dialysis program: teachers dedicated to working with patients (G9a), a standard review of school performance and patient's Individualized Education Program (G9b), summer camp (G9c), quality of life assessment (G9d), and Child Life specialists (G9e).

Liver Transplant Program (Gastroenterology & GI Surgery)

In Gastroenterology & GI Surgery, hospitals received up to 5 points for having a liver transplant program. Hospitals received 1 point for having a UNOS-recognized liver transplant program (D20), 1 point for having at least 1 transplant hepatologist (D19.1) and up to 3 points based on the number of unique patients who received a liver transplant in the past 5 years (D21a and D22a): 1 point for 1-9 patients, 2 points for 10-19 patients, or 3 points for 20 or more patients.

Lung Transplant Program (Pulmonology & Lung Surgery)

In Pulmonology & Lung Surgery, hospitals received up to 5 points for having a lung transplant program. Hospitals received 1 point for offering a UNOS-recognized lung transplant program (J46). Hospitals received 1 point for performing one lung transplant between January 2014 and June 2016 or 2 points for performing two or more lung transplants between January 2014 and June 2016 (J48a). Hospitals received 1 point for performing one lung transplant between July 2016

^{***} For in-person interpreter services having at least 50% certified through the National Board of Certification for Medical Interpreters or the Certification Commission for Healthcare Interpreters.

and December 2018 or 2 points for performing two or more lung transplants between July 2016 and December 2018 (J47a).

Neonatal Transport (Neonatology)

In neonatology, hospitals received up to 12 points for ensuring the safety of newborns during transport. Hospitals received up to 5 points for providing a neonatal-specific transport team with each of the following:

- A medical director board-certified in Neonatal-Perinatal Medicine (F13a)
- At least 2 clinicians (e.g., RN, RT, MD, DO, NNP, PA) on each transport who are non-drivers (F13b)
- All RN's and RT's have at least 1 year of NICU level III or IV experience (F13c)
- Neonatal transport team is immediately available 24/7 to respond to emergent neonatal transports (F13d)
- Active servo-controlled cooling on transport for term and near term infants with moderate to severe hypoxic ischemic encephalopathy (F13e).

Hospitals received 1 point for tracking temperature at admission for infants actively cooled during transport for the management of moderate to severe hypoxic ischemic encephalopathy (F13.1). Hospitals received an additional 1 point if data on active cooling was systematically collected and reported to a benchmarking registry (e.g., CHNC) or for internal process improvement work (F13.3).

Hospitals received 1 point if the NICU has the capability of providing inhaled nitric oxide therapy during transport with high-risk pre-ECMO patients whenever indicated (F13.4).

Hospitals received 1 point for tracking time to vehicular dispatch for neonatal transport (F13.5). Hospitals received 2 additional points for having $\geq 80\%$ of neonatal transports dispatched within 30 minutes of the call being logged as received or 1 point for having ≥ 40 and $< 80\%$ dispatched within 30 minutes (F13.6). Hospitals received an additional 1 point if data on emergent neonatal transport was systematically collected and reported to a benchmarking registry (e.g., CHNC) or for internal process improvement work (F13.7).

Nurse Staffing (All Specialties)

This measure is a relative ratio of the number of nurses to the average daily patient census. The numerator is the number of on-staff registered nurses (RNs) hospital-wide (other than in

Neonatology) who are dedicated to inpatient pediatric clinical care, expressed as FTEs (A2)†††. Nurses must have an RN degree from an approved nursing school and hold a current state license. The denominator is the average daily number of pediatric inpatients (A1)‡‡‡. The source was the Pediatric Hospital Survey. In Neonatology, the measure counted only nurses dedicated specifically to the NICU (F4a) and the average daily census comprised only NICU patients (F6). For scoring purposes, nurse-patient values were capped at 4.0 in all specialties to prevent skewness.

Percent of Dialysis Patients Who Had Transplants (Nephrology)

Hospitals received up to 6 points in Nephrology based on the percentage of end-stage renal disease (ESRD) patients with a completed CMS-2728 (Medical Evidence) form receiving hemodialysis or peritoneal dialysis who were over 10 kg (G20.1) who received kidney transplants within the past 2 years (G21). Patients were evaluated separately by age group: children under 5 and children aged 5-19. For each age group, hospitals received up to 3 points for having a higher percentage of patients receiving transplants as follows: 1 point if $\geq 20\%$ and $< 40\%$, 2 points if $\geq 40\%$ and $< 60\%$, or 3 points if $\geq 60\%$.

Provides Advanced Palliative Care Program (Cancer)

In Cancer, hospitals received up to 8 points for palliative care. Hospitals could receive up to 4 points for offering the following pain control programs: patient-controlled analgesia (B29a), nurse-controlled analgesia (B29b), pediatric pain service consults (B29c), and pediatric outpatient pain management services (B29d).

Hospitals received 1 point for offering a qualified palliative care program onsite (B29.1). A qualified program is defined as one that is organized and staffed for children nearing the end of life or living with conditions that limit lifespan or quality of life. It is intended to minimize pain and discomfort, provide emotional and spiritual support for children and their families, assist with financial guidance and social services and support decision-making. The program must include at least one physician providing direct patient care as well as a nurse coordinator and either a social worker, certified child life specialist or pastoral counselor, and all staff must have training in palliative care.

Hospitals received 1 point for having at least 1 physician board-certified in Hospice and Palliative Medicine (B29.2).

††† Hospital are directed to calculate FTEs based on total paid hours for the period of review divided by 2080.

‡‡‡ This includes inpatient days divided by the number of days that the hospital was open (e.g. 365); hospitals are directed to include all patients admitted to the hospital, including short stays and observation stays.

Hospitals could receive up to 2 points based on the percentage of patients with advanced and refractory cancer who were referred to the palliative care program (B30): 1 point for $\geq 50\%$ or $< 75\%$ or 2 points for $\geq 75\%$.

Recognized as Nurse Magnet Hospital (All Specialties)

The Nurse Magnet status measure is a formal designation by the Magnet Recognition Program®. The program was developed by the American Nurses Credentialing Center (ANCC) to recognize healthcare organizations that meet certain quality indicators on specific standards of nursing excellence. The list of Magnet-recognized facilities is updated throughout the year as organizations apply for designation and re-designation status. Hospitals received credit based on their Magnet Recognition status as of March 13, 2020. The current list of Magnet-recognized organizations is shown at <http://www.nursecredentialing.org/Magnet/FindaMagnetFacility>.

Hospitals received 1 point for being recognized as a Nurse Magnet hospital. For children's hospitals that are part of a special merger^{§§§} or a multiplex healthcare system, the primary hospital is required to have Magnet Recognition status for the combination hospital to receive 1 point. If there is no defined primary hospital, then if either hospital in the special merger has Magnet Recognition status then both receive credit.

Specialized Clinics and Programs (Cancer, Diabetes & Endocrinology, Gastroenterology & GI Surgery, Neonatology, Neurology & Neurosurgery, Orthopedics, Urology)

Cancer (6 points). Hospitals received 1 point for each of the following specialized treatment programs: clinical brain tumor program (B10a), solid tumor program (B10b), clinical leukemia/lymphoma program (B10c), comprehensive longer-term survivors program (B10d), histiocytosis program (B10e), and molecular oncology/targeted therapy program (B10f).

Diabetes & Endocrinology (22 points). Hospitals received up to 2 points for specialized treatment programs for endocrine patients. Hospitals received one point for having the following onsite programs and an additional point for each program if pediatric endocrinologists regularly attended the program: lipid disorders (C46a), hypertension (C46b), comprehensive weight management (C46c), Turner syndrome (C46d), cystic fibrosis-related diabetes (C46e), gender dysphoria (C46f), disorders of sexual development (C46g), thyroid nodules (C46h), 22q11.2 Deletion Syndrome (C46i), Muscular Dystrophy (C46j) or Prader Willi Syndrome (C46k).

^{§§§} In a special merger, two separate hospitals operate as one and their data are combined for analysis. Boston Children's Hospital and Dana-Farber Cancer Center are an example in pediatric Cancer. Specialty or secondary hospitals that are combined with the primary hospital are noted on the US News website for that hospital.

Gastroenterology & GI Surgery (12 points). Hospitals received 1 point for each of the following interdisciplinary treatment programs for gastrointestinal disorders with at least 10 patients in the last calendar year: intestinal rehabilitation (D10a), cystic fibrosis treatment (D10b), total parenteral nutrition (TPN) (D10c), aerodigestive (D10j), pancreatic disease (D10k), and anorectal or colorectal program (D10l). Hospitals received 1 point for each of the following interdisciplinary treatment programs for gastrointestinal disorders with at least 20 patients in the last calendar year: pediatric intensive feeding (D10d), multidisciplinary childhood obesity management (D10e), inflammatory bowel disease (D10f), multidisciplinary allergic gastrointestinal disease (D10g), chronic liver disease (D10h), and neurogastrointestinal (D10i).

Neonatology (17 points). Hospitals received 1 point for having a cardiac ICU to care for newborn infants (<28 days) that need specialized care for heart conditions (F17), and up to 16 additional points for providing specialized treatment teams or clinics to deal with particularly challenging conditions. Hospitals received 1 point for each of the following: craniofacial team (F14a), spina bifida team (F14b), comprehensive retinopathy of prematurity program (F14c), neonatal-neurointensive care program (F14e), palliative care program (F14f), micrognathia team (F14g), chronic lung disease team (F14h), congenital diaphragmatic hernia team (F14i), chronic pulmonary hypertension team (F14j), neonatal dialysis team (F14k), multidisciplinary team for follow-up with congenital diaphragmatic hernia patients after discharge (F14l), metabolic team (F15a), bowel rehabilitation team (F15b), home ventilator management team (F15c), neurodevelopmental follow-up clinic for premature/high-risk NICU patients (F15d) and neurodevelopmental clinic for high-risk congenital heart neonatal patients (F15e).

Neurology & Neurosurgery (19 points). Hospitals received up to 20 points for access to specialized treatment clinics or programs for pediatric neurological disorders. To receive credit, a hospital had to have an organized program that included a medical director and nursing coordinator. One point was awarded for each of the following multidisciplinary clinics or programs: cerebral palsy/spasticity (H12a), cerebrovascular/stroke multidisciplinary (H12b), craniofacial surgical (H12c), surgical movement disorders (H12d), neurofibromatosis (H12e), neuromuscular (H12f), neuro-oncology (H12g), spina bifida (H12h), tuberous sclerosis (H12i), brachial plexus (H12j), genetic metabolic (H12k), neonatal neurology (H12l), head trauma/post-concussion (H12m), new-onset seizures (H12n), diagnostic neuro-fetal (H12o), headache (H12p), pain (H12q), demyelinating disorders (H12r), and autism/neurodevelopmental disorders (H12s).

Orthopedics (12 points). Hospitals received up to 12 points for providing specialized treatment clinics or programs to treat significant conditions. To receive credit, the clinic had to be attended regularly by the pediatric orthopedic service and see a minimum of 25 patients in the last calendar year. Hospitals received 1 point for each of the following clinics or programs: spina bifida

(I15a), spasticity (I15b), skeletal dysplasia (I15c), brachial plexus (I15d), neurofibromatosis (I15e), muscle disease (I15f), pain (I15g), sports medicine (I15h), sports concussion program (I15i), arthrogyposis (I15j), limb deficiency/limb reconstruction/prosthetics (I15k), and skeletal health/metabolic bone health (I15l).

Urology (10 points). Hospitals received 1 point for each of the following specialized treatment clinics or programs to treat significant urological conditions or issues: spina bifida (K10a), comprehensive stone program (K10b), prenatal program (K10c), differences in sex development program (K10d), exstrophy/cloaca/GU sinus program (K10e), end stage renal disease program for patients who require a kidney transplant (K10f), and fertility preservation/oncofertility program (K10g). Hospitals also received 1 point for offering a Transitional Care Program for patients transitioning to adult urology with congenital conditions affecting the genitourinary (GU) system (K10.1). Hospitals received an additional 1 point for being an integrated member of the pediatric oncology treatment program (K10.3). Hospitals received an additional 1 point for providing a voiding dysfunction program that treated at least 120 patients in the last calendar year (K17).

Success in Helping Patients Manage Their Asthma (Pulmonology & Lung Surgery)

In Pulmonology & Lung Surgery, hospitals received up to 13 points for management of asthma patients. Hospitals received up to 4 points for their treatment of severe asthma patients: hospitals received 1 point for having a program with dedicated staff (at least 0.5 FTE) to identify and treat patients with severe, high-risk, difficult-to-control or life-threatening asthma (J7), 1 point for having a written protocol for evaluation of patients with severe asthma (J8), 1 point for monitoring medication adherence in severe asthma patients (J8.1), and 1 point for having access to at least 1.0 FTE of Certified Asthma Educators in the last calendar year (J8.2).

Hospitals received up to 9 additional points based on the percentage of patients with a primary diagnosis of asthma following 3 specific protocols. The protocols evaluated were as follows: providing eligible outpatients in subspecialty care clinics with a documented assessment of asthma control (e.g., ACT, ATAQ) (J10e/J10d), completing an outpatient follow-up visit within 90 days of discharge (J10c/J10b), and successfully managing outpatients so that they were not admitted for care related to their asthma (J10b/J10a^{****}). For the first two protocols, up to 3 points were awarded for the percentage of patients following the protocol: 1 point for $\geq 50\%$ and $< 75\%$, 2 points for $\geq 75\%$ and $< 90\%$ or 3 points for $\geq 90\%$. For the last protocol, points were awarded for a *lower*

**** This survey item was reverse scored to reward hospitals for having FEWER outpatients admitted for asthma-related care.

percentage of patients being admitted for care related to their asthma: 3 points for $\leq 10\%$ of patients admitted, 2 points for $>10\%$ and $\leq 25\%$ or 1 point for $>25\%$ and $\leq 50\%$.

Success in Managing Neuromuscular Weakness Disorder (Pulmonology & Lung Surgery)

In Pulmonology & Lung Surgery, hospitals received up to 3 points for the percentage of muscular dystrophy patients who had pulmonary function testing in the past calendar year (J30). Points were based on the percentage of patients as follows: 1 point for $\geq 50\%$ and $< 75\%$, 2 points for $\geq 75\%$ or $< 90\%$, and 3 points for $\geq 90\%$.

Tracking Growth Metrics for Treated Patients (Neonatology)

Hospitals received up to 7 points in Neonatology for recording growth metrics on infants within 7 days prior to discharge or transfer, including weight (F40a), length (F40b) and head circumference (F40c). For each of the 3 growth metrics, points were awarded as follows: 1 point for recording metrics on $\geq 60\%$ and $< 90\%$ of infants; or 2 points for recording metrics on $\geq 90\%$ of infants. Hospitals received an additional 1 point for measuring infant length using a length board (F41).

Volume of Patients (All Specialties)

Unless noted otherwise, volume measures indicate the number of unique patients in the past calendar year who had the specified diagnoses or conditions or who received the specified procedures or treatments. If data were unavailable for the most recent calendar year, hospitals were instructed to use data from the most recent 12 months data were available.

Low-, medium- and high-volume categories were created for most measures, based on the distribution of volume across all hospitals. For other measures, categories were based on conceptual thresholds for the number of patients or procedures needed to indicate a sufficient level of experience. We assigned points based on categories rather than on continuous values to ensure that one or two hospitals with extremely high volumes did not skew scoring. Hospitals that had zero volume or that did not respond received 0 points. For almost all measures, hospitals in the lowest-volume category received 1 point, medium-volume hospitals received 2 points and high-volume hospitals received 3 points. An exception is two of the items in the Number of Patients in Orthopedics which received 2 points, 4 points or 6 points respectively for low, medium or high volume. The increased points reflects these items' increased importance relative to other items in the measure.

In addition, for items with extremely low volume, such as GI and urological surgical procedures, the measure was divided only into low and medium volumes, with a maximum of 2 points. *Table 7* identifies the volume measures used by specialty and the points assigned to volume scores within a certain range.

Table 7. Volume Measures by Specialty

Cancer Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of new patients, 2 years (B6) (max points = 3)	1-99	100-399	400+
Number of patients (max points = 30)			
• Leukemia (B27a1)	1-149	150-299	300+
• Brain tumors/Central Nervous System (B27b1)	1-99	100-199	200+
• Neuroblastoma (B27c1)	1-24	25-49	50+
• Bone tumors (B27d1)	1-19	20-39	40+
• Soft tissue sarcomas (B27e1)	1-19	20-39	40+
• Wilms' tumor (B27f1)	1-14	15-29	30+
• Liver tumors (B27g1)	1-5	6-11	12+
• Retinoblastoma (B27h1)	1-5	6-11	12+
• Germ cell tumors (B27i1)	1-7	8-15	16+
• Lymphoma (B27j1)	1-29	30-59	60+
Number of surgeries** (B27), (max points = 16)			
• Brain tumors (B27b2)	1-29	30+	NA
• Neuroblastoma (B27c2)	1-4	5+	NA
• Bone tumors (B27d2)	1-9	10+	NA
• Soft tissue sarcomas (B27e2)	1-9	10+	NA
• Wilms' tumor (B27f2)	1-4	5+	NA
• Liver tumors (B27g2)	1-3	4+	NA
• Retinoblastoma (B27h2)	1-2	3+	NA
• Germ cell tumors (B27i2)	1-3	4+	NA

* Parenthetical references indicate related survey questions.

** Volume represents procedures, not patients.

(continued)

Table 7. Volume Measures by Specialty (continued)

Cardiology & Heart Surgery Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of catheter procedures** (max points = 18)			
• Diagnostic catheterization only (hemodynamic and/or angiographic evaluation) (E11b, E12b)	1-99	100-199	200+
• Endomyocardial biopsy (with or without a diagnostic procedure and without an additional intervention) (E11c, E12c)	1-59	60-119	120+
• Catheter ablation procedures (E11d, E12d)	1-4	5-9	10+
• Interventional procedure (with or without a diagnostic procedure or biopsy) (E11e, E12e)	1-199	200-399	400+
• Transcatheter pulmonary valve replacements (E11f, E12f)	1-4	5-9	10+
• Placement of permanent transvenous pacing (E15)	1-19	20-39	40+
Number of Norwood or hybrid surgeries (max points = 12)			
• Patients receiving hybrid or Norwood Stage 1, year 1 (E40a)	1-6	7-13	14+
• Patients receiving hybrid or Norwood Stage 1, year 2 (E40b)	1-6	7-13	14+
• Patients receiving hybrid or Norwood Stage 1, year 3 (E40c)	1-6	7-13	14+
• Patients receiving hybrid or Norwood Stage 1, year 4 (E40d)	1-6	7-13	14+
Number of surgeries*** (max points = 12)			
• STAT ^{††††} Level 2: Years 1-4 (E42)	1-299	300-599	600+
• STAT Level 3: Years 1-4 (E42)	1-149	150-299	300+
• STAT Level 4: Years 1-4 (E42)	1-149	150-299	300+
• STAT Level 5: Years 1-4 (E42)	1-59	60-119	120+

* Parenthetical references indicate related survey questions. (continued)

** Volume represents procedures, not patients. For the first five metrics (E11a-f, E12a-f), volumes from NCDR report are used if available for all quarters in the most recent year.

*** Volume represents procedures, not patients.

†††† Society of Thoracic Surgery & European Association for Cardio-Thoracic Surgery Congenital Heart Surgery Mortality Categories (STAT)

Table 7. Volume Measures by Specialty (continued)

Diabetes & Endocrinology Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of patients (max points = 36)			
• Type 1 diabetes outpatient visits (C28.1a)	1-499	500+	NA
• Type 2 diabetes outpatient visits (C28.1b)	1-249	250+	NA
• Diabetes-related care admissions for Type 1 patients (C28.1c)	1+	NA	NA
• Diabetes-related care admissions for Type 2 patients (C28.1d)	1+	NA	NA
• Patients with a genetically confirmed form of genetic diabetes MODY or NDM on private insurance (C28.2a)	1-9	10+	NA
• Patients with a genetically confirmed form of genetic diabetes MODY or NDM on Medicaid insurance (C28.2b)	1-9	10+	NA
• Congenital adrenal hyperplasia (C47a)	1-39	40+	NA
• CNS and endocrine tumors (C47b)	1-99	100+	NA
• Diabetes insipidus (C47c)	1-24	25+	NA
• Hypopituitarism (C47d)	1-99	100+	NA
• Turner Syndrome (C47e)	1-24	25+	NA
• Noonan Syndrome (C47f)	1-24	25+	NA
• Gender dysphoria (C47g)	1-24	25+	NA
• Disorders of sexual development (C47h)	1-24	25+	NA
• Bone disease (including metabolic and genetic conditions) (C47i)	1-24	25+	NA
• Non-diabetes related hypoglycemia (C47j)	1-99	100+	NA
• Polycystic ovarian syndrome (C47k)	1-79	80+	NA
• Nondiabetes endocrine disorders outpatients (C57a1)	1-1,999	2,000+	NA
• Nondiabetes endocrine disorders inpatients (C57b1)	1-124	125+	NA

* Parenthetical references indicate related survey questions.

** Volume represents procedures, not patients.

(continued)

Table 7. Volume Measures by Specialty (continued)

Diabetes & Endocrinology, continued Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of patients undergoing procedures** (max points = 14)			
• Ratio of patients with growth hormone deficiency who received a brain or pituitary MRI (2 years) (C49a/C48)	50%-74.9%	75%+	NA
• Ratio of patients with growth hormone deficiency who received a growth hormone therapy (C49b/C48)	50%-74.9%	75%+	NA
• Ratio of patients with growth hormone deficiency who received a serum IGF-1 measurement (C49c/C48)	50%-74.9%	75%+	NA
• Thyroid cancer surgery (C51.1a)	1-3	4+	NA
• Parathyroid surgery (C51.1b)	1	2+	NA
• Brain tumor surgery involving hypothalamus or pituitary (C51.1c)	1-3	4+	NA
• Abdominal endocrine surgery (C51.1d)	1	2+	NA
Gastroenterology & GI Surgery Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of noninvasive procedures** (max points = 18)			
• Capsule endoscopy (D11a)	1-19	20+	NA
• Endoscopic band ligation/sclerotherapy (D11b)	1-4	5+	NA
• Esophageal impedance or high resolution esophageal manometry (D11c)	1-49	50+	NA
• Endoscopic retrograde cholangiopancreatography (D11d)	1-19	20+	NA
• Antroduodenal and full colonic motility studies (D11e)	1-4	5+	NA
• Esophageal dilation (D11f)	1-29	30+	NA
• Alternative hemostasis therapies (D11g)	1-7	8+	NA
• Deep enteroscopy-single or double balloon (D11h)	1-3	4+	NA
• Endoscopic ultrasound (D11i)	1-3	4+	NA

* Parenthetical references indicate related survey questions.

** Volume represents procedures, not patients.

(continued)

Table 7. Volume Measures by Specialty (continued)

Gastroenterology & GI Surgery, continued Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of patients (max points = 27)			
• Pseudo-obstruction (D13a)	1-12	13-24	25+
• Chronic intestinal failure patients who require TPN for 2 months or more (D13b)	1-39	40-79	80+
• Chronic liver disease (D13c)	1-99	100-199	200+
• Acute recurring or chronic pancreatitis (D13d)	1-49	50-99	100+
• Biliary atresia (D13e)	1-14	15-29	30+
• Portal hypertension (D13f)	1-19	20-39	40+
• Celiac disease (D13g)	1-99	100-199	200+
• Eosinophilic esophagitis (D13h)	1-99	100-199	200+
• Congenital anorectal or colorectal disorders (such as Hirschsprung disease, Imperforate anus, and Cloaca) (D13i)	1-49	50-99	100+
Number of patients undergoing surgeries (max points = 12)			
• Hepatopertoenterostomy or Kasai procedure on a patient with biliary atresia (D17a)	1	2+	NA
• Laparoscopic gastrointestinal, hepatic, and pancreatic surgery (D17b)	1-24	25+	NA
• Bariatric surgery (D17c)	1-4	5+	NA
• Posterior sagittal anorectoplasties (D17d)	1-9	10+	NA
• Open and laparoscopic abdominal surgeries for inflammatory bowel disease (IBD) (D17e and D17f)	1-11	12+	NA
• Esophageal atresia repair (D17g)	1-7	8+	NA

* Parenthetical references indicate related survey questions.

(continued)

** Volume represents procedures, not patients.

Table 7. Volume Measures by Specialty (continued)

Neonatology Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of patients (max points = 33)			
• Congenital diaphragmatic hernia (F16a)	1-14	15-29	30+
• Hirschsprung's disease (F16b)	1-11	12-23	24+
• Hypothermia treatment and either hypoxic ischemic encephalopathy or severe birth asphyxia (F16c)	1-49	50-99	100+
• Open neural tube defect treatment (F16d)	1-14	15-29	30+
• Gastroschisis (F16e)	1-17	18-35	36+
• Tracheoesophageal fistula (F16f)	1-15	16-31	32+
• Omphalocele (F16g)	1-9	10-19	20+
• Duodenal atresia (F16h)	1-11	12-23	24+
• Anorectal malformation (F16i)	1-19	20-39	40+
• Extracorporeal life support therapy (F16j)	1-14	15-29	30+
• Jejunal atresia or ileal atresia (F16k)	1-9	10-19	20+
Nephrology Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of dialysis patients, 2 years (max points = 12)			
• End-stage renal disease (ESRD) patients < 5 years of age on hemodialysis or peritoneal dialysis (G20a)	1-4	5-9	10+
• ESRD patients 5-19 years of age on hemodialysis or peritoneal dialysis (G20b)	1-14	15-29	30+
• Dialysis treatment volume in days (previous year) (G8a)	1-249	250-499	500+
• Dialysis treatment volume in days (current year) (G8b)	1-249	250-499	500+
Number of kidney biopsies, 2 years (max points = 6)			
• Native kidney percutaneous biopsies (G14a)	1-25	26-75	76+
• Percutaneous kidney transplant biopsies (G27)	1-19	20-39	40+
Number of kidney transplants (max points = 6)			
• Deceased-donor kidney transplant patients (G32.1a1 and G32.2a1)	1-8	9-17	18+
• Living-donor kidney transplant patients (G32.1a2 and G32.2a2)	1-7	8-16	17+

n/a = not applicable.

* Parenthetical references indicate related survey questions.

** Volume represents procedures, not patients.

(continued)

Table 7. Volume Measures by Specialty (continued)

Nephrology, continued Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of patients, 2 years (max points = 14)			
• Acute kidney injury (G6)	1-199	200-399	400+
• Systemic lupus erythematosus with renal involvement (G16a)	1-8	9-16	17+
• Focal segmental glomerulosclerosis (G16b)	1-5	6-10	11+
• Inpatient admissions and consultations (G18.1)	1-299	300+	NA
• New outpatient evaluations/consultations (G18.3)	1-499	500-999	1,000+
Neurology and Neurosurgery Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of patients undergoing epilepsy workups and treatments** (max points = 12)			
• Initial medical evaluations with patients newly diagnosed with epilepsy (H9a)	1-599	600-1,199	1,200+
• Standard EEG evaluations (H9b)	1-999	1,000-1,999	2,000+
• Long-term video EEG evaluations (H9c)	1-449	450-899	900+
• Number of first-time surgical procedures for epilepsy (H9d)	1-9	10-19	20+
Number of patients undergoing surgeries (max points = 42)			
• Surgical resection or laser ablation for epilepsy (H8)	1-7	8-15	16+
• Brain tumors (benign/malignant) (H16a)	1-24	25-49	50+
• Craniosynostosis (H16b)	1-19	20-39	40+
• Hydrocephalus shunt procedures (H16c)	1-49	50-99	100+
• Medically intractable epilepsy (H16d)	1-11	12-23	24+
• Spinal dysraphism (H16e)	1-19	20-39	40+
• Chiari I malformation (H16f)	1-14	15-29	30+
• Endoscopic treatment of hydrocephalus (H16g)	1-29	30-59	60+
• Brachial plexus exploration/reconstruction (H16h)	1-2	3-5	6+
• Spasticity (H16i)	1-11	12-23	24+

* Parenthetical references indicate related survey questions.

** Volume represents procedures, not patients.

(continued)

Table 7. Volume Measures by Specialty (continued)

Neurology and Neurosurgery, continued Volume Measures *	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of patients undergoing surgeries, continued			
• Vascular cases including endovascular procedures, excluding diagnostic angiograms conducted onsite at the children’s hospital or an adjoining facility with pediatric anesthesia assisting with the case (H16j)	1-9	10-19	20+
• Deep brain stimulation for primary dystonia (H16k)	1-4	5-9	10+
• Responsive neurostimulation (RNS) for medically intractable epilepsy (H16l)	1-2	3-5	6+
• Craniofacial procedures performed by pediatric neurosurgeons (H33)	1-29	30-59	60+
Orthopedics Volume Measures*	Low Volume (2 points)	Medium Volume (4 points)	High Volume (6 points)
Number of patients (max points = 24)			
• Patients transferred from another hospital for inpatient care (I14.1a)	1-149	150-299	300+
• Pediatric trauma patients who received pediatric orthopedic trauma surgery within 72 hours of admission (I14.1b)	1-299	300-599	600+
	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
• Pediatric trauma inpatients with fractures or musculoskeletal injuries (I14a)	1-999	1,000-1,999	2,000+
• Pediatric trauma outpatients with fractures or musculoskeletal injuries (I14b)	1-1,999	2,000-3,999	4,000+
• Scoliosis correction patients (I31a-b)	1-149	150-299	300+
• Single event multi-level surgery (I45)	1-19	20-39	40+

* Parenthetical references indicate related survey questions.

(continued)

** Volume represents procedures, not patients.

Table 7. Volume Measures by Specialty (continued)

Orthopedics Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of procedures and surgeries** (max points = 26)			
• Motion laboratory diagnostic clinical evaluations (I20a)	1-24	25-49	50+
• Motion laboratory diagnostic research evaluations (I20b)	1-24	25-49	50+
• Open reduction developmental dysplasia of the hip (I24a)	1-7	8-15	16+
• Ponsetti treatment for clubfoot in patients < 1 years old (I24b)	1-99	100-199	200+
• Bernese pelvic osteotomy in patients < 18 years old (I24c)	1-6	7-13	14+
• Cast treatment for infantile scoliosis < 5 years old (I24d)	1-7	8-15	16+
• ACL reconstruction (males < 14 years old or females < 12 years old) (I24e)	1-9	10-19	20+
• Femoral and tibial leg lengthening surgery (I24f)	1-3	4-7	8+
• Pollicization hand surgeries (I24g)	1	2+	NA
Pulmonology & Lung Surgery Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of tests and noninvasive procedures** (max points = 12)			
• 12- or 32- channel polysomnographic studies (J36)	1-699	700-1,399	1,400+
• Patients receiving home nocturnal PAP or bilevel therapy (J37)	1-89	90-179	180+
• Patients receiving non-invasive positive pressure ventilation support (J39)	1-34	35-69	70+
• Bronchoscopy and laryngoscopy (J49)	1-199	200-399	400+
Number of patients (max points = 17)			
• CF patients (J24a)	1-124	125-249	250+
• Neuromuscular weakness disorders (J29)	1-49	50-99	100+
• Ventilator dependent patients, 3 years (J40)	1-69	70-139	140+
• Rare lung disease (J26)	1-39	40-79	80+
• Lung disease of prematurity (J27)	1-59	60-119	120+
• Lung transplants, 3 years (J46.1)	1	2+	NA

* Parenthetical references indicate related survey questions.

(continued)

** Volume represents procedures, not patients.

Table 7. Volume Measures by Specialty (continued)

Urology Volume Measures*	Low Volume (1 point)	Medium Volume (2 points)	High Volume (3 points)
Number of minimally invasive procedures (max points = 9)			
• Stone treatment/shock wave lithotripsy (K11a)	1-14	15-29	30+
• Laparoscopic (or open surgery) orchiopexy/orchidectomy (K11b)	1-13	14-27	28+
• Laparoscopic cyst ablation, pyeloplasty, nephrectomy, partial nephrectomy, heminephrectomy, ureteral reimplantation, or ureteroureterostomy (K11c)	1-13	14-27	28+
Number of patients (max points = 27)			
• Pediatric urology outpatients (2 years), (K8b)	1-4,999	5,000-9,999	10,000+
• Pediatric urology surgical cases** (2 years) (K9)	1-999	1,000-1,999	2,000+
• Spina bifida program (K10a)	1-149	150-299	300+
• Comprehensive stone program (K10b)	1-74	75-149	150+
• Prenatal program (K10c)	1-49	50-99	100+
• Disorders of sexual differentiation program (K10d)	1-49	50-99	100+
• Exstrophy/cloaca/GU sinus program (K10e)	1-39	40-79	80+
• End stage renal disease program (K10f)	1-14	15-29	30+
• Oncofertility program (K10g)	6-9	10-19	20+
Number of surgeries (max points = 20)			
• Radical or partial nephrectomy for malignancies** (K12a)	1-2	3+	NA
• Open pyeloplasty, nephrectomy and partial nephrectomy, heminephrectomy, ureteral reimplantation, ureteroureterostomy for benign disease** (K12b)	1-19	20+	NA
• Radical prostatectomy for rhabdomyosarcoma, partial/total cystectomy or retroperitoneal lymph node dissection for cancer diagnoses	1-19	20+	NA
• Patients receiving exstrophy closures (K13a)	1-2	3+	NA
• Patients receiving reconstructive open procedures for incontinence or hostile bladder - open (K13b)	1-19	20+	NA
• Patients receiving posterior urethral valve ablation (K13c)	1-5	6+	NA
• Patients receiving proximal urethroplasty or complex revision for hypospadias (K13d)	1-44	45+	NA
• Distal hypospadias – primary repairs and not re-operative cases** (K14a)	1-59	60-119	120+
• Pyeloplasty** (K14b)	1-14	15-29	30+

n/a = not applicable.

* Parenthetical references indicate related survey questions.

** Volume represents procedures, not patients.

B. Normalization

As of the 2012-13 rankings, all structural measures have been normalized prior to weighting. Normalization transforms index values into a distribution between 0 and 1 based on the range of possible values for a given measure. The formula for normalization is provided in Equation (1):

$$\text{Equation (1)} \quad \text{Normalized Value} = X_i / (\text{Maximum}_i - \text{Minimum}_i)$$

where

X_i = the value for measure i and

Maximum_i = the highest *possible* value for measure i .

Minimum_i = the lowest *possible* value for measure i .

For example, the Urology patient volume measure is worth a maximum of 33 points. If a hospital received 18 out of 33 points, its normalized value for Urology patient volume would be $18/(33-0) = 0.55$. For nurse staffing, which does not have an absolute maximum, we cap the maximum value at 4.0 to reduce skewness in the data.

C. Weighting

For the 2012-13 rankings, we convened a special panel to provide feedback on the weighting of each measure within the three major rankings components. This evaluation was conducted both across specialties to build in a degree of consistency in weighting, and within specialties to identify keys to quality in a particular specialty. Overall, the weights were determined using input from the project team and working groups based on how important each measure was in defining the Donabedian components of quality of care within hospitals. The weights have been revised slightly for 2020-21, based on changes to the measures used in each specialty.

Table 8 shows the relative weight, by specialty, for each measure that makes up the structural component of the specialty rankings. The combined structural components comprise 33.3% of the overall score in each specialty. To determine the total structural points for a hospital, multiply the normalized value of each measure by the measure weight. In the example provided under normalization (Section IV.B), a hospital that received 18 out of 33 points for Urology patient volume would have a normalized score of 0.55. The relative weight for patient volume in Urology is 1. Therefore, the hospital would have a total of 0.55 for patient volume. Do this for all measures in a specialty, and then sum the values to determine the total points received. To determine the percent of the overall score for a given measure, divide the individual measure relative weight by the total weight for that specialty and multiply by 33.3 (since the combined structural components comprise 33.3% of the overall score in each specialty).

Table 8. Relative Weights of Individual Structural Measures by Specialty

Measure	Cancer	Cardiology & Heart Surgery	Diabetes & Endocrinology	Gastroenterology & GI Surgery	Neonatology	Nephrology	Neurology & Neurosurgery	Orthopedics	Pulmonology & Lung Surgery	Urology
Accredited by FACT	1.5									
Active fellowship program	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.5	1.25
Adoption of health information technology	1	1	1	1	1	1	1	1	1	1
Adult congenital heart program		1.25								
Advanced clinical services offered	1	1	1	1	1	1	1	1	1	1
Advanced technologies available	1	1	1	1	1	1	1	1	1	1
Bone marrow transplant services	1									
Clinical support services offered	1	1	1	1	1	1	1	1	1	1
Commitment to clinical research	1.5	1.25	1.25	1.25	1.25	1	1.25	1.25	1.5	1.5
Commitment to quality improvement	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Congenital heart program		1.25								
ECMO availability					1					
Enlists families in structuring care	1.5	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25	1.25
Has fulltime subspecialists available	1	1	1	1	1	1	1	1	1	1
Help for families	1	1	1	1	1	1	1	1	1	1
Neonatal Transport					1					
Nurse staffing	2	2	2	2	2	2	2	2	2	2
Percent of dialysis patients who had transplants						1.25				
Provides advanced palliative care	1.5									
Recognized as a Nurse Magnet hospital	2	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Specialized clinics and programs	1.25		1	1	1		1	1		1
Success in helping patients manage their asthma									1.25	

(continued)

Table 8. Relative Weights of Individual Structural Measures by Specialty, Continued

Measure	Cancer	Cardiology & Heart Surgery	Diabetes & Endocrinology	Gastroenterology & GI Surgery	Neonatology	Nephrology	Neurology & Neurosurgery	Orthopedics	Pulmonology & Lung Surgery	Urology
Success in managing neuro-muscular weakness disorder									1.25	
Tracking growth metrics for treated patients					1					
Transplant program (heart, liver, lung)		0.33		0.28					0.30	
Volume: Number of catheter procedures		1								
Volume: Number of dialysis patients						0.67				
Volume: Number of epilepsy workups and treatment							1			
Volume: Number of kidney biopsies						0.67				
Volume: Number of kidney transplants						0.27				
Volume: Number of minimally invasive procedures										1
Volume: Number of new patients	1									
Volume: Number of Norwood or hybrid surgeries		1								
Volume: Number of patients	1		1	1	1.5	0.67		1.5	1	1
Volume: Number of procedures			1							
Volume: Number of procedures and surgeries								1.5		
Volume: Number of surgeries	1	1.5		1			1			1
Volume: Number of tests and noninvasive procedures				1					1	
Total	24.00	21.08	17.75	19.03	20.25	18.03	17.75	18.75	20.05	19.00

V. Process

The process component in Best Children’s Hospitals is represented by three measures—commitment to best practices, ability to prevent infections, and expert opinion of pediatric specialists. The combined process measures are worth 33.3% of the overall score in all specialties except for pediatric cardiology. In pediatric cardiology, the process component is worth 28.3% of the total score.

A. Commitment to Best Practices

This measure evaluates hospitals’ commitment to following and implementing best practices. Best practices were identified for all specialties. *Table 9* identifies the best practices identified for each specialty and the number of points awarded.

Table 9. Commitment to Best Practices by Specialty

Cancer* (47 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
Offering the following to reduce radiation exposure to patients and staff (A10.1): <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with the American College of Radiology guidelines • Participation in the ACR CT dose index registry OR use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from American College of Radiology (ACR) • Accreditation in nuclear medicine from American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7
Having patients undergoing imaging studies (e.g., MRI, CT or voiding cystourethrogram scans) provided access to a certified child life specialist (A10.5)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty, continued

Cancer, continued* (47 points)	Points
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team training • Team training includes clear instructions and demonstration of roles and lines of communication • Team training is videotaped to allow review of performance and needs for improvement • Team training includes critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team training ends with development of an action plan to address problems identified during the training or simulation 	5
Having at least 0.5 FTE nurse practitioners, physician assistants or clinical nurses devoted to case management for each of the following patient populations (B4): <ul style="list-style-type: none"> • Hematologic malignancies • Solid tumors • Brain tumors • Stem cell transplants 	4
Offering an institutional code team to immediately address emergencies in outpatient cancer treatment clinics (B5)	1
Offering a parent advisory committee that meets at least twice a year (B11.3)	1
Participating in morbidity and mortality conferences at least quarterly (B12)	1
Having multidisciplinary tumor boards that meet at least monthly to discuss the following patient populations in active treatment (B13): <ul style="list-style-type: none"> • Hematologic malignancy • Solid tumor • Brain tumor • Stem cell transplant 	4
Promoting ease of access through the following mechanisms (B14): <ul style="list-style-type: none"> • Offering onsite direct oncology-specific patient care from hematology/oncology providers during evenings and weekends • A coordinated outreach program that provides community-based follow-up care • Multidisciplinary clinics allowing patients to see multiple care providers in a single visit 	3
Submitting data to the Center for International Blood & Marrow Transplant Research (CIBMTR) or the Stem Cell Therapeutic Outcome Database (SCTOD) (B20)	1
Patients have thyroidectomies performed by a high-volume thyroid surgeon (>25 thyroid resections per year) (B27.1)	1
Percentage of patients receiving radical nephrectomy for Wilms tumor underwent lymph node sampling during the procedure (B27.2): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 80\%$ • 2 points for $\geq 80\%$ 	2

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty, continued

Cancer, continued* (47 points)	Points
Percentage of patients 3-5 years post-completion of therapy seen in a formally structured late effects or off-therapy clinic from over the past 3 years (B28): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ 	2
Percentage of patients with certain cancer diagnoses, 1-3 years post-treatment, who had documentation of a formal neuropsychological evaluation conducted since the completion of therapy (B28.1): <ul style="list-style-type: none"> • 1 point for $\geq 25\%$ & $< 75\%$ • 2 points for $\geq 75\%$ 	2
Percentage of school-age patients with certain cancer diagnoses were formally assessed for school intervention services since diagnosis and before the end of the last calendar year (B28.2): <ul style="list-style-type: none"> • 1 point for $\geq 25\%$ & $< 75\%$ • 2 points for $\geq 75\%$ 	2
$\geq 75\%$ of pediatric brain tumor patients (from B27b) were enrolled in a formal, comprehensive neuro-oncology clinic for their care coordination (B28.3)	1
Percentage of patients presenting with febrile neutropenia who receive intravenous antibiotics within one hour of initial triage (B31.1): <ul style="list-style-type: none"> • 1 point for $\geq 75\%$ & $< 85\%$ • 2 points for $\geq 85\%$ & $< 95\%$ • 3 points for $\geq 95\%$ 	3
Cardiology & Heart Surgery* (57 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
Offering the following to reduce radiation exposure to patients and staff (A10.1): <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with the American College of Radiology guidelines • Participation in the ACR CT dose index registry OR use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty, continued

Cardiology & Heart Surgery, continued* (57 points)	Points
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from American College of Radiology (ACR) • Accreditation in nuclear medicine from American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team training • Team training includes clear instructions and demonstration of roles and lines of communication • Team training is videotaped to allow review of performance and needs for improvement • Team training includes critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team training ends with development of an action plan to address problems identified during the training or simulation 	5
Number of pediatric cardiothoracic surgeons with subspecialty certification in congenital heart surgery from the American Board of Thoracic Surgery (E2.1) or certification from foreign organizations (e.g., the Fellowship of the Royal Colleges of Surgeons (FRCS)), practice exclusively or primarily (greater than 90%) in congenital heart surgery and are not eligible for Subspecialty-Certification in Congenital Heart Surgery from the ABTS (E2.2): <ul style="list-style-type: none"> • 1 point for 1 surgeon • 2 points for 2+ surgeons 	2
Having IMPACT Registry data available in the NCDR database for each of the following quarters (E9): <ul style="list-style-type: none"> • 2018 Quarter 3 • 2018 Quarter 4 • 2019 Quarter 1 • 2019 Quarter 2 	4
Having at least one of the following types of data submitted to the NCDR database in 2019 (E10): <ul style="list-style-type: none"> • Diagnostic catheterizations • Interventional catheterizations • Myocardial biopsies • Catheter ablations 	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty, continued

Cardiology & Heart Surgery, continued* (57 points)	Points
Offering lead extraction for pacemaker and automatic implantable cardioverter defibrillator (ICD/AICD) leads onsite (E15.1) <ul style="list-style-type: none"> • 1 point for offsite • 2 points for onsite 	2
Participating in the STS Public Reporting On-Line Program (E18.3)	2
Offering the following conferences/programs (E27): <ul style="list-style-type: none"> • Multidisciplinary morbidity and mortality conferences • Multidisciplinary maternal/fetal medicine conferences • Active home surveillance program for infants after Stage 1 palliation for hypoplastic left heart syndrome • A neurodevelopmental follow-up program for children with complex congenital heart disease or at risk for adverse neurodevelopmental outcomes • Patient planning conference • Support groups for patients and families with congenital heart conditions • Multidisciplinary management program for complex congenital heart disease patients who experience long term change to physical or cognitive functioning 	7
Engaging in the following surgical safety procedures for cardiac surgical procedures (E35): <ul style="list-style-type: none"> • Conventional pre-procedural "time-out" • Pre-procedural briefings • Post-procedural debriefings • Implementation of a hand-off protocol or briefing 	4
Engaging in the following surgical safety procedures for cardiac catheterization procedures (E35.1): <ul style="list-style-type: none"> • Conventional pre-procedural "time-out" • Pre-procedural briefings • Implementation of a hand-off protocol or briefing 	3
Using clinical practice guidelines to manage perioperative and postoperative care for the following patient populations (E36): <ul style="list-style-type: none"> • Single ventricle/shunt management • Two-ventricle repairs • Infant feeding • Anticoagulation with Coumadin • Sedation and pain management 	5
Routinely tracking and reporting every occurrence of the following surgical admission outcomes parameters to the STS database (E37): <ul style="list-style-type: none"> • Unplanned reoperation during the same hospital admission • Re-exploration for bleeding • Deep sternal wound infection/mediastinitis requiring debridement • Arrhythmia necessitating pacemaker or permanent pacemaker 	4
Routinely tracking and submitting to IMPACT every unplanned cardiothoracic procedure, vascular surgical procedure, other surgical procedure or cardiac catheterization due to a catheterization complication (E37.1)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty, continued

Cardiology & Heart Surgery, continued* (57 points)	Points
Percent of hybrid and Norwood Stage 1 surgery patients alive 1 year after surgery who had a neurodevelopment evaluation prior to 24 months of age (E41): <ul style="list-style-type: none"> • At least 75% of patients in evaluation (Year 1) • At least 75% of patients in evaluation (Year 2) • At least 75% of patients in evaluation (Year 3) • At least 75% of patients in evaluation (Year 4) 	4
Diabetes & Endocrinology* (119 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
Offering the following to reduce radiation exposure to patients and staff (A10.1): <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with the American College of Radiology guidelines • Participation in the ACR CT dose index registry OR use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from the American College of Radiology (ACR) • Accreditation in nuclear medicine from the American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from the ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7
Having patients undergoing imaging studies (e.g., MRI, CT or voiding cystourethrogram scans) provided access to a certified child life specialist (A10.5)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty, continued

Diabetes & Endocrinology, continued* (119 points)	Points
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team trainings • Team trainings include clear instructions and demonstration of roles and lines of communication • Team trainings are videotaped to allow review of performance and needs for improvement • Team trainings include critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team trainings end with the development of an action plan to address problems identified during the training or simulation 	5
Diabetes staff taking a leadership role in organizing and running a diabetes camp (C10)	1
Having pediatric diabetes staff take a leadership role in a formal advocacy effort supporting the rights of patients (C11)	1
Hosting or actively involved in organizing a diabetes-specific technology education program (C13)	1
Administering a formal, written assessment of diabetes management knowledge after initial education and yearly thereafter (C15)	1
Diabetes education program recognized by American Diabetes Association or American Association of Diabetes Educators (C14)	1
Percent of diabetes patients on insulin therapy admitted as inpatients to other services, were seen by providers in the pediatric diabetes program (C16 and C16.1): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ & $< 90\%$ • 3 points for $\geq 90\%$ 	3
Having a formal written transition program to prepare pediatric patients for the transition to an adult diabetes program (C17)	1
Having $\geq 90\%$ of diabetes outpatients receive a written (or electronic) report of their diagnosis/findings and a treatment plan at the conclusion of their most recent visit (C18a)	1
Having $\geq 90\%$ of diabetes inpatients receiving a written (or electronic) report of their diagnosis/findings and a treatment plan at the conclusion of their most recent visit (C18b)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty, continued

Diabetes & Endocrinology, continued* (119 points)	Points
Always including the following elements in summaries given to patients in outpatient clinic visits (C19): <ul style="list-style-type: none"> • Complete insulin dosages • Blood glucose testing and record-keeping recommendations • A1c values from today and/or percent of time "in range" (70 – 180 mg/dl) from at least 1 weeks of CGM data • Next visit date and time • Information on when and how to contact the Diabetes Center • Referrals made for laboratory, ophthalmological, dental and mental health before next visit • Behavioral goals 	7
Having a clinical database of attributes of current, active diabetes patients that is used for quality assessment and improvement (C20)	1
Having a written plan to review inpatient incidents of insulin-related medication errors and adverse drug events requiring IV glucose treatment (C21)	1
Having written consensus protocols for management of the following patient populations (C22): <ul style="list-style-type: none"> • Glucagon mini-dose for families • Periodic screening for complications of diabetes in the outpatient clinic • Evaluation of hyperglycemia in critically ill inpatients • Outpatient management of Type 2 diabetes patients • Outpatient management of pre-diabetes patients who typically have obesity and insulin resistance 	5
Performing care review for all patients admitted with a primary diagnosis of diabetes at an interdisciplinary team prior to discharge (C23)	1
Having regularly scheduled interdisciplinary care conferences to discuss diabetes patients with poor control (C25 and C26): <ul style="list-style-type: none"> • 1 point for 1-11 times/year • 2 points for 12+ times/year 	2
Having written protocols for identifying "high risk" patients and enrolling them in special pathways (C27)	1
Interacting with clinical laboratory or pathology service to review lab findings, problems and updates (C28)	1
Asking about the number of hospital admissions, emergency visits or urgent care visits since the last diabetes outpatient visit (C29)	1
Tracking the number of hospital admissions, emergency visits or urgent care visits since the last diabetes outpatient visit in EMR (C29.1)	1
Percentage of primary diabetes care patients with face-to-face visit with nutritionist or certified diabetes educator for medical nutrition therapy (C30a): <ul style="list-style-type: none"> • 1 point for ≥ 50% & < 75% • 2 points for ≥ 75% 	2

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty, continued

Diabetes & Endocrinology, continued* (119 points)	Points
Percentage of primary diabetes care patients with face-to-face visit with CDE or equivalent for diabetes education (C30b): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ 	2
Percentage of primary diabetes care patients with face-to-face visit with a social worker or psychologist for an assessment (C30c): <ul style="list-style-type: none"> • 1 point for $\geq 25\%$ & $< 50\%$ • 2 points for $\geq 50\%$ 	2
Percentage of Type 1 primary care diabetes patients with a TSH documented in their medical chart in past 2 years (C31a): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ & $< 90\%$ • 3 points for $\geq 90\%$ 	3
Percentage of Type 1 primary care diabetes patients over 10 years of age who had a lipid profile within the past 5 years (C31b): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ & $< 90\%$ • 3 points for $\geq 90\%$ 	3
Percentage of Type 1 primary care diabetes patients over 10 years of age (with diabetes for at least 5 years) who received a microalbuminuria screening in the past year (C31c): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ & $< 90\%$ • 3 points for $\geq 90\%$ 	3
Percentage of Type 1 primary care diabetes patients over 10 years of age (with diabetes for at least 5 years) who received a dilated retinal or non-mydratiac camera examination in the past year (C31d): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ & $< 90\%$ • 3 points for $\geq 90\%$ 	3
Percentage of Type 2 primary care diabetes patients who had a lipid profile performed in the past year (C31e): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ & $< 90\%$ • 3 points for $\geq 90\%$ 	3
Percentage of Type 2 primary care diabetes patients who received a microalbuminuria screening in the past year (C31f): <ul style="list-style-type: none"> • 1 point for $\geq 40\%$ & $< 60\%$ • 2 points for $\geq 60\%$ & $< 80\%$ • 3 points for $\geq 80\%$ 	3

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty, continued

Diabetes & Endocrinology, continued* (119 points)	Points
<p>Percentage of Type 2 primary care diabetes patients who received a dilated retinal or non-mydriatic camera examination in the past 2 calendar years (C31g):</p> <ul style="list-style-type: none"> • 1 point for $\geq 40\%$ & $< 60\%$ • 2 points for $\geq 60\%$ & $< 80\%$ • 3 points for $\geq 80\%$ 	3
<p>Percentage of Type 1 primary care diabetes patients that are < 18 years of age who were treated in the past 12 months or longer who scheduled for 4 or more outpatient clinic visits in past 12 months (C32a):</p> <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ & $< 90\%$ • 3 points for $\geq 90\%$ 	3
<p>Percentage of Type 1 primary care diabetes patients treated in the past 12 months or longer who attended 4 or more outpatient clinic visits (C32b):</p> <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ & $< 90\%$ • 3 points for $\geq 90\%$ 	3
<p>Average number of documented interpretations of continuous glucose monitoring (CGM) readings associated with ambulatory visits for primary Type 1 diabetes care pediatric patients that are < 18 years of age for private and Medicaid insurance in the past year (C32.1, C32.2):</p> <p>Private insurance</p> <ul style="list-style-type: none"> • 1 point for $>.5$ and <1 CGM per patient • 2 points for ≥ 1 CGM per patient <p>Medicaid insurance</p> <ul style="list-style-type: none"> • 1 point for $>.25$ and $<.75$ CGM per patient • 2 points for $\geq .75$ CGM per patient 	4
<p>Percentage of Type 1 primary care diabetes patients that are < 18 years of age on an insulin pump in the past calendar year for private and Medicaid insurance (C33):</p> <p>Private insurance</p> <ul style="list-style-type: none"> • 1 point $\geq 20\%$ & $< 60\%$ • 2 points $\geq 60\%$ <p>Medicaid insurance</p> <ul style="list-style-type: none"> • 1 point $\geq 20\%$ & $< 40\%$ • 2 points $\geq 40\%$ 	4
<p>Percentage of Type 1 and Type 2 primary diabetes care patients aged 13 to < 18 screened for depression in the past calendar year (C34):</p> <ul style="list-style-type: none"> • 1 point for $\geq 25\%$ & $< 50\%$ • 2 points for $\geq 50\%$ 	2

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Diabetes & Endocrinology, continued* (119 points)	Points
Percentage of Type 1 and Type 2 primary diabetes care patients who had an abnormal score on their depression screen (see C34) were either referred for assessment by a mental health professional (social worker, licensed counselor, psychologist, or psychiatrist) or are already under the care of a mental health professional (C34.1): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ 	2
Percentage of Type 1 diabetes outpatients with daily glucose blood glucose measurements available for review for the past 2 weeks (C36): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ & $< 90\%$ • 3 points for $\geq 90\%$ 	3
Having a written curriculum for diabetes self-management education that addresses self-care behaviors (C37)	1
Tracking the number of school days missed for diabetes-related reasons (C38)	1
Providing a dedicated team of Type 2 diabetes providers (C40)	1
Distributing patient education materials that address the details of their conditions to patients with the following conditions (C44): <ul style="list-style-type: none"> • Adrenal insufficiency • Congenital hypothyroidism • Diabetes insipidus 	3
Distributing patient education materials to patients that address the potential side effects of taking the following medications (C45): <ul style="list-style-type: none"> • Anti-thyroid medication • Growth hormone • Cortisol • Oral contraceptive pills 	4
Using a clinical database used by the program to evaluate performance (C54 and C54.1)	1
Discussing thyroid cancer patient cases in active treatment at a tumor board at least once a quarter (C56)	1
Percentage of patients admitted to the hospital in the past year with an endocrine disorder that have an admission or consultation note written by a physician in the pediatric endocrinology program (C58): <ul style="list-style-type: none"> • 1 point for $< 50\%$ • 2 points for $\geq 50\%$ 	2

* Parenthetical references indicate related survey questions.

(continued)

Diabetes & Endocrinology, continued* (119 points)	Points
<p>Having a system in place to alert providers that the following types of patients have not returned for care (C63):</p> <ul style="list-style-type: none"> • Type 1 diabetes • Congenital hypothyroidism • Congenital adrenal hyperplasia • Growth hormone therapy • Precocious puberty on therapy • Hyperthyroidism on anti-thyroid medication <p>Hospitals received 1 point for 1-3 types and 2 points for 4-6 types.</p>	2
<p>Participating in multidisciplinary evaluation and management of the following types of patients (C64):</p> <ul style="list-style-type: none"> • Endocrine complications in hematology/oncology patients • Endocrine complications in post-transplant patients • Metabolic bone disease and osteogenesis imperfecta • Inborn errors of metabolism or evaluation of hypoglycemia <p>Hospitals received 1 point for 1-2 types and 2 points for 3-4 types.</p>	2
<p>Hosting or conducting the following conferences or educational programs in the last year (C65):</p> <ul style="list-style-type: none"> • Joint case conferences with Internal Medicine • Joint case conferences with genetics program • Pediatric endocrinology case conference • Pediatric endocrinology journal club • CME-granting education activity conferences <p>Hospitals received 1 point for 1-34 conferences and 2 points for 35 or more conferences.</p>	2
Gastroenterology & GI Surgery* (35 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
<p>Offering the following to reduce radiation exposure to patients and staff (A10.1):</p> <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with American College of Radiology guidelines • Participation in the ACR CT dose index registry or use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Gastroenterology & GI Surgery, continued* (35 points)	Points
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from American College of Radiology (ACR) • Accreditation in nuclear medicine from American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7
Having patients undergoing imaging studies (e.g., MRI, CT or voiding cystourethrogram scans) provided access to a certified child life specialist (A10.5)	1
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team training • Team training includes clear instructions and demonstration of roles and lines of communication • Team training is videotaped to allow review of performance and needs for improvement • Team training includes critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team training ends with development of an action plan to address problems identified during the training or simulation 	5
Average “third next available” appointment time for new patients for an outpatient office visit (D6): <ul style="list-style-type: none"> • 1 point for >7 & ≤ 30 days • 2 points for ≤ 7 days 	2
Providing educational programs for the following disease-specific GI conditions (D9): <ul style="list-style-type: none"> • Inflammatory bowel disease, Crohn’s disease or colitis • Celiac disease • Liver disease • Eosinophilic esophagitis • Chronic intestinal failure 	5

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Gastroenterology & GI Surgery, continued* (35 points)	Points
Providing the following diagnostic and therapeutic procedures (D11.1) <ul style="list-style-type: none"> • Interventional radiology embolization for gastrointestinal bleeding • Interventional radiology for image guided liver biopsies • Interventional radiology performance of transjugular intrahepatic portosystemic shunt (TIPS) • Interventional radiology performance of transjugular (TJ) liver biopsies • Interventional radiology performance of hepatic vein wedge pressure measurement 	5
Having regular, multidisciplinary morbidity and mortality conferences for pediatric GI patients (D26)	1
Having a standard mechanism to determine if complications have occurred in patients who underwent outpatient GI procedures (D27)	1
Having 1 or more IRB-approved protocols that provide GI patients access to drugs, biologics or devices through compassionate use (D28)	1
Having the Pediatric GI program serve as a referral center for inflammatory bowel disease (IBD) patients for second or subsequent evaluations (D33.1)	1
Neonatology* (99 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
Offering the following to reduce radiation exposure to patients and staff (A10.1): <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with American College of Radiology guidelines • Participation in the ACR CT dose index registry OR use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from American College of Radiology (ACR) • Accreditation in nuclear medicine from American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Neonatology, continued* (99 points)	Points
Having patients undergoing imaging studies (e.g., MRI, CT or voiding cystourethrogram scans) provided access to a certified child life specialist (A10.5)	1
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team training • Team training includes clear instructions and demonstration of roles and lines of communication • Team training is videotaped to allow review of performance and needs for improvement • Team training includes critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team training ends with development of an action plan to address problems identified during the training or simulation 	5
Having in-house 24x7 coverage provided by board-certified or board-eligible neonatologists (F2.1)	1
Patient load per nurse practitioner or physician assistant (F3): <ul style="list-style-type: none"> • 1 point for ≥ 9 patients • 2 points for < 9 patients 	2
Having at least 50% of NICU staff who support the lactation program FTEs with IBCLC and/or CBC certification (F4c/F4b)	1
Percentage of eligible direct clinical care RNs who are nationally certified in neonatal intensive care (F4.1): <ul style="list-style-type: none"> • 1 point for $\geq 25\%$ & $< 50\%$ • 2 points for $\geq 50\%$ 	2
Patient load per neonatologist (F5): <ul style="list-style-type: none"> • 1 point for ≥ 18 • 2 points for < 18 	2
Patient load per Licensed independent contractor (attending, fellow, resident or advanced practitioner) on the night shift (F5.1): <ul style="list-style-type: none"> • 1 point for ≥ 15 • 2 points for < 15 	2
Patient load per nutritionist (F7.1): <ul style="list-style-type: none"> • 1 point for ≥ 20 • 2 points for < 20 	2

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Neonatology, continued* (99 points)	Points
Providing the following elements of a "Safe Sleep" program (F8.1): <ul style="list-style-type: none"> • Mandatory Safe Sleep Education for NICU Staff • Required documentation that Safe Sleep has been discussed with family prior to discharge • Policy in place for use of devices (swings, infant seats, etc.) • NICU-based written criteria for Safe Sleep within NICS • Safe Sleep policy includes a minimum length of time prior to discharge to implement safe sleep practices if no contraindications exist • Safe Sleep policy indicates that infants greater than 32 weeks post-conceptual gestational age should be placed in supine position if no contraindications exist 	6
Engaging in Safe Sleep auditing to ensure that patient sleep environment and position is appropriate for safe sleep practice (F8.2)	1
Monitoring compliance with NICU-based Safe Sleep policy at least weekly or at least bi-weekly (F8.3)	1
Patient load per staff person social workers (F11.1): <ul style="list-style-type: none"> • 1 point for ≥ 15 patients • 2 points for < 15 patients 	2
Tracking the proportion of infants discharged from NICU on partial or full mother's own milk (F10)	1
Offering a dedicated area within the facility for milk and formula preparation (F10.2)	1
Offering the following for nutrition and breastfeeding (F10.3): <ul style="list-style-type: none"> • Cohort of NICU RNs specially trained in lactation counseling • NICU-specific breast milk committee that meets at least monthly • Process to facilitate obtaining a breast pump (within 48 hours of identified need) for home use • NICU specific risk reduction program that includes process designed to reduce breast milk errors • Donor breast milk program with written institution-specific criteria for the initiation and discontinuation of donor breast milk 	5
Employing the following risk-reduction practices (F10.4): <ul style="list-style-type: none"> • Bar coding system for correct breast milk identification • Dedicated breast milk technician who prepares milk for proper identification and distribution 	2
Tracking breast milk administration error rate (F10.5)	1
Having a mandatory neonatal consult or a neonatologist co-managing care for surgical patients in the NICU (F16.1)	1
Having at least 75% of anesthesiologists with board-certification or are board-eligible in pediatric anesthesia (F16.2)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Neonatology, continued* (99 points)	Points
Engaging in the following interaction with hospital's CICU (F18): <ul style="list-style-type: none"> All preterm cardiac patients <28 days of age receive a neonatology consult All newborn cardiac patients <28 days of age (preterm and full term) receiving a neonatology consult 	2
Providing a percutaneous intravenous central catheter (PICC) team with specialized training to place and maintain PICC lines in NICU patients (F20)	1
Availability of PICC line placement services (F20.1): <ul style="list-style-type: none"> 1 point for day shift 2 points for 24/7 coverage 	2
Mandating that core NICU staff participate in the following training protocols at least once every 2 years (F22): <ul style="list-style-type: none"> Neonatal unplanned code response Arrhythmia treatment including use of defibrillator Simulation of emergency evacuation of the NICU ECMO emergency simulation training Exchange transfusion simulation or just in time training Other training 	6
Having at least 75% of neonatal fellows complete training in the following procedure protocols (F23.1): <ul style="list-style-type: none"> Chest tube placement Intubation Neonatal resuscitation program 	3
Having at least 75% of current attending physicians in the Level IV NICU who have completed simulation or other training to refresh their skills with each of the following procedures (F23.2): <ul style="list-style-type: none"> chest tube placement pericardiocentesis abdominal paracentesis double volume exchange transfusion cardioversion 	5
Number of standardized hand-off tools used by physicians and advanced practitioners to inform clinical staff during shift transitions (F29.1): <ul style="list-style-type: none"> 1 point for 1-2 tools 2 points for 3-4 tools 	2
Number of standardized hand-off tools used by nurses to inform clinical staff during shift transitions (F30.1): <ul style="list-style-type: none"> 1 point for 1 tool 2 points for 2-3 tools 	2
Tracking patients' first postoperative temperatures and using it as a quality metric (F31)	1
Tracking unintended extubation of NICU patients (F32)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Neonatology, continued* (99 points)	Points
Frequency of quality review process (F32.2): <ul style="list-style-type: none"> • 1 point for a retrospective multidisciplinary review weekly or monthly • 1 point for a prospective mini-root cause analysis review within 12 hours 	2
Conducting multidisciplinary review of all unplanned readmissions to determine if preventable (F33)	1
Having a formal program for reviewing neonatal transfer cases received from other hospitals that includes a formal feedback mechanism to the referring facility (F33.1)	1
Providing the following for very-low-birth-weight and low gestational age infants (F34): <ul style="list-style-type: none"> • Starter protein solution available on day of admission • Very low birth weight feeding protocol • "Kangaroo care" routinely provided for infants receiving mechanical ventilation • Dedicated team of bedside RNs with additional training in the care of preterm infants that care for VLBW infants in your NICU • Respiratory support weaning protocol 	5
Having or being associated with a fetal diagnosis and counselling program either onsite or at another facility (F34.1)	1
Providing mandatory prenatal consultation when the postnatal patient management plan requires care in the Level IV NICU (F34.2)	1
Holding multidisciplinary patient management conferences to discuss plans for the delivery and early NICU management of fetuses with congenital abnormalities (F34.3): <ul style="list-style-type: none"> • 2 points for meeting at least weekly or at least monthly • 1 point for meeting less frequently than monthly 	2
Offering family meetings or counseling that includes neonatologists, genetic counselors and relevant specialists when fetuses are expected to require care in the Level IV NICU (F34.4)	1
Offering a fetal MRI program for assessment of fetal neurologic, thoracic, and abdominal anomalies (F34.5)	1
Having a medication error reporting system/database (F35)	1
Offering the following to evaluate and reduce medication errors (F35.1): <ul style="list-style-type: none"> • NICU-specific multidisciplinary committee, including a residency trained NICU clinical pharmacist • Residency trained NICU clinical pharmacist who participates in multidisciplinary rounds • Access to an up-to-date electronic version of a neonatal-specific drug information reference which includes the ability to check IV compatibility • Utilize medication administration technology including barcode administration and smart pumps with a dedicated neonatal drug library • Formalized process for discharge medication counseling 	5

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Nephrology* (59 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
Offering the following to reduce radiation exposure to patients and staff (A10.1): <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with American College of Radiology guidelines • Participation in the ACR CT dose index registry OR use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from American College of Radiology (ACR) • Accreditation in nuclear medicine from American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7
Having patients undergoing imaging studies (e.g., MRI, CT or voiding cystourethrogram scans) provided access to a certified child life specialist (A10.5)	1
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team training • Team training includes clear instructions and demonstration of roles and lines of communication • Team training is videotaped to allow review of performance and needs for improvement • Team training includes critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team training ends with development of an action plan to address problems identified during the training or simulation 	5
Percentage of school-age pediatric dialysis patients enrolled in a school or vocational rehabilitation program (G10): <ul style="list-style-type: none"> • 1 point for <50% • 2 points for ≥ 50% 	2

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Nephrology, continued* (59 points)	Points
At least 50% of native kidney biopsies were performed by a pediatric nephrologist or pediatric nephrology fellow using real-time ultrasound or ultrasound guidance (G14.1)	1
Have access to an interventional radiologist to perform image guided renal biopsies, nephrostomies, and AV fistula/graft management (G16.1)	1
Participating in regular interdisciplinary clinical conferences to review and coordinate the care of patients in the following specialties (G17): <ul style="list-style-type: none"> • Urology/uroradiology • Renal pathology • Rheumatology • Fetal health 	4
Providing the following services in support of the pediatric dialysis unit (G19): <ul style="list-style-type: none"> • Designated medical director board-certified in pediatric nephrology with 0.25 or more FTE support for this position • Quality Assurance Performance Improvement activities reviewed independently from the adult dialysis service • Pediatric maintenance dialysis patients receive treatment in a unit independent from adult patients • Dedicated nursing staff with formal training in pediatric dialysis • At-home maintenance hemodialysis program for adolescents (maintained more than 3 months consecutively at home) that is either standalone or conducted in conjunction with an adult program • At-home maintenance peritoneal dialysis program 	6
Availability and prescription of therapeutic plasma exchange to patients (G19.1): <ul style="list-style-type: none"> • Available and prescribed by Pediatric Nephrology (2 points) • Available but NOT prescribed by Pediatric Nephrology (1 point) 	2
Availability of ABPM to patients and report generated (G19.2): <ul style="list-style-type: none"> • Available and report generated by Pediatric Nephrology (2 points) • Available but report NOT generated by Pediatric Nephrology (1 point) 	2
Ratio of accesses received in the last 2 years per patient for permanent hemodialysis vascular central venous catheters placed in children < 5 years of age (G22a): <ul style="list-style-type: none"> • 1 point for > 1.5 & ≤ 4 accesses per patient • 2 points ≥ 1 & ≤ 1.5 accesses per patient 	2
Ratio of accesses received in the last 2 years per patient for permanent hemodialysis vascular central venous catheters placed in children, 5-19 years of age (G22b): <ul style="list-style-type: none"> • 1 point for > 1.5 & ≤ 4 accesses per patient • 2 points ≥ 1 & ≤ 1.5 accesses per patient 	2

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Nephrology, continued* (59 points)	Points
Ratio of accesses received in the last 2 years per patient for hemodialysis AV fistula/graft access placements in children, 10-19 years of age on maintenance dialysis (G22c): <ul style="list-style-type: none"> • 1 point for > 1.25 & ≤ 4 accesses per patient • 2 points for ≥ 1 & ≤ 1.25 accesses per patient 	2
Ratio of accesses received in the last 2 years per patient for peritoneal dialysis catheters placed in children < 5 (G22d): <ul style="list-style-type: none"> • 1 point for > 1.25 & ≤ 4 accesses per patient • 2 points for ≥ 1 & ≤ 1.25 accesses per patient 	2
Ratio of catheters placed per patient for peritoneal dialysis catheters placed in children and adolescents, 5-19 (G22e): <ul style="list-style-type: none"> • 1 point for > 1.25 & ≤ 4 catheters per patient • 2 points for ≥ 1 & ≤ 1.25 catheters per patient 	2
Offering a formal transition program for kidney transplant patients from pediatric to adult care when needed (G25)	1
Offering a formal transition program for dialysis patients into adult care when needed (G26)	1
At least 50% of percutaneous kidney biopsies were performed by a pediatric nephrologist or pediatric nephrology fellow using real-time ultrasound or ultrasound guidance (G27.1)	1
Percentage of living donor nephrectomies conducted via laparoscopic procedure (G29): <ul style="list-style-type: none"> • 1 point for 70-79% • 2 points for 80-89% • 3 points for ≥ 90% 	3
Percentage of kidney transplant patients <18 years of age that were preemptive (G31): <ul style="list-style-type: none"> • 1 point for 10-20% • 2 points for >20% 	2
Offering the following programs to support pediatric patients undergoing kidney transplant (G33): <ul style="list-style-type: none"> • Quality of life assessment • Child life program for kidney transplant patients • Transplant pharmacist 	3
Maintaining a database of current kidney transplant patients with clinical data to allow for quality assessment and improvement of care (G38)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Neurology & Neurosurgery* (37 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
Offering the following to reduce radiation exposure to patients and staff (A10.1): <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with American College of Radiology guidelines • Participation in the ACR CT dose index registry OR use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from American College of Radiology (ACR) • Accreditation in nuclear medicine from American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7
Having patients undergoing imaging studies (e.g., MRI, CT or voiding cystourethrogram scans) provided access to a certified child life specialist (A10.5)	1
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team training • Team training includes clear instructions and demonstration of roles and lines of communication • Team training is videotaped to allow review of performance and needs for improvement • Team training includes critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team training ends with development of an action plan to address problems identified during the training or simulation 	5
Having at least 50% of patients receiving a surgical resection or laser ablation for epilepsy have intraoperative electrocorticography and/or extraoperative monitoring (H8 and H8.1)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Neurology & Neurosurgery, continued* (37 points)	Points
Having \geq 75% of EEG tests incorporated into the patients' medical chart within designated timeframes (H10): <ul style="list-style-type: none"> • Standard EEG medical evaluations interpreted and recorded within 36 hours of being conducted • Long-term vEEG evaluations interpreted and recorded within 5 days from completion of the study 	2
Offering neuropsychological evaluations for surgical patients with the following diagnoses (H15): <ul style="list-style-type: none"> • Benign or malignant brain tumors postoperative • Traumatic brain injury/concussion postoperative • Medically intractable epilepsy postoperative • Craniofacial disorders postoperative • Congenital heart disease postoperative • Stroke • Demyelinating diseases 	7
Participating in the following nationally audited research programs that focus on outcome measures specific to neurology and neurosurgery (H19): <ul style="list-style-type: none"> • Neurocritical Care Research Group • International Pediatric Stroke Study • 	2
Participating in community outreach programs to improve health in the community (H20.1)	1
Providing a transition of care program that helps patients prepare for and then move from pediatric to adult care providers (H20.2)	1
Engaging in the following activities (H22): <ul style="list-style-type: none"> • Maintaining a surgical mortality database • Holding regular mortality and morbidity conferences • Interdisciplinary care conferences held monthly or more often 	3
Having an epilepsy program designated Level IV by National Association of Epilepsy Centers (H32)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Orthopedics* (82 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
Offering in the following to reduce radiation exposure to patients and staff (A10.1): <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with American College of Radiology guidelines • Participation in the ACR CT dose index registry OR use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from American College of Radiology (ACR) • Accreditation in nuclear medicine from American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7
Having patients undergoing imaging studies (e.g., MRI, CT or voiding cystourethrogram scans) provided access to a certified child life specialist (A10.5)	1
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team trainings • Team training includes clear instructions and demonstration of roles and lines of communication • Team training is videotaped to allow review of performance and needs for improvement • Team training includes critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team training ends with development of an action plan to address problems identified during the training or simulation 	5
Percentage of nurse practitioners and physician assistants receiving pediatric orthopedic surgery-related continuing education credit or continuing medical credit (I3.1): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ 	2

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Orthopedics, continued* (82 points)	Points
Percentage of RNs receiving pediatric orthopedic surgery-related continuing education credit or continuing medical credit (I4.1a): <ul style="list-style-type: none"> • 1 point for ≥50% & <75% • 2 points for ≥75% 	2
Percentage of medical assistants receiving pediatric orthopedic surgery-related continuing education credit or continuing medical credit (I4.1b): <ul style="list-style-type: none"> • 1 point for ≥50% & <75% • 2 points for ≥75% 	2
Number of pediatric orthopedic surgeons who are active or candidate members of the Pediatric Orthopaedic Society of North America (I5): <ul style="list-style-type: none"> • 1 point for 1-2 • 2 points for 3+ 	2
Having at least 75% of patient MRI and CT examinations read by pediatric radiologists with musculoskeletal imaging expertise (I9.1)	1
Providing pediatric imaging center with the following services (I10): <ul style="list-style-type: none"> • Pediatric protocols to reduce radiation exposure • Digitally stored test results, images, and medical records accessible from locations offsite or away from the hospital • Intraoperative navigation system • Low-dose, three-dimensional upright body imaging for evaluating idiopathic scoliosis • Pediatric anesthesia services to support sedation and general anesthesia for imaging in very young children • MR arthrography • Image guided thermal ablation of bone tumors • Interventional radiology image guided bone biopsies • Medical director for musculoskeletal imaging 	9
All musculoskeletal cancer patients discussed at a tumor board at least once a quarter (I17)	1
More than 75% of tumor boards attended by a pediatric (medical) oncologist (I17.1a)	1
More than 75% of tumor boards attended by an orthopaedic (surgical) oncologist (I17.1b)	1
Participating in regular multidisciplinary morbidity and mortality conferences (I18)	1
Having a designated inpatient trauma operating room that 100% guarantees orthopedics a “first case of the day start” (I28)	1
Having a policy in place that provides even greater operating room access based on periodic demand (I29)	1
Having a preoperative coordinated care review process led by a nursing coordinator that meets at least monthly to evaluate high-risk patients and prepare them for surgery and hospitalization (I30)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Orthopedics, continued* (82 points)	Points
Having surgeons who perform surgical correct for scoliosis participate in the "Surgeon Performance Program" provided by the Setting Scoliosis Straight Foundation (SSSF) (I32.1)	1
Having access to at least 1 of the following types of anesthesiologists (I34): <ul style="list-style-type: none"> • Pediatric anesthesiologist • Pediatric spine anesthesiologist 	1
Having at least 1 surgical correction for scoliosis case that was staffed by either a pediatric anesthesiologist or a pediatric spine anesthesiologist (I35)	1
Having at least 50% of pediatric spine anesthesiologists who were the anesthesiologist of record for 20 or more surgical corrections for scoliosis cases in the past 2 years (I35.1)	1
Percentage of surgical spine patients 8 or older completing SRS-22 or SRS-30 (I36a): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ 	2
Percentage of surgical spine patients 8 or older completing CPCHILD (I36b): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ 	2
Percentage of surgical spine patients 8 or older completing another orthopedic patient reported outcomes questionnaire (I36c): <ul style="list-style-type: none"> • 1 point for $\geq 50\%$ & $< 75\%$ • 2 points for $\geq 75\%$ 	2
Percentage compliance with written checklists/guidelines for patients with neurological injury associated with surgery for idiopathic scoliosis (I37a): <ul style="list-style-type: none"> • 1: $\geq 70\%$ & $< 85\%$ • 2: $\geq 85\%$ 	2
Percentage compliance with written checklists/guidelines for patients with neurovascular injuries associated with supracondylar fractures or dislocation of the knee (I37b): <ul style="list-style-type: none"> • 1: $\geq 70\%$ & $< 85\%$ • 2: $\geq 85\%$ 	2
Percentage compliance with written checklists/guidelines for patients with spinal trauma resulting in acute spinal cord injury (I37c): <ul style="list-style-type: none"> • 1: $\geq 70\%$ & $< 85\%$ • 2: $\geq 85\%$ 	2
Having at least 1 in-service presentation or formal lecture to an RN audience (I39)	1
Establishing a professional relationship with one or more prosthetic/orthotics providers such that they attend clinic on a regular basis (I40)	1
Having a fixed surgery support team that are dedicated to working with pediatric orthopedic surgeons (I41)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Orthopedics, continued* (82 points)	Points
Having exactly the same fixed surgery support team working together during normal working hours (I41.1): <ul style="list-style-type: none"> • 4 points for > 75% of the time • 1 point for 50-75% of the time 	4
Providing afterhours or weekend "on call" service for a fixed surgery support team (I42)	1
Having exactly the same fixed surgery support team working together after hours or on weekends (I42.1): <ul style="list-style-type: none"> • 4 points for > 75% of the time • 1 point for 50-75% of the time 	4
Rate of single event multi-level surgery patients who received a Multimodal pain management (I46a/I45): <ul style="list-style-type: none"> • 1 point for $\geq 75\%$ & $< 90\%$ • 2 points for $\geq 90\%$ 	2
Rate of single event multi-level surgery patients who received a postoperative assessment by anesthetic/pain team (I46b/I45): <ul style="list-style-type: none"> • 1 point for $\geq 75\%$ & $< 90\%$ • 2 points for $\geq 90\%$ 	2
Hosting or being actively involved in organizing a cerebral palsy support group (I47)	1
Having the following elements of a Narcotic Stewardship program (I48): <ul style="list-style-type: none"> • A non-narcotic pathway in place for patients being admitted for orthopedic surgery • 'Right Size' opioid prescribing recommendations based on patient age and procedure for orthopedic surgical patients • Narcotic safety education provided to families of orthopedic surgical patients with instructions on how to safely dispose of unused narcotics • Plan to ensure tracking of potential pain medication seeking or opioid addition in orthopedic surgical patients • A system to automatically limit the number of narcotic tablets prescribed to orthopedic surgical patients following treatment for supracondylar fracture of the humerus or isolated femoral shaft fractures 	5

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Pulmonology & Lung Surgery* (48 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
Offering the following to reduce radiation exposure to patients and staff (A10.1): <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with American College of Radiology guidelines • Participation in the ACR CT dose index registry OR use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from American College of Radiology (ACR) • Accreditation in nuclear medicine from American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7
Having patients undergoing imaging studies (e.g., MRI, CT or voiding cystourethrogram scans) provided access to a certified child life specialist (A10.5)	1
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team training • Team training includes clear instructions and demonstration of roles and lines of communication • Team training is videotaped to allow review of performance and needs for improvement • Team training includes critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team training end with development of an action plan to address problems identified during the training or simulation 	5
Screening all pulmonology patients for tobacco smoke exposure and actively counseling family members who smoke (J6)	1
Having access to a thorough onsite assessment of patients' home environment and offer guidance for reducing exposures that contribute to asthma (J9)	1

* Parenthetical references indicate related survey questions.

(continued)

Table 9. Commitment to Best Practices by Specialty (continued)

Pulmonology & Lung Surgery, continued* (48 points)	Points
Having written consensus protocols for the following conditions (J6.1): <ul style="list-style-type: none"> • Asthma exacerbations • Bronchiolitis • Croup • Cystic fibrosis • Uncomplicated pneumonia • Complicated pneumonia • Initiation of tracheostomy of home ventilator support • Tracheostomy or ventilator-dependent patients • Pneumothorax care pathway • Acute chest syndrome • Other care pathways including airway emergencies such as foreign body, epiglottitis/tracheitis, or inhalation injury 	11
Having a formal plan to actively transition CF patients from pediatric care to adult care (J25)	1
Provide bronchial artery embolization for CF patients (J25.1)	1
Having a pediatric sleep disorders clinic that addresses the needs of patients with ventilation or other sleep disorders and manages the patient's positive airway pressure (J38)	1
Having multidisciplinary care team to coordinate the care of long-term ventilator-dependent patients with the following members (J42): <ul style="list-style-type: none"> • ENT or critical care physician • Pulmonologist • Physiatrist • Respiratory therapist • Social worker • Dietician 	6
Participating in formal programs for the outpatient management of pediatric patients with the following conditions (J50): <ul style="list-style-type: none"> • Sickle cell anemia • Aerodigestive disorders • Craniofacial disorders • Pulmonary hypertension 	4
Having a protocol for preparing and assisting in the transition of patients from pediatric to adult pulmonology (J53)	1
Providing financial support for staff to attend extramural continuing education (J54)	1
Interpreting exercise challenge and bronchoprovocation testing (J56)	1

* Parenthetical references indicate related survey questions.

Table 9. Commitment to Best Practices by Specialty (continued)

Urology* (42 points)	Points
Having at least 2.0 FTE pediatric interventional radiologists (A4.1)	1
Offering the following to reduce radiation exposure to patients and staff (A10.1): <ul style="list-style-type: none"> • Provide a designated medical director of radiology to oversee quality and safe practices • Iterative reconstruction software on all computed tomography (CT) scanners • MRI safety program compliant with American College of Radiology • Participation in the ACR CT dose index registry OR use of dose monitoring software for tracking pediatric patients undergoing CT scans 	4
Using computerized tomography (CT) protocols that adjust milliampere-seconds (mAs) and peak kilovolts (kVp) (A10.2)	1
Maintaining the following certifications (A10.3): <ul style="list-style-type: none"> • Accreditation in computerized tomography (CT) imaging from American College of Radiology (ACR) • Accreditation in nuclear medicine from American College of Radiology (ACR) or the Intersocietal Accreditation Commission (IAC) • All sonographers have an ultrasound certificate from the American Registry of Diagnostic Medical Sonographers (ARDMS) the American Registry of Radiologic Technologists (ARRT) • One or more pediatric sonographers have a pediatric ultrasound certificate from the American Registry for Diagnostic Medical Sonographers (ARDMS) • Program accreditation in ultrasound from ACR or AIUM • Accreditation in MRI from ACR • At least one full-time Medical Physicist on-site to monitor and optimize pediatric imaging 	7
Having patients undergoing imaging studies (e.g., MRI, CT or voiding cystourethrogram scans) provided access to a certified child life specialist (A10.5)	1
Engaging in activities designed to ensure high reliability (A39): <ul style="list-style-type: none"> • All clinical staff are trained in code response using simulations or other team training • Team training includes clear instructions and demonstration of roles and lines of communication • Team training is videotaped to allow review of performance and needs for improvement • Team training includes critical event debriefing or team discussions that focus on identifying what worked well and where improvement is needed • All team training ends with development of action plan to address problems identified during the training or simulation 	5
Having regular morbidity and mortality conferences to discuss pediatric urology patients (K6a)	1
Having regular case conferences to discuss surgical management of complex cases (K6b)	1
Having regular multi-specialty case review and/or process review conferences for all surgical services (K6c)	1

* Parenthetical references indicate related survey questions.

Table 9. Commitment to Best Practices by Specialty (continued)

Urology, continued* (42 points)	Points
Having the following elements of a Narcotic Stewardship program (K10.2): <ul style="list-style-type: none"> • A non-narcotic pathway in place for patients being admitted for genitourinary (GU) surgery • 'Right Size' opioid prescribing recommendations based on patient age and procedure • Narcotic safety education provided to families with instructions on how to safely dispose of unused narcotics • Plan to ensure tracking of potential pain medication seeking or opioid addition patients 	4
Monitoring reconstructive procedure for incontinence or hostile bladder for the following operative complications (K13.1): <ul style="list-style-type: none"> • Monitor the renal function of patients with reconstructed GU tracts with eGFR at least once during the 12-month reporting period • Hydronephrosis based on ultrasound or nuclear medicine testing • Continence • Stomal complications 	4
Offering the following weekend, afterhours, and satellite outpatient clinics for elective care (K21): <ul style="list-style-type: none"> • Scheduled (non-emergency) weekday evening clinic appointments (after 5pm) • Scheduled (non-emergency) weekend surgical appointments • Scheduled (non-emergency) weekend clinic appointments • Satellite clinics in the community 	4
Having the following protocols in place (K22): <ul style="list-style-type: none"> • Protocol for teaching home intermittent catheterization • Radiation reduction/safety protocols for urology patients (e.g., retrograde pyelogram protocol, ureteroscopy protocol) • Standardized clinical pathway for children presenting with acute stone pain to the Emergency Department • Standardized clinical pathway for postoperative patients • Educational materials for patients and families on urological conditions that are updated on a regular basis • Child life specialists for perioperative care in the operating room and for office procedures • Sedation services for VCUG/noxious procedures 	7
Making use of a patient portal to enable families to access electronic medical records and communicate with their physicians and medical staff (K23)	1

* Parenthetical references indicate related survey questions.

B. Ability to Prevent Infections

Incorporating infection-preventing measures captures the commitment of a hospital to identifying and implementing proven means of reducing the risk of various infections.

All-Specialty Infection-Preventing Measures

A core set of submeasures for all specialties was worth up to 31 points, as shown in *Table 10*. Specialty-specific measures in all specialties allowed an additional 1-29 points, depending on the specialty.

Table 10. Core Infection-Preventing Measures, All Specialties (31 points)

All Specialties* (31 points)	Points
Percentage of compliant hand hygiene observations for inpatient care in the past 12 months (F37.1 for Neonatology, A25 for all other specialties): <ul style="list-style-type: none"> 1 point for $\geq 80\%$ & $< 90\%$ 2 points for $\geq 90\%$ 	2
Providing at least 0.2 FTE financial support per 100 beds for a pediatric infectious disease specialist to serve as the medical director of the infection prevention program (A26.1, A1.2)	1
Providing at least 1.0 FTE infection preventionists per 100 beds (A27, A1.2): <ul style="list-style-type: none"> 2 points for ≥ 1.45 FTE per 100 beds 1 point for ≥ 1.0 FTE per 100 beds and < 1.45 FTE per 100 beds 	2
Having at least 1 infection preventionist certified by the Certification Board in Infection Control (A27.1)	1
Ensuring that at least 90% of the following staff received influenza vaccination ^{††††} (A28): <ul style="list-style-type: none"> Physicians (including attending physicians, fellows, residents) Nursing staff and mid-level providers 	2
Ensuring that at least 95% of the following staff received Tdap vaccination (A29): <ul style="list-style-type: none"> Physicians (including attending, fellows, residents) Nursing staff and mid-level providers 	2
Requiring all volunteers to receive or provide documentation of: <ul style="list-style-type: none"> Influenza vaccination (A29.1) Tdap vaccination (A29.2) 	2
Offering an influenza vaccination program for families and primary caregivers (A29.3)	1
Offering an adult Tdap booster program for families and caregivers (A29.4)	1

* Parenthetical references indicate related survey questions.

(continued)

^{††††} “Eligible” healthcare providers, as defined by NHSN, includes all providers who have worked at the facility for at least 1 working day between October 1 and December 31 during the reporting period, regardless of clinical responsibility or patient contact. This includes healthcare providers who joined after October 1 or left before December 31, or who were on extended leave during part of the reporting period. Working for any number of hours a day counts as one working day. Providers should be counted as individuals rather than full-time equivalents. Licensed practitioners who receive a direct paycheck from the reporting facility, or who are owners of the reporting facility, should be counted as employees. Due to the timing of the Hospital Survey, please submit data for October 1 through December 31 only. Include staff who refuse immunizations for personal reasons as eligible healthcare providers. For more information see NHSN guidelines: <https://www.cdc.gov/nhsn/pdfs/hps-manual/vaccination/hps-flu-vaccine-protocol.pdf>

Table 10. Core Infection-Preventing Measures, All Specialties, continued (31 points)

All Specialties*, continued (31 points)	Points
Having the following elements of antimicrobial stewardship program: <ul style="list-style-type: none"> • Actively monitoring internal days of therapy (DOT) of antibiotic use per 1,000 patients (A31a) • Restriction or pre-authorization of selected antimicrobial agents to prevent potential resistance from overuse (A31b) • Implementing prospective review and real time intervention regarding antimicrobial use or “handshake stewardship” (A31c) • Use of clinical guidelines in prescribing antimicrobials (A31d) • IV to PO conversion program available to ensure correct dosage (A31e) • Dedicated pharmacist to antimicrobial stewardship program (ASP) (A31.1a): <ul style="list-style-type: none"> ○ At least 1.0 FTE support for hospitals with at least 250 beds ○ At least 0.5 FTE support for hospitals with less than 250 beds • At least 0.3 FTE support for the role of medical director of the pediatric ASP program (A31.1b) • At least 0.2 FTE support for a dedicated analyst to support ASP program (A31.1c) • Formal policy on the use of antimicrobials (A32a) • Letter of support for the ASP by hospital administration (A32b) • An ASP committee that meets at least quarterly (A32c) • Regular tracking and reporting of ASP data to hospital clinicians (A32d) • Annual and ongoing education to hospital staff regarding ASP (A32e) 	13
Using the following interventions to reduce indwelling urinary catheter utilization in ICU settings (A34.1) <ul style="list-style-type: none"> • Written indications for insertion and/or removal of indwelling urinary catheters • Routine removal of urinary catheters following surgery • Bladder scanning • Non-indwelling catheter (e.g., in and out or straight catheter) for urinary retention 	4

* Parenthetical references indicate related survey questions.

Specialty-Specific Infection-Preventing Measures

Cancer (6 additional points). Hospitals receive 1 point for auditing hand hygiene compliance rates (A24) via direct observation, a hybrid of direct observation and electronic monitoring, or via electronic monitoring. Hospitals received 1 point for having a formal program to prevent hospital-acquired pressure injuries (A36). Hospitals received 1 point for actively tracking seasonal influenza vaccinations in cancer patients on active chemotherapy (B32). Up to 3 additional points were awarded according to the percentage vaccinated (B33): 1 point for $\geq 50\%$ and $< 75\%$, 2 points for $\geq 75\%$ or $< 90\%$, and 3 points for $\geq 90\%$.

Cardiology & Heart Surgery (10 additional points). Hospitals receive 1 point for auditing hand hygiene compliance rates (A24) via direct observation, a hybrid of direct observation and electronic monitoring, or via electronic monitoring. Hospitals received 1 point for having a formal program to prevent hospital-acquired pressure injuries (A36). Hospitals received 1 point for

monitoring compliance with preoperative antibiotic prophylaxis for a sample of cases or 2 points for monitoring compliance for all cardiothoracic surgeries (E32). Up to 2 additional points were awarded according to the percentage of compliance (E33): 1 point if $\geq 75\%$ and $< 90\%$, or 2 points if $\geq 90\%$.

Hospitals received up to 4 additional points for engaging in the following surgical site infection prevention procedures: pre-operative bath (E31a), no use of razor for hair removal (E31b), preparation of skin at surgical site with alcohol containing agent (E31c), and screening for and appropriately decolonizing Staph Aureus utilizing a nasal antiseptic OR perform universal decolonization on all patient undergoing a cardiac surgical procedure (E31d).

Diabetes & Endocrinology (6 additional points). Hospitals receive 1 point for auditing hand hygiene compliance rates (A24) via direct observation, a hybrid of direct observation and electronic monitoring, or via electronic monitoring. Hospitals received 1 point for having a formal program to prevent hospital-acquired pressure injuries (A36). Hospitals received 1 point for actively tracking seasonal influenza vaccinations in diabetes outpatients (C42). Up to 3 additional points were awarded according to the percentage vaccinated (C43): 1 point for $\geq 50\%$ and $< 75\%$, 2 points for $\geq 75\%$ and $< 90\%$, or 3 points for $\geq 90\%$.

Gastroenterology & GI Surgery (12 additional points). Hospitals receive 1 point for auditing hand hygiene compliance rates (A24) via direct observation, a hybrid of direct observation and electronic monitoring, or via electronic monitoring. Hospitals received 1 point for having a formal program to prevent hospital-acquired pressure injuries (A36). Hospitals received up to 2 points for actively tracking seasonal influenza vaccinations for chronic intestinal failure patients (D18) and post-liver transplant patients (D23). Up to 3 points each were awarded based on the percentage of both chronic intestinal failure patients (D19) and liver-transplant patients (D24) vaccinated (6 points total): 1 point for $\geq 50\%$ and $< 75\%$, 2 points for $\geq 75\%$ and $< 90\%$, or 3 points for $\geq 90\%$. Hospitals received up to 2 additional points for implementing strategies for preventing central-line associated bloodstream infections for total parenteral nutrition patients (D37): 1 point for implementing one or two strategies, or 2 points for implementing 3 or more strategies.

Neonatology (12 additional points). Hospitals received 1 point for having a formal program to prevent hospital-acquired pressure injuries (A36). Hospitals received 1 point for auditing hand hygiene compliance rates via electronic monitoring and/or direct observation (F37). Hospitals received 1 point for having written standardized guidelines and 1 point for monitoring compliance with the guidelines for antibiotic use in each the following situations (up to 10 additional points): surgical NEC repair or drain placement (F38a), small bowel atresia repair (F38b), gastroschisis abdominal closure (F38c), medical necrotizing enterocolitis (F38d), culture negative sepsis (F38e).

Nephrology (29 additional points). Hospitals receive 1 point for auditing hand hygiene compliance rates (A24) via direct observation, a hybrid of direct observation and electronic monitoring, or via electronic monitoring. Hospitals received 1 point for having a formal program to prevent hospital-acquired pressure injuries (A36). Hospitals received 1 point each (up to 6 points) for actively tracking seasonal influenza and pneumococcal vaccinations for hemodialysis patients (G12a, G13a), peritoneal dialysis patients (G12b, G13b) and kidney transplant patients (G34, G35). Up to 3 additional points were awarded for each of the 6 groups (up to 18 points) according to the percentage up to date on their vaccinations (G12a, G12b, G13a, G13b, G34.1, G35.1): 1 point for $\geq 50\%$ and $< 75\%$, 2 points for $\geq 75\%$ and $< 90\%$, or 3 points for $\geq 90\%$.

Hospitals received up to 3 points for employing the following strategies to prevent pediatric outpatient hemodialysis catheter associated BSI events: focused patient and parent education about infection prevention (G36a), formal collaboration between infection prevention and dialysis providers (G36b), and formal improvement project focused on reducing CLABSIs in these patients (G36c).

Neurology & Neurosurgery (10 additional points). Hospitals receive 1 point for auditing hand hygiene compliance rates (A24) via direct observation, a hybrid of direct observation and electronic monitoring, or via electronic monitoring. Hospitals received 1 point for having a formal program to prevent hospital-acquired pressure injuries (A36). Hospitals received 1 point for monitoring compliance with preoperative antibiotic prophylaxis for a sample of cases or 2 points for monitoring compliance for all ventricular surgeries (H25). Up to 2 additional points were awarded based on the percentage of compliance (H26): 1 point if $\geq 75\%$ and $< 90\%$, or 2 points if $\geq 90\%$. Hospitals received 1 point for actively tracking SSIs for ventricular shunt surgeries (H27). Up to 3 points were awarded for the rate of surgical site infections per 100 ventricular shunt surgeries performed in the prior year (H28). Points were awarded as follows: 1 point if $> 6\%$ and $\leq 10\%$, 2 points if $> 3\%$ and $\leq 6\%$, or 3 points if $\leq 3\%$.

Orthopedics (7 additional points). Hospitals receive 1 point for auditing hand hygiene compliance rates (A24) via direct observation, a hybrid of direct observation and electronic monitoring, or via electronic monitoring. Hospitals received 1 point for having a formal program to prevent hospital-acquired pressure injuries (A36). Hospitals received 1 point for monitoring compliance with preoperative antibiotic prophylaxis for a sample of cases or 2 points for monitoring compliance for all spinal fusion surgeries (I21). Up to 2 additional points were awarded according to the percentage of compliance (I22): 1 point if $\geq 75\%$ and $< 90\%$, or 2 points if $\geq 90\%$. Hospitals received 1 point for monitoring SSIs for spinal fusion surgeries using an established standard program (I23 and I23.1).

Pulmonology & Lung Surgery (19 additional points). Hospitals receive 1 point for auditing hand hygiene compliance rates (A24) via direct observation, a hybrid of direct observation and electronic monitoring, or via electronic monitoring. Hospitals received 1 point for having a formal program to prevent hospital-acquired pressure injuries (A36). Hospitals received 1 point each (up to 4 points) for actively tracking seasonal influenza vaccinations for asthma patients (J14), cystic fibrosis patients (J18), neuromuscular weakness disorder patients (J33) or ventilator-dependent patients (J43). Up to 3 additional points were awarded for each of the 4 groups (up to 12 points) according to the percentage vaccinated (J15, J19, J34, J44): 1 point for $\geq 50\%$ and $< 75\%$, 2 points for $\geq 75\%$ and $< 90\%$, or 3 points for $\geq 90\%$. Hospitals received 1 additional point for having at least 75% of vaccine-eligible patients treated between October 1 and December 31 receive a seasonal influenza vaccine (J27.1).

Urology (1 additional point). Hospitals receive 1 point for auditing hand hygiene compliance rates (A24) via direct observation, a hybrid of direct observation and electronic monitoring, or via electronic monitoring.

C. Expert Opinion with Pediatric Specialists

Expert opinion can be viewed as a form of peer review of the hospital’s capabilities across a wide variety of processes related to quality of care. For all specialties, expert opinion scores were based on responses to the physician surveys conducted in 2018, 2019 and 2020. Scores were calculated separately in each year, and averaged such that each year’s scores are given equal weighting in the final expert opinion score, as shown in *Table 11*.

Table 11. Expert Opinion Weight by Survey Year

Sample Source	Expert Opinion Weight	Overall Weight
2020 Physician Survey	33.3%	5.0%
2019 Physician Survey	33.3%	5.0%
2018 Physician Survey	33.3%	5.0%
Total	100.0%	15.0% ^{§§§§}

^{§§§§} In Cardiology & Heart Surgery, the overall weight for reputation was reduced to 8.5% starting in the 2017-18 rankings.

The sections below describe the approach used for the 2020 survey, which was similar to the 2018 and 2019 surveys. The approaches used for the 2018 and 2019 surveys are provided in the corresponding methodology reports for those years, available at www.rti.org/besthospitals.

2020 Survey Approach

Sample Selection

Starting with the 2015-16 rankings, the sample has been drawn from a database of U.S. physicians compiled by Doximity. Similar to the AMA Physician Masterfile, which was used as the sampling frame in previous years, Doximity's comprehensive Physician Database includes every practicing U.S. physician, identified by National Provider Identifier (NPI) number. Its sources include the U.S. Department of Health and Human Services NPI Registry, specialty boards (e.g., the American Board of Medical Specialties and the American Board of Surgery) and state medical boards. Doximity's proprietary database is augmented by more than 750,000 registered and verified physician members who review and update their profiles to provide another set of primary data.

Table 12 provides the population counts of pediatric specialists in the database by Doximity members and nonmembers.

Table 12. Population Counts by Best Hospitals Specialty, Doximity Members and Nonmembers

Best Children’s Hospitals Specialty	Subspecialties	Doximity Members		Doximity Nonmembers	
		Count	%	Count	%
Cancer	Pediatric Hematology-Oncology	2214	81.5%	501	18.5%
Cardiology & Heart Surgery****	Pediatric Cardiology, Pediatric Cardiac Surgery, and Pediatric Thoracic Surgery	2587	85.5%	440	14.5%
Diabetes & Endocrinology	Pediatric Endocrinology	1187	79.3%	309	20.7%
Gastroenterology & GI Surgery	Pediatric Gastroenterology and Pediatric Transplant Hepatology	1318	84.8%	237	15.2%
Neonatology	Neonatal-Perinatal Medicine	4166	76.2%	1304	23.8%
Nephrology††††	Pediatric Nephrology	619	80.5%	150	19.5%
Neurology & Neurosurgery****	Child Neurology and Pediatric Neurological Surgery*	2177	80.8%	516	19.2%
Orthopedics§§§§	Pediatric Orthopedic Surgery	909	85.8%	150	14.2%
Pulmonology & Lung Surgery	Pediatric Pulmonary	969	86.4%	153	13.6%
Urology*****	Pediatric Urology	219	59.3%	150	40.7%

* These specialists were drawn from the American Board of Pediatric Neurological Surgery (www.abpns.org).

**** To ensure both cardiologists and heart surgeons are represented in the nonmember sample, we sample 110 cardiologists and 40 heart surgeons. However, the actual total number of Doximity members in Pediatric Cardiac Surgery and Pediatric Thoracic Surgery was 219 and the total number of nonmembers was 14. To meet the subsample of 40 heart surgeons, 26 Doximity members with specialties in Pediatric Cardiac Surgery and Pediatric Thoracic Surgery were selected to receive the nonmember survey. These physicians’ scores are included in the nonmember reputation values.

†††† The actual total number of Doximity members in Nephrology was 623 and the total number of nonmembers was 144. However, 6 member nephrologists were randomly selected to be included in the mail survey so that at least 150 physicians received the mail survey. These physicians’ scores are included in the nonmember reputation values.

**** To ensure both neurologists and neurosurgeons are represented in the nonmember sample, we sample 110 neurologists and 40 neurosurgeons. However, the actual total number of Doximity members in Pediatric Neurosurgery was 276 and the total number of nonmembers was 31. To meet the subsample of 40 neurosurgeons, 9 Doximity members with specialties in Pediatric Neurosurgery were selected to receive the nonmember survey. These physicians’ scores are included in the nonmember reputation values.

§§§§ The actual total number of Doximity members in Orthopedics was 926 and the total number of nonmembers was 138. However, 12 member nephrologists were randomly selected to be included in the mail survey so that at least 150 physicians received the mail survey. These physicians’ scores are included in the nonmember reputation values.

***** The actual total number of Doximity members in Urology was 330 and the total number of nonmembers was 39. However, 111 member urologists were randomly selected to be included in the mail survey so that at least 150 physicians received the mail survey. These physicians’ scores are included in the nonmember reputation values.

Data Collection Procedures

Doximity members and nonmembers were surveyed separately, as described below.

Member survey. The Doximity member survey identified a total of 16,365 physicians eligible in one of the 10 pediatric specialties as of November 1, 2019. In February, physicians received an initial email invitation with a link to the survey. The survey asked for names of up to 10 hospitals in the physician's specialty that provide the best care to patients with serious conditions, without considering location or expense. Nonresponding physicians received up to two follow-up email reminders with a link to the survey. In addition, survey-eligible Doximity members – i.e., those who were board certified in a relevant specialty – received alerts upon login to Doximity.com or use of the Doximity app inviting them to participate.

Nonmember survey. The nonmember survey was conducted by randomly sampling 1,500 Doximity nonmembers – 150 specialists in each of the 10 specialty areas^{†††††}. Stratifying by census region (https://www2.census.gov/geo/pdfs/maps-data/maps/reference/us_regdiv.pdf), we selected physicians in each region proportional to the size of the population. For example, if 40% of all Doximity nonmembers in a specialty were from the South, then 40% of our sample would have included physicians in that region. Sampling physicians proportional to population size allowed us to minimize the weights needed to produce expert opinion scores that are representative of the nation.

Sampled physicians were asked to complete a brief survey containing a single nomination element. The survey of nonmembers was identical to the survey of Doximity members but was conducted via mail instead of the web. It asked physicians to supply the names of up to 10 hospitals in their specialty that provide the best care to patients with serious conditions, without considering location or expense. A copy of the mailed survey is available in *Appendix C*.

Up to four mailings were sent to sampled Doximity nonmembers. Each mailing included a cover letter, questionnaire, and business reply envelope. The first survey mailing also included a small token incentive. The survey was conducted from January 8 through March 20, 2020.

^{†††††} A few specialties had nonmember populations smaller than the required sample size for the Doximity nonmember survey. To ensure equivalent sample sizes across all specialties, 26 Doximity members in Pediatric Thoracic Surgery, 6 Doximity members in Pediatric Nephrology, 9 Doximity members in Pediatric Neurosurgery, 12 Doximity members in Pediatric Orthopedics, and 111 Doximity members in Pediatric Urology were included in the nonmember sample. These physicians were treated as nonmembers for the purposes of weighting and scoring.

Response Rates

The overall response rate for the 2018, 2019 and 2020 surveys was 30.0% using the American Association of Public Opinion Research (AAPOR) standard response rate 6^{####}, which treats undeliverables as ineligible. The 2020 combined response rate for the Doximity member and nonmember surveys was 28.7% using AAPOR standard response rate 6. Below we provide more detail on the response rates to the 2020 Doximity member and nonmember surveys.

Member survey. Of 16,365 Doximity members, 4,737 completed the web survey by March 20, 2020. The final response rate was 28.9% using AAPOR standard response rate 6. **Table 13** shows response rates by region and specialty.

Table 13. Member Survey Response Rates (%) by Region and Specialty, 2020

Specialty	Midwest	Northeast	South	West	Total
Cancer	37.9	28.3	34.1	25.6	31.8
Cardiology & Heart Surgery	37.9	35.5	38.2	29.1	35.7
Diabetes & Endocrinology	35.4	29.5	33.9	27.3	31.8
Gastroenterology & GI Surgery	35.9	34.9	28.5	28.2	31.9
Neonatology	24.9	22.7	17.1	15.1	19.5
Nephrology	43.9	30.5	33.2	23.7	33.0
Neurology & Neurosurgery	35.8	26.4	25.7	25.5	28.0
Orthopedics	39.8	38.9	31.8	25.2	33.3
Pulmonology & Lung Surgery	35.3	26.3	28.1	28.9	29.6
Urology	53.8	45.7	35.7	37.5	43.4
Total	34.5	29.1	28.3	23.9	28.9

Nonmember survey. Of 1,500 nonmember physicians sampled for this year's report, 321 were deemed ineligible after determining that they were no longer actively practicing or because we were unable to verify their eligibility. Of the remaining 1,179 physicians, 290 returned the completed questionnaire. The final response rate was 24.6% using the AAPOR standard response rate 6. **Table 14** shows response rates by region and specialty.

Definitions available at https://www.aapor.org/AAPOR_Main/media/publications/Standard-Definitions20169theditionfinal.pdf

Table 14. Nonmember Survey Response Rates (%) by Region and Specialty, 2020

Specialty	Midwest	Northeast	South	West	Total
Cancer	30.0	11.1	25.7	20.7	21.6
Cardiology & Heart Surgery	11.8	25.0	23.8	25.9	22.9
Diabetes & Endocrinology	25.0	13.3	27.0	12.0	19.8
Gastroenterology & GI Surgery	23.5	17.2	32.5	26.7	25.9
Neonatology	8.3	19.0	7.3	12.5	10.9
Nephrology	38.2	16.0	11.5	23.8	23.6
Neurology & Neurosurgery	17.2	7.4	15.8	25.9	16.5
Orthopedics	38.9	33.3	31.8	17.1	29.1
Pulmonology & Lung Surgery	23.1	21.2	18.6	25.0	21.4
Urology	62.5	56.5	47.4	47.1	50.8
Total	27.1	21.7	24.5	25.6	24.6

Survey Response Weighting

For the Doximity member survey, we used post-stratification weights for age by gender (55+ male, <55 male, and female^{§§§§§§}) as well as census region. Weights were constructed and applied to each physician’s survey response to make nominations representative of Doximity members at the national level. Since all Doximity members were surveyed, weights were used only to adjust for differences in nonresponse by region and demographics. In each specialty, the sample for the Nonmember physician survey was stratified only by census region (Midwest, Northeast, South and West). The sample size in each specialty was too small to stratify by the demographic characteristics used in the Doximity sample. Weights were constructed and applied to each physician’s survey response to make nominations representative of Doximity nonmembers at the national level. Weights were based on probability of selection within each unique specialty-region combination, adjusting to account for nonresponders.

Expert opinion scores were tabulated separately for Doximity members and nonmembers, and then combined to create the 2020 expert opinion score. The weight is based on the proportion of Doximity members and nonmembers in the population so the expert opinion score is representative of all physicians in the nation (see Table 12). Expert opinion scores for each of the past three years were then averaged to create the final, weighted expert opinion values that appear in the methodology report.


§§§§§§ Age categories were collapsed for females because there were too few female physicians over 55 in the sample.

Log Transformation

Weighted three-year expert opinion values are displayed in the ranking tables. Before the expert opinion data were combined into the Index of Hospital Quality (IHQ), log transformation was implemented to adjust for the skewed distribution of values. By its nature, a survey that solicits recommendations for “best hospitals” will result in data that do not follow a normal distribution. Relatively few hospitals will receive even one recommendation, and of the hospitals recommended, even fewer will receive a substantial number of nominations. Since other ranking components such as structural measures and mortality are not similarly skewed, expert opinion would have a greater impact on the final rankings than is warranted if left unadjusted.

Log transformation reshapes the distribution to more closely match expert opinion data to those of other components. The transformation is applied to the weighted expert opinion data. The transformed data are then normalized and multiplied by 100 to provide scores ranging from 0 to 100. *Figure 1* demonstrates the effect of the transformation.

Figure 1. Impact of Log Transformation on Expert Opinion



The transformed expert opinion scores are mostly higher than the untransformed scores, but the relative increases are larger for low scores than for high ones. For example, an untransformed expert opinion value of 1% has a transformed score of 4 (4 times greater), an untransformed value of 10% has a transformed score of 29 (2.9 times greater), and an untransformed value of 60% has a transformed score of 81 (1.4 times greater). Skewness is thus reduced, and the impact of expert opinion on final standing in the rankings is slightly diminished.

D. Normalization and Weighting

The process component, which consists of commitment to best practices, infection-prevention program, and expert opinion, is worth one-third (33.3%) of the overall score in each specialty except for pediatric cardiology & heart surgery. The overall measure weight and the process component weight for all other specialties is provided in *Table 15*.

Table 15. Weight of Individual Process Measures (All Specialties Except Pediatric Cardiology & Heart Surgery)

Process Measure	Overall Weight	Process Component Weight
Commitment to Best Practices	9.2%	27.5%
Ability to Prevent Infections	9.2%	27.5%
Expert Opinion with Pediatric Specialists	15.0%	45.0%
Total	33.3%*	100%

*Numbers do not add up to 33.3 percent due to rounding.

In pediatric cardiology and heart surgery only, the overall weight for expert opinion was 8.5% and the other two measures included in process (Commitment to Best Practices and Ability to Prevent Infections Throughout Hospital) are each 9.9% of the overall weight. The total weight given to the process component is 28.3%. The other 5 percentage points were added to the outcomes component (worth 38.3% in this specialty only).

As with the other components, individual process measures were normalized before being combined in the Index of Hospital Quality (IHQ). Normalization, as described in *Section IV.B*, transforms a measure's index values into a distribution between 0 and 1 based on the range of possible values. The range of expert opinion scores is from 0% (no nominations) to 100% (every surveyed physician nominated the hospital). Starting with the 2013-14 rankings, the normalized expert opinion score has determined the number of points hospitals received for expert opinion. After log transformation, if the highest expert opinion score in a given specialty is 80, for example, the hospital with that score receives a normalized score of 0.80. Because expert opinion is worth

15% of the overall score, the hospital receives 0.80 x 15, or 12 points, for expert opinion. In past years, hospitals with the highest expert opinion scores received the full point total.

VI. Outcomes

For the Best Hospitals adult specialty rankings, risk-adjusted mortality 30 days after admission is a key outcome measure. Other measures now used by healthcare researchers as quality indicators include readmissions following surgical or hospital discharge, patient functional status (or improvement), infection rates, and medical complications.*****

Because of the absence of comprehensive national sources of pediatric outcomes data comparable to the Medicare Provider Analysis and Review (MedPAR) data used in the adult rankings, outcomes-related data are obtained directly from pediatric hospitals through the Pediatric Hospital Survey. Such data include BSI rates, transplant survival rates, mitigation of adverse events, and surgical outcomes. Other data will be added over time to address the need for relevant outcomes measures and to provide a more complete picture of pediatric hospital care. Measures for the 2020-21 rankings were developed from recommendations by expert advisory panels, as previously described. Details on specific outcomes measures, how they were calculated and how they were scored are provided below.

A. Outcome Measures

Outcome measures are listed below, by specialty. Scoring rules used to assign points to hospitals for these outcomes are also described below. For all outcomes measures, a higher number of points indicates better outcomes (e.g., higher survival, lower mortality, fewer complications).

Cancer

Ability to Prevent Infections in Intensive Care Units (15 points). The rate of infections in intensive care units (ICUs) is considered a good benchmark of patient safety and outcome because such infections in hospital-based care should be minimal. Central line-associated bloodstream infection (CLABSI) rates were calculated as the number of BSIs per 1,000 central-line days during the previous 12 months.

CLABSI (A33) rates were tracked for all pediatric ICUs and all oncology/stem cell transplant patients (B22). Hospitals were rewarded for lower rates of infections.

***** For more information on hospital quality measures and updates on national quality of hospital care initiatives, see reports from the Agency for Healthcare Research and Quality (AHRQ) at <http://www.qualitymeasures.ahrq.gov/> and the Joint Commission at <http://www.jointcommission.org/>.

For pediatric ICU CLABSI rates, hospitals received up to 5 points. Hospitals that report to NHSN and receive a Standardized Infection Ratio (SIR) for their pediatric ICU are scored based on their SIR value and upper bound as follows: 5 points if SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is ≤ 1.5 , 4 points if the SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is > 1.5 , 3 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is ≤ 1.5 , 2 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is > 1.5 , or 1 point if the SIR value is > 1.25 . Hospitals that do not participate in the NHSN SIR program receive points based on their unadjusted CLABSI rates as follows: 1 point for > 2.0 and ≤ 3.0 infections per 1,000 patient days, 2 points for > 1.5 and ≤ 2.0 infections per 1,000 patient days, 3 points for > 1.0 and ≤ 1.5 infections, per 1,000 patient days, 4 points for > 0.5 and ≤ 1.0 infections per 1,000 patient days, or 5 points for ≤ 0.5 infections per 1,000 patient days.

Finally, for oncology/stem cell transplant patients CLABSI rates, hospitals received up to 10 points per group. Hospitals that report to NHSN and receive a Standardized Infection Ratio (SIR) for their oncology/stem cell transplant unit are scored based on their SIR value and upper bound as follows: 10 points if SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is ≤ 1.5 , 8 points if the SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is > 1.5 , 6 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is ≤ 1.5 , 4 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is > 1.5 , or 1 point if the SIR value is > 1.25 . Hospitals that do not participate in the NHSN SIR program receive points based on CLABSI rates: 2 points for > 4.0 and ≤ 6.0 infections per 1,000 patient days, 4 points for > 2.0 and ≤ 4.0 infections per 1,000 patient days, 6 points for > 1.0 and ≤ 2.0 infections per 1,000 patient days, 8 points for > 0.5 and ≤ 1.0 infections per 1,000 patient days, or 10 points for ≤ 0.5 infections per 1,000 patient days.

Ability to Prevent Pressure Injuries (5 points). Hospitals received up to 3 points for lower rates of Stage III, Stage IV and unstageable hospital-acquired pressure injuries (A38.1 and A38.2). For both Stage III and Stage IV, hospitals received 1 point for a pressure ulcer rate of ≤ 0.1 per 100 patient admissions assessed over the last four quarters. For unstageable pressure ulcers, hospitals received 1 point for a pressure ulcer rate of ≤ 0.2 per 100 patient admissions assessed over the last four quarters. Hospitals also received up to 2 points for assessing a larger percentage of their total inpatients for pressure ulcers: 1 point for $\geq 50\%$ or $< 75\%$ and 2 points for $\geq 75\%$.

Five-Year Cancer Survival (15 points). This measure evaluated the percentage of pediatric patients at least 18 months old with all subtypes and risk levels of acute lymphoblastic leukemia (ALL) (B35a), acute myeloid leukemia (AML) (B35b), Stage L1 neuroblastoma (B35c), Stage L2/M neuroblastoma (B35d), and medulloblastoma (B35e) who were alive after 5 years of treatment in the

pediatric cancer program. For each of the five measures, hospitals could receive up to 3 points for having a high percentage of 5-year survivors. For ALL, points were awarded as follows: 1 point for $\geq 70\%$ and $< 85\%$ survival, 2 points for ≥ 85 and $< 95\%$ survival, or 3 points for $\geq 95\%$ survival. For AML and stage L2/M neuroblastoma, points were awarded as follows: 1 point for $\geq 35\%$ and $< 50\%$ survival, 2 points for ≥ 50 and $< 60\%$ survival, or 3 points for $\geq 60\%$ survival. For Stage L1 neuroblastoma, points were awarded as follows: 1 point for $\geq 60\%$ and $< 85\%$ survival, 2 points for ≥ 85 and $< 95\%$ survival, or 3 points for $\geq 95\%$ survival. For medulloblastoma, points were awarded as follows: 1 point for $\geq 70\%$ and $< 80\%$ survival, 2 points for ≥ 80 and $< 90\%$ survival, or 3 points for $\geq 90\%$ survival.

Survival After Bone Marrow Transplant (6 points). This measure assessed the percentage of pediatric patients aged 20 years or younger receiving allogeneic bone marrow (including cord blood and stem cell) transplants (BMTs) in the past 5 years who survived for at least 100 days following transplant (B20.1). Hospitals could receive up to 3 points for survival rates for sibling-matched (HLA-identical) allogeneic transplants (B20.1a, B20.1b): 1 point for $\geq 75\%$ and $< 90\%$ survival, 2 points for $\geq 90\%$ and $< 95\%$ survival, or 3 points for $\geq 95\%$ survival. Hospitals could receive up to 3 points for survival rates for matched unrelated allogeneic transplants (B20.1c, B20.1d): 1 point for $\geq 75\%$ and $< 85\%$ survival, 2 points for $\geq 85\%$ and $< 95\%$ survival, or 3 points for $\geq 95\%$ survival.

Cardiology & Heart Surgery

Ability to Prevent Infections in Intensive Care Units (5 points). The rate was calculated as the number of CLABSI (A33) infections per 1,000 central-line days in pediatric ICUs during the previous 12 months. Hospitals were rewarded for lower rates of infections. Hospitals received up to 5 points. Hospitals that report to NHSN and receive a Standardized Infection Ratio (SIR) for their pediatric ICU are scored based on their SIR value and upper bound as follows: 5 points if SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is ≤ 1.5 , 4 points if the SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is > 1.5 , 3 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is ≤ 1.5 , 2 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is > 1.5 , or 1 point if the SIR value is > 1.25 . Hospitals that do not participate in the NHSN SIR program receive points based on CLABSI rates as follows: 1 point for > 2.0 and ≤ 3.0 infections per 1,000 patient days, 2 points for > 1.5 and ≤ 2.0 infections, 3 points for > 1.0 and ≤ 1.5 infections, 4 points for > 0.5 and ≤ 1.0 infections, or 5 points for ≤ 0.5 infections.

Ability to Prevent Pressure Injuries (5 points). Hospitals received up to 3 points for lower rates of Stage III, Stage IV and unstageable hospital-acquired pressure injuries (A38.1 and A38.2). For Stage III and IV, hospitals received 1 point for having a pressure ulcer rate of ≤ 0.1 per

100 patient admissions assessed over the last four quarters. For unstageable pressure ulcers, hospitals received 1 point for a pressure ulcer rate of ≤ 0.2 per 100 patient admissions assessed over the last four quarters. Hospitals also received up to 2 points for assessing a larger percentage of their total inpatients for pressure ulcers: 1 point for $\geq 50\%$ and $< 75\%$ or 2 points for $\geq 75\%$.

Median Post-Operative Length of Stay (9 points). Hospitals can receive up to 9 points for success in reducing the time that congenital heart patients spend in the hospital related to their heart condition for simple to moderately complex cases. For STAT Level 1 (E45a), hospitals received 3 points for having a median post-operative length of stay of ≤ 3 days, 2 points for a median post-operative length of stay > 3 and ≤ 5 days, and 1 point for a median post-operative length of stay > 5 and ≤ 8 days. For each STAT Level 2 and Level 3 (E45b, E45c), hospitals received 3 points each for a median post-operative length of stay of ≤ 7 days, 2 points each for a median post-operative length of stay > 7 and ≤ 14 days, and 1 point each for a median post-operative length of stay > 14 and ≤ 21 days.

Survival After Congenital Heart Surgery (5 points). Starting with the 2017-18 rankings, hospitals now receive points for risk-adjusted survival after heart surgery. The Society of Thoracic Surgeons (STS) Congenital Heart Surgery Database (CHSD) provides an adjusted mortality rate (AMR) using a mortality risk model that incorporates a hospital's patient mix to adjust scores based on known risk factors such as the patient's age, weight, procedure type, prior cardiothoracic operations, non-cardiac congenital anatomic abnormalities, chromosomal abnormalities, syndromes, and preoperative risk factors^{†††††}.


The adjusted mortality rate (AMR) produced for each hospital estimates what the hospital's mortality rate would be if that hospital's case mix was the same as the overall case mix (across all hospitals). The AMR is calculated as the observed mortality rate divided by the expected mortality rate for that case mix, multiplied by the overall STS mortality rate for all hospitals. Operative mortality is defined in all STS databases as (a) all deaths occurring during the hospitalization in which the operation was performed, plus (b) all deaths, occurring after discharge from the hospital but before the end of the 30th postoperative day^{7,8}. Lower scores indicate lower than expected mortality rates. The STS reports also include confidence intervals surrounding the AMR scores.

In an effort to account for the uncertainty in the AMR measure, the upper limit of the confidence interval was incorporated into the points starting with the 2018-19 rankings. However, the measure in the 2018-19 rankings had a distinct cut point at the national average. This meant that a hospital who had a 4-year combined AMR just higher than the national average, and a confidence interval similar to a hospital with an AMR estimate just lower than the national average received

^{†††††} For more information, please see: <https://publicreporting.sts.org/chsd-risk-model>

different scores. For the 2019-20 rankings, a new method of handling the data was utilized to eliminate this difference in performance scores for hospitals near the national average. To accomplish this, we calculated values of 1 and 2 standard errors from the national average dependent on total patient volume across the 4-years using a funnel plot methodology (see **Figure 2**). The standard errors were calculated using the standard statistical formula of $\sqrt{\frac{p*(1-p)}{n}}$, where n represents total patient volume, and p represents the national AMR value.

Figure 2. Example funnel plot of AMR and patient volume with 1 and 2 standard error lines used for point boundaries



After establishing the ranges for scoring of the AMR using the funnel plot methodology, points were awarded based on the comparison of the hospital’s 4-year combined AMR and CI bounds (E43) in the following manner:

- 5 points were awarded to hospitals who had a 4-year combined AMR lower than 2 standard errors from the national average, or who had an upper CI bound less than the national average;
- 4 points were awarded to hospitals who had a 4-year combined AMR between 1 and 2 standard errors lower than the national average;

- 3 points were awarded to hospitals who had a 4-year combined AMR within 1 standard error of the national average;
- 2 points were awarded to hospitals who had a 4-year combined AMR between 1 and 2 standard errors higher than the national average, or who had a 4-year combined AMR higher than 2 standard errors from the national average but also had a lower CI bound less than the national average; and
- 1 point was awarded to hospitals who had a 4-year combined AMR higher than 2 standard errors from the national average.

Hospitals that do not receive a risk-adjusted mortality rate (AMR) from the STS can receive up to 2.5 points for the measure of patient survival after complex heart procedures. This measure represents the rate of operative mortality (patient deaths) following moderately complex to very difficult heart surgery procedures at pediatric hospitals in the four most recent reporting periods (E42). An overall survival rate was computed based on data from STAT levels 1-5 for the past 4 years. Points were assigned as follows: 0.5 points for operative mortality rates $> 4.0\%$ and $\leq 5.0\%$, 1 point for rates $> 3.0\%$ and $\leq 4.0\%$, 1.5 points for rates $> 2.0\%$ and $\leq 3.0\%$, 2 points for rates $> 1.0\%$ and $\leq 2.0\%$, or 2.5 points for rates $\leq 1.0\%$.

Survival After Certain Complex Heart Procedures (9 points). Hospitals received 9 points for lower rates of reoperation and support after initial surgeries for each of three types of surgeries: Tetralogy of Fallot (TOF) repair (E37.2), Arterial Switch Operations for Transposition of the Great Arteries with intact ventricular septum (TGA, IVS) (E37.3), and Ventricular Septal Defect (VSD) repair surgery (E37.4). For each type, hospitals received 1 point for $> 4\%$ and $\leq 8\%$ reoperation or support after the initial surgery, 2 points for $> 2\%$ and $\leq 4\%$ reoperation or support, or 3 points for $\leq 2\%$ reoperation or support.

Survival After Heart Transplant (6 points). Hospitals received up to 3 points based on the ratio of observed^{#####} to expected survival rates for pediatric patients at 1 and 3 years following heart transplant (6 points total) (E23 and E24). The expected survival rate is calculated from statistical models that take into account various factors of both recipients and donors that affect success. A ratio of observed (unadjusted probability of survival) to expected (adjusted probability of survival) survival rates greater than 1.0 indicates that more patients survived than expected, and a ratio of less than 1.0 indicates that fewer patients survived than expected. Points were awarded for

The SRTR now uses “estimated” rather than “observed” survival in its public reports. This report uses “observed” for consistency with other Best Children’s Hospitals measures.

both 1-year and 3-year ratios as follows: 1 point for ratios ≥ 0.80 and < 0.90 , 2 points for ratios ≥ 0.90 and $< 1.$, or 3 points for ratios $\geq 1.$

Survival After Norwood/Hybrid Surgery (24 points). Hospitals received up to 12 points based on the percentage of patients who received the hybrid or Norwood Stage 1 procedure in the last 4 years and were alive without a heart transplant at 1 year of age (E40.1); the denominator for this calculation includes all patients who received the hybrid or Norwood Stage 1 procedure at each heart center and patients who were in the program but died prior to surgical intervention. Up to 3 points were awarded for each of the four reporting years for 1-year survival rates. Points were awarded as follows: 1 point for survival rates $\geq 25\%$ and $< 50\%$, 2 points for survival rates $\geq 50\%$ and $< 75\%$, or 3 points for survival rates $\geq 75\%$. Hospitals received up to 12 points based on the percentage of patients who received the hybrid or Norwood Stage 1 procedure in the last 4 years and were alive regardless of whether they had a heart transplant or not at 1 year of age (E40.2); the denominator for this calculation includes all patients who received the hybrid or Norwood Stage 1 procedure at each heart center and patients who were in the program but died prior to surgical intervention. Up to 3 points were awarded for each of the four reporting years for 1-year survival rates. Points were awarded as follows: 1 point for survival rates $\geq 25\%$ and $< 50\%$, 2 points for survival rates $\geq 50\%$ and $< 75\%$, or 3 points for survival rates $\geq 75\%$.

Diabetes & Endocrinology

Success in Hypothyroid Management (6 points). Hospitals received up to 6 points for hypothyroid management (C59, C59.1). Hospitals received points for having a higher percentage of new congenital hypothyroid patients referred at < 21 days of age who received a confirmatory serum TSH $> 50\text{uIU/ml}$ and began thyroid hormone therapy also before 21 days of age. Points were awarded as follows: 1 point for $\geq 90\%$ and $< 95\%$ of patients beginning therapy, 2 points for $\geq 95\%$ and $< 99\%$ of patients beginning therapy, or 3 points for $\geq 99\%$ patients beginning therapy.


Hospitals received up to 3 additional points for having a higher percentage of congenital hypothyroidism patients < 3 years of age at the time of their visit with at least 2 TSH values within normal ranges (C59.1). Points were awarded as follows: 1 point for $\geq 75\%$ and $< 90\%$ of patients with at least 2 TSH values in normal ranges, 2 points for $\geq 90\%$ and $< 95\%$ of patients with at least 2 TSH values in normal ranges, or 3 points for $\geq 95\%$ of patients with at least 2 TSH values in normal ranges.

Success in Managing Diabetes (48 points). This measure evaluated adverse events in Type 1 and Type 2 diabetes outpatients, rate of optimal hemoglobin A1c levels in primary care Type 1 diabetes outpatients, and inpatient admissions for Type 1 and Type 2 primary care diabetes patients.

To increase the statistical reliability of hospitals' scores on this measure, the 2020-21 rankings implemented a funnel plot methodology to assign points to hospitals for each element of the Success in Managing Diabetes measure. To accomplish this, we calculated value of 1 standard error from the national average dependent on patient volume across the 4-years (see **Figure 3**).

The standard errors were calculated using the standard statistical formula of $\sqrt{\frac{p*(1-p)}{n}}$, where n represents patient volume, and p represents the national ratio value of each metric. After establishing the ranges for scoring of using the funnel plot methodology, points were awarded based on the comparison of the hospital's metric to the national average.

Figure 3. Example funnel plot of optimal A1C values for patients 13-17 years of age on private insurance with Type 1 diabetes and patient volume with 1 standard error lines used for point boundaries



Type 1 diabetes patients < 18 years of age, were evaluated to determine the percentage that achieved optimal control (i.e., at or below 7.5%) for two types of payers (private insurance and Medicaid) and three age groups (0-5 years of age—National average: private insurance = 50.87%,

Medicaid = 29.90%; 6-12 years of age—National average: private insurance = 48.00%, Medicaid = 27.79%; and 13-17 years of age—National average: private insurance = 42.86%, Medicaid = 26.60%) in the last calendar year. Increases in A1c values increase the risk of microvascular complications in patients. Hospitals received up to 3 points in each of the six groups (18 points total) for higher percentages of patients with optimal A1c values (C35.1). Points were awarded as follows:

- 3 points were awarded to hospitals who had a percentage of patients with optimal A1c values greater than or equal to the 1 standard error upper bound from the national average;
- 2 points were awarded to hospitals who had a percentage of patients with optimal A1c values greater than or equal to the 1 standard error lower bound and less than the 1 standard error upper bound from the national average;
- 1 point was awarded to hospitals who had a percentage of patients with optimal A1c values less than the 1 standard error lower bound from the national average.

Hospitals that had missing data in one of the six populations, but otherwise had data for all other elements, received a score based on their overall average for that population. For example, if a hospital did not have any patients 0-5 years of age in the private insurance group, their score for that population, is based on their overall percentage of patients with optimal A1c values across the other five populations. Hospitals who did not have any patients in two or more of the six populations, received 0 points for that population.

Successful management of Type 1 and Type 2 diabetes patients is reflected by the type of primary care these patients receive. Hospitals were rewarded for lower incidence of inpatient admissions and visits to the ER/urgent care for diabetes-related causes for two types of payers (private insurance and Medicaid) (C29.2). For inpatient admissions for Type 1 (C29.2c—National average: private insurance = 4.12%, Medicaid = 10.12%) and Type 2 (C29.2d—National average: private insurance = 2.65%, Medicaid = 4.59%) primary care diabetes patients for each insured group, and for ER/urgent care visits for Type 1 (C29.2e—National average: private insurance = 3.56%, Medicaid = 7.91%) and Type 2 (C29.2f—National average: private insurance = 2.96%, Medicaid = 5.48%) primary care diabetes patients, hospitals were awarded up to 3 points for each insured group (private insurance and Medicaid) for up to a total of 24 points. Points were awarded as follows:

- 3 points were awarded to hospitals who had a ratio of inpatient admissions or ER/urgent care visits less than or equal to the 1 standard error lower bound from the national average;

- 2 points were awarded to hospitals who had a ratio of inpatient admissions or ER/urgent care visits greater than the 1 standard error lower bound and less than or equal to the 1 standard error upper bound from the national average;
- 1 point was awarded to hospitals who had a ratio of inpatient admissions or ER/urgent care visits greater than the 1 standard error upper bound from the national average.

Hospitals received up to 3 points for LDL cholesterol management (C41.1) for each insured group (private insurance—National average = 91.99% and Medicaid—National average = 85.77%) for up to a total of 6 additional points. Hospitals were rewarded according to the percentage of patients with LDL cholesterol values less than 130 at the most recent measurement. Points were awarded as follows:

- 3 points were awarded to hospitals who had a percentage of patients with LDL cholesterol values less than 130 at the most recent measurement greater than or equal to the 1 standard error upper bound from the national average;
- 2 points were awarded to hospitals who had a percentage of patients with LDL cholesterol values less than 130 at the most recent measurement greater than or equal to the 1 standard error lower bound and less than the 1 standard error upper bound from the national average;
- 1 point was awarded to hospitals who had a percentage of patients with LDL cholesterol values less than 130 at the most recent measurement less than the 1 standard error lower bound from the national average.

Gastroenterology & GI Surgery

Ability to Prevent Infections in Intensive Care Units (5 points). The rate was calculated as the number of CLABSI (A33) infections per 1,000 central-line days in pediatric ICUs during the previous 12 months. Hospitals were rewarded for lower rates of infections. Hospitals received up to 5 points. Hospitals that report to NHSN and receive a Standardized Infection Ratio (SIR) for their pediatric ICU are scored based on their SIR value and upper bound as follows: 5 points if SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is ≤ 1.5 , 4 points if the SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is > 1.5 , 3 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is ≤ 1.5 , 2 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is > 1.5 , or 1 point if the SIR value is > 1.25 . Hospitals that do not participate in the NHSN SIR program receive points based on their unadjusted CLABSI rates as follows: 1 point for > 2.0 and ≤ 3.0 infections per 1,000 patient

days, 2 points for > 1.5 and ≤ 2.0 infections, 3 points for > 1.0 and ≤ 1.5 infections, 4 points for > 0.5 and ≤ 1.0 infections, or 5 points for ≤ 0.5 infections.

Ability to Prevent Pressure Injuries (5 points). Hospitals received up to 3 points for lower rates of Stage III, Stage IV and unstageable hospital-acquired pressure injuries (A38.1 and A38.2). For Stage III and IV, hospitals received 1 point for having a pressure ulcer rate of ≤ 0.1 per 100 patient admissions assessed over the last four quarters. For unstageable pressure ulcers, hospitals received 1 point for a pressure ulcer rate of ≤ 0.2 per 100 patient admissions assessed over the last four quarters. Hospitals also received up to 2 points for assessing a larger percentage of their total inpatients for pressure ulcers: 1 point for $\geq 50\%$ and $< 75\%$ or 2 points for $\geq 75\%$.

Success of Certain GI-Related Treatments (9 points). This measure comprises of three items: percentage of patients receiving endoscopic procedures with severe complications (D29), percentage of patients receiving successful Kasai procedures (i.e., improvement total in bilirubin, no synthetic dysfunction, no surgical complications and delayed need for liver transplant) within 90 days of birth (D30.1) and percentage of patients treated for inflammatory bowel disease (IBD) experiencing prednisone-free remission (D32 and D33). Up to 3 points were awarded for each type of procedure. For endoscopic procedures, points were awarded for fewer complications as follows: 1 point for $> 3\%$ and $\leq 5\%$ complications, 2 points for $> 1\%$ and $\leq 3\%$ complications, or 3 points for $\leq 1\%$ complications. For Kasai procedure success, points were awarded as follows: 1 point for $\geq 30\%$ and $< 45\%$, 2 points for $\geq 45\%$ and $< 60\%$, or 3 points for $\geq 60\%$. For IBD prednisone-free remission at the most recent visit, points were awarded as follows: 1 point for $\geq 55\%$ and $< 70\%$ success, 2 points for $\geq 70\%$ and $< 80\%$ success, or 3 points for $\geq 80\%$ success.

Survival After Liver Transplant (6 points). Hospitals received up to 3 points based on the ratio of observed^{§§§§§§} to expected survival rates for pediatric patients at 1 and 3 years after isolated liver transplant (6 points total) (D21 and D22). The expected survival rate is calculated from statistical models that take into account various factors of both recipients and donors that affect success. A ratio of observed (unadjusted probability of survival) to expected (adjusted probability of survival) survival rates greater than 1.0 indicates more patients survived than expected, and a ratio of less than 1.0 indicates that fewer patients survived than expected. Points were awarded as follows: 1 point for ratios ≥ 0.80 and < 0.90 , 2 points for ratios ≥ 0.90 and < 1 , or 3 points for ratios ≥ 1 .

§§§§§§ The SRTR now uses “estimated” rather than “observed” survival in its public reports. This report uses “observed” for consistency with other Best Children’s Hospitals measures.

Neonatology

Ability to Prevent Infections in Neonatal Intensive Care Unit (5 points)*****. The rate was calculated as the number of BSIs per 1,000 central-line days during the previous 12 months (F26.1). Hospitals were rewarded for lower CLABSI rates, which is calculated as the number of BSI events divided by the number of central line days and multiplied by 1,000. Hospitals received up to 5 points. Hospitals that report to NHSN and receive a Standardized Infection Ratio (SIR) for their NICU are scored based on their SIR value and upper bound as follows: 5 points if SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is ≤ 1.5 , 4 points if the SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is > 1.5 , 3 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is ≤ 1.5 , 2 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is > 1.5 , or 1 point if the SIR value is > 1.25 . Hospitals that do not participate in the NHSN SIR program receive points based on their unadjusted CLABSI rates as follows: 1 point for > 2.0 and ≤ 3.0 infections per 1,000 patient days, 2 points for > 1.5 and ≤ 2.0 infections, 3 points for > 1.0 and ≤ 1.5 , 4 points for > 0.5 and ≤ 1.0 infections, or 5 points for ≤ 0.5 infections.

Keeping Breathing Tube in Place (5 points). Hospitals were rewarded for having a lower rate of unintended extubations in infants without tracheostomy. The rate was calculated as the number of unintended extubations per 100 patient ventilator days (F32.1). Hospitals received up to 5 points as follows: 1 point for > 3.0 and ≤ 5.0 extubations per 100 days, 2 points for > 2.0 and ≤ 3.0 extubations, 3 points for > 1.0 and ≤ 2.0 extubations, 4 points for > 0.5 and ≤ 1.0 extubations, or 5 points for ≤ 0.5 extubations.

Matching Breast Milk With Correct Infants (4 points). Hospitals were rewarded for having a lower rate of breast milk administration errors, such as a newborn receiving the wrong breast milk. The rate was calculated as the number of breast milk administration errors per 1,000 breast feeding patient days (F10.6). Hospitals received up to 4 points as follows: 1 point for > 1.0 and ≤ 3.0 errors per 1,000 breast milk feeding patient days, 2 points for > 0.5 and ≤ 1.0 errors, 3 points for > 0.25 and ≤ 0.5 errors, or 4 points for ≤ 0.25 errors.

NICU Temperature Management (12 points). Hospitals were rewarded for having success in managing temperature at the time of admission and post-operatively for patients in the NICU. Hospitals received up to 3 points for having a lower percentage of patients with a first recorded NICU temperature of < 36.0 degrees centigrade in two patient populations: infants with an admission weight of ≤ 1500 grams; and infants with an admission weight of > 1500 (6 points total).

***** In the 2017-18 rankings we attempted to address year-to-year variability in the measurement of BSIs, by incorporating data from the last three years of reporting in the Pediatric Hospital Survey. After discussions with hospitals and the working groups, we have returned to awarding points based on only the most recent year of data.

For each of these two categories, hospitals were awarded 3 points for having $\leq 10\%$ of infants with a temperature of < 36.0 degrees centigrade, 2 points for $> 10\%$ and $\leq 20\%$ of infants with a temperature of < 36.0 degrees centigrade, or 1 point for $> 20\%$ and $\leq 40\%$ of infants with a temperature of < 36.0 degrees centigrade. Because temperature management success rates did not differ significantly between inborn infants (i.e., infants delivered in the hospital where the Level IV NICU is located or at a hospital physically connected to it) and outborn infants (i.e., infants born at another facility and requiring vehicle transfer to hospital or infants previously at home), these populations were pooled within each weight category.

Hospitals could receive an additional 3 points based on the admission temperature of infants who were actively cooled during transport for the management of moderate to severe hypoxic ischemic encephalopathy in the past 3 years (F13.2). Hospitals received 3 points if $\leq 10\%$ of infants had an admission temperature < 33.0 degrees centigrade, 2 points if $> 10\%$ and $\leq 20\%$ of infants had an admission temperature < 33.0 degrees centigrade, or 1 point if $> 20\%$ and $\leq 30\%$ of infants had that temperature upon admission. Since this temperature represents overcooling, the goal is to reduce the percentage of patients with admission temperatures of 33.0 centigrade or less.

Hospitals received up to 3 points based the first postoperative temperature within 60 minutes of return to the NICU after surgery (F31.1). Hospitals received points based on the percent of infants with the first postoperative temperature < 36.0 degrees centigrade. Points were awarded as follows: 3 points if $\leq 5\%$ of infants had the first postoperative temperature < 36.0 degrees centigrade, 2 points if $> 5\%$ and $\leq 10\%$ of infants had the first postoperative temperature < 36.0 degrees centigrade, or 1 point if $> 10\%$ and $\leq 15\%$ of infants had the first postoperative temperature < 36.0 degrees centigrade.

Taking Breast Milk When Discharged (3 points). Hospitals were rewarded for having higher rates of infants admitted at less than 7 days of age being discharged to home, foster care, or a chronic care facility from the NICU (before 120 days) on partial or full mother's own milk (F10.1). Points were awarded as follows: 1 point for $> 0\%$ and $< 60\%$, 2 points for $\geq 60\%$ and $< 80\%$, or 3 points for $\geq 80\%$.

Nephrology

Ability to Prevent Biopsy-Related Complications (6 points). This item measures the percentage of native kidney percutaneous biopsy procedures (G14) and percutaneous kidney transplant biopsies (G27) that resulted in a biopsy complication requiring admission, readmission or a lengthened stay (G15 and G27.2). For both rates, hospitals receive more points for having lower complication rates, as follows: 1 point for complication rates $> 5\%$ and $\leq 10\%$, 2 points for complication rates $> 2\%$ and $\leq 5\%$, or 3 points for complication rates $\leq 2\%$.

Ability to Prevent Dialysis-Related Infections (9 points). Hospitals received 6 points based on a lower peritonitis rate (months of dialyses/cases of peritonitis) for pediatric outpatients on maintenance peritoneal dialysis (G24.1). For each of the past 2 years, up to 3 points were awarded: 1 point for a peritonitis rate of ≥ 1 and < 10 months between peritonitis cases, 2 points for a rate of ≥ 10 and < 20 months between cases, or 3 points for a rate of ≥ 20 months between cases.

Hospitals could receive an additional 3 points for having lower hemodialysis catheter-associated BSIs for outpatients on maintenance hemodialysis in the last 2 years (G37). Hospitals received points for each year as follows: 1 point for ≥ 4.0 and < 8.0 infections per 100 patient months, and 2 points for ≥ 2.0 and < 4.0 infections, or 3 points for < 2.0 infections.

Ability to Prevent Infections in Intensive Care Units (5 points). The rate was calculated as the number of CLABSI (A33) infections per 1,000 central-line days in pediatric ICUs during the previous 12 months. Hospitals were rewarded for lower rates of infections. Hospitals received up to 5 points. Hospitals that report to NHSN and receive a Standardized Infection Ratio (SIR) for their pediatric ICU are scored based on their SIR value and upper bound as follows: 5 points if SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is ≤ 1.5 , 4 points if the SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is > 1.5 , 3 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is ≤ 1.5 , 2 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is > 1.5 , or 1 point if the SIR value is > 1.25 . Hospitals that do not participate in the NHSN SIR program receive points based on their unadjusted CLABSI rates as follows: 1 point for > 2.0 and ≤ 3.0 infections per 1,000 patient days, 2 points for > 1.5 and ≤ 2.0 infections, 3 points for > 1.0 and ≤ 1.5 infections, 4 points for > 0.5 and ≤ 1.0 infections, or 5 points for ≤ 0.5 infections.

Ability to Prevent Pressure Injuries (5 points). Hospitals received up to 3 points for lower rates of Stage III, Stage IV and unstageable hospital-acquired pressure injuries (A38.1 and A38.2). For Stage III and IV, hospitals received 1 point for having a pressure ulcer rate of ≤ 0.1 per 100 patient admissions assessed over the last four quarters. For unstageable pressure ulcers, hospitals received 1 point for a pressure ulcer rate of ≤ 0.2 per 100 patient admissions assessed over the last four quarters. Hospitals also received up to 2 points for assessing a larger percentage of their total inpatients for pressure ulcers: 1 point for $\geq 50\%$ and $< 75\%$ or 2 points for $\geq 75\%$.

Success in Managing Dialysis Patients (12 points). This measure evaluates outcomes for patients on maintenance dialysis during the past 2 calendar years (G23). Hospitals received up to 8 points for higher percentage of patients with these favorable outcomes: monthly Kt/V values of > 1.2 for patients who received hemodialysis three times a week, and percentage of total Kt/V values of ≥ 1.8 for patients receiving peritoneal dialysis. For both outcomes in each of the past 2

years points were awarded as follows: 1 point for desirable outcome rates $\geq 80\%$ and $< 90\%$ or 2 points for desirable outcome rates $\geq 90\%$.

Hospitals received up to an additional 4 points based on the percentage of end-stage renal disease patients receiving hemodialysis and/or peritoneal dialysis for at least 3 consecutive months who survived (G20). Rates were divided into two submeasures based on age: infants and children under 5 years of age and children and adolescents aged 5-19. For children under 5 years of age, up to 2 points per item were awarded: 1 point for survival rates $\geq 80\%$ and $< 90\%$ or 2 points for survival rates $\geq 90\%$. For children and adolescents aged 5-19, points were awarded as follows: 1 point for survival rates $\geq 85\%$ and $< 95\%$ or 2 points for survival rates $\geq 95\%$.

Survival after Kidney Transplant (24 points). Hospitals received up to 24 points based on observed^{††††††††} survival rates at 1 and 3 years of the kidney and of the patient for deceased-donor and living-donor kidney transplants (24 points total) (G32.1b, G32.2b, G32.3b, and G32.4b). A total of eight observed survival rates, each worth up to 3 points were included: 1- and 3-year graft survival rates (deceased donor), 1- and 3-year graft survival rates (living donor), 1- and 3-year patient survival rates (deceased donor), and 1- and 3-year patient survival rates (living donor). Points were awarded in each of the eight groups based on the observed probability of survival (unadjusted probability of survival) as follows: 1 point for rates ≥ 0.50 and < 0.80 , 2 points for rates ≥ 0.80 and < 0.90 , or 3 points for rates ≥ 0.90 .

Neurology & Neurosurgery

Ability to Prevent Surgical Complications (22 points). This measure rewards hospitals for having lower readmission rates. Hospitals received up to 10 points total for having a lower percentage of patients readmitted for any cause within 30 days of the following four surgical procedures: craniotomy (H17a), spinal surgery for dysraphism (H17b), Chiari decompression (H17c), shunt placement (H17d), and intrathecal baclofen pump insertion (H17e). Points were awarded in each group as follows: 1 point for $> 5\%$ and $\leq 15\%$ readmission rate or 2 points for $\leq 5\%$ readmission rate.

Hospitals received up to 3 points for having lower unplanned returns to the operating room within 30 days for patients receiving new/initial neurosurgical shunt placements (H29). Points were awarded as follows: 1 point for $> 5\%$ and $\leq 15\%$ unplanned return rate, 2 points for $> 3\%$ and $\leq 5\%$ unplanned return rate, or 3 points for $\leq 3\%$ unplanned return rate.

^{††††††††} The SRTR now uses “estimated” rather than “observed” survival in its public reports. This report uses “observed” for consistency with other Best Children’s Hospitals measures.

Hospitals received up to 3 points for having a lower percentage of unplanned returns to the operating room within 30 days of receiving a craniotomy (H17.1). Points were awarded as follows: 1 point for > 5% and ≤ 15% readmission rate, 2 points for > 3% and ≤ 5% readmission rate, or 3 points for ≤ 3% readmission rate.

Hospitals received up to 3 points for having a lower complication rate—returns to the OR for unplanned revision surgery within 12 months—for craniofacial procedures performed (H33 and H34). Points were awarded as follows: 1 point for > 5% and ≤ 15% complication rate, 2 points for > 3% and ≤ 5% complication rate, or 3 points for ≤ 3% complication rate.

Hospitals received up to 3 points for having lower 30-day complication rates for epilepsy surgical resection or laser ablation in patients with convulsive disorders (H8 and H8.2). Points were awarded as follows: 1 point for > 5% and ≤ 10% complication rate, 2 points for > 3% and ≤ 5% complication rate, or 3 points for ≤ 3% complication rate.

Success in Controlling Epilepsy (6 points). Hospitals received up to 6 points for the percentage of patients receiving three specific treatments for epilepsy (temporal lobe epilepsy surgery, extra-temporal lobe epilepsy surgery, and functional hemispherectomy) who achieved Engel Class 1 after 12 months. Hospitals were rewarded for higher rates of success. For temporal lobe epilepsy surgery including laser ablation (H31a): 1 point for seizure-free rates ≥ 50% and < 80% or 2 points for seizure-free rates ≥ 80%. For extra-temporal lobe epilepsy surgery including laser ablation (H31b): 1 point for seizure-free rates ≥ 30% and < 60% or 2 points for seizure-free rates ≥ 60%. For functional hemispherectomy (H31c): 1 point for seizure-free rates ≥ 75% and < 90% or 2 points for seizure-free rates ≥ 90%.

Survival After Surgery (12 points). Hospitals received up to 12 points for surgical survival rates for six significant neurological disorders or procedures, including brain tumors (H16a), craniosynostosis (H16b), hydrocephalus patient shunts (H16c), medically intractable epilepsy (H16d), spinal dysraphism (H16e), and Chiari I malformation (H16f). Lower mortality rates indicate better performance (i.e., a lower rate of death following surgery). Points were awarded as follows: 1 point for survival rates ≥ 95% and < 99% or 2 points for survival rates ≥ 99%.

Orthopedics

Ability to Prevent Surgical Complications (15 points). Hospitals received up to 12 points based on the rate of adverse outcomes for patients who received surgical correction for two types of scoliosis: idiopathic scoliosis (I31a) and neuromuscular scoliosis in patients with Cerebral Palsy who have a Gross Motor Function Classification System (GMFCS) function level IV or V (I31b). Two adverse outcomes were measured for both types of scoliosis: unplanned admissions within 30 days

of procedure (for any reason) and reoperation (for any cause) within 90 days (I32). Hospitals who had 40 or more idiopathic scoliosis patients or 20 or more neuromuscular scoliosis patients with Cerebral Palsy received up to 3 points in each of the four categories, with more points for better performance (i.e., lower levels of adverse events): 1 point for complication rate $> 7\%$ and $\leq 10\%$, 2 points for complication rate $> 3\%$ and $\leq 7\%$ or 3 points for complication rate $\leq 3\%$.

Hospitals received an additional 3 points for rates of allogenic blood transfusions for adolescent patients with idiopathic scoliosis with major Cobb angle of 45-70 degrees who received posterior spinal fusion and instrumentation. Points were awarded as follows: 3 points for $\leq 5\%$ of patients receiving allogenic blood transfusion, 2 points for $> 5\%$ and $\leq 10\%$ of patients receiving allogenic blood transfusion, or 1 point for $> 10\%$ and $\leq 15\%$ of patients receiving allogenic blood transfusion.

Speed and Success in Treating Complex Fractures (8 points). Hospitals received up to 4 points for having a higher percentage of patients with an operating room start time within 18 hours of check-in with the Emergency Department for two conditions: operative reduction and fixation of supracondylar fracture (I25) of the humerus and femoral shaft fracture (I26). Hospitals received points for supracondylar fractures as follows: 1 point for $\geq 75\%$ and $< 90\%$ of patients with operating room start times within 18 hours or 2 points for $\geq 90\%$. Hospitals received points for femoral shaft fractures as follows: 1 point for $\geq 60\%$ and $< 80\%$ of patients with operating room start times within 18 hours or 2 points for $\geq 80\%$. Hospitals received an additional 2 points for fewer procedures performed on patients with supracondylar fractures using a formal open procedure (I25.1/I25): 2 points for $\leq 5\%$, or 1 point for $> 5\%$ and $\leq 10\%$.

Hospitals received up to 2 additional points for conducting radiographically-assisted reductions (without requiring hospital admission) of displaced forearm fractures in patients under 14 years of age (I27). Hospitals received points as follows: 1 point for $\geq 70\%$ and $< 90\%$ of patients without requiring hospital admission or 2 points for $\geq 90\%$.

Pulmonology & Lung Surgery

Ability to Prevent Infections in Intensive Care Units (5 points). The rate was calculated as the number of CLABSI (A33) infections per 1,000 central-line days in pediatric ICUs during the previous 12 months. Hospitals were rewarded for lower rates of infections. Hospitals received up to 5 points. Hospitals that report to NHSN and receive a Standardized Infection Ratio (SIR) for their pediatric ICU are scored based on their SIR value and upper bound as follows: 5 points if SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is ≤ 1.5 , 4 points if the SIR value is ≤ 0.75 and the upper bound of the 95% confidence interval is > 1.5 , 3 points if the SIR value is ≤ 1.25 and the upper bound of the 95% confidence interval is ≤ 1.5 , 2 points if the SIR value is ≤ 1.25 and

the upper bound of the 95% confidence interval is > 1.5 , or 1 point if the SIR value is > 1.25 . Hospitals that do not participate in the NHSN SIR program receive points based on unadjusted CLABSI rates as follows: 1 point for > 2.0 and ≤ 3.0 infections per 1,000 patient days, 2 points for > 1.5 and ≤ 2.0 infections, 3 points for > 1.0 and ≤ 1.5 infections, 4 points for > 0.5 and ≤ 1.0 infections, or 5 points for ≤ 0.5 infections.

Ability to Prevent Pressure Injuries (5 points). Hospitals received up to 3 points for having lower rates of Stage III, Stage IV and unstageable hospital-acquired pressure injuries (A38.1 and A38.2). For Stage III and IV, hospitals received 1 point for having a pressure ulcer rate of ≤ 0.1 per 100 patient admissions assessed over the last four quarters. For unstageable pressure ulcers, hospitals received 1 point for a pressure ulcer rate of ≤ 0.2 per 100 patient admissions assessed over the last four quarters. Hospitals also received up to 2 points for assessing a larger percentage of their total inpatients for pressure ulcers: 1 point for $\geq 50\%$ and $< 75\%$ or 2 points for $\geq 75\%$.

Success in Managing Cystic Fibrosis Patients (19 points). Hospitals received up to 14 points for representing better outcomes for patients with cystic fibrosis. Hospitals received up to 12 points (3 points for each item) for improving the functional status of cystic fibrosis patients' median body mass index (BMI) for patients 2-19 (J24b), median forced expiratory volume (FEV₁) for patients 6-12 (J24c), median forced expiratory volume (FEV) for patients 13-17 (J24d), the percentage of children 7-17 who met treatment guidelines for CF patients (at least four outpatient visits, one culture, two spirometries or PFTs) (J24e), and median weight-for-length percentile for CF patients 24 months of age or less (J24f). More points indicate better outcomes or better functional status. For BMI, points were awarded as follows: 1 point for median BMI percentile ≥ 40 and $< 45\%$, 2 points for median BMI percentile $\geq 45\%$ and $< 50\%$, or 3 points for median BMI percentile $\geq 50\%$. For the FEV₁ measure, points were awarded as follows: 1 point for median FEV₁ ≥ 80 and $< 90\%$, 2 points for median FEV₁ $\geq 90\%$ and $< 100\%$, or 3 points for median FEV₁ $\geq 100\%$. For the percentage of children meeting treatment guidelines, points were awarded as follows: 1 point for ≥ 50 and $< 75\%$, 2 points for $\geq 75\%$ and $< 90\%$, or 3 points for median FEV₁ $\geq 90\%$. For median weight-for-length percentile for CF patients 24 months of age or less, points were awarded as follows: 1 point for ≥ 10 and $< 25\%$, 2 points for $\geq 25\%$ and $< 50\%$, or 3 points for median $\geq 50\%$.

Hospitals received up to 2 additional points for meeting performance benchmarks for cystic fibrosis. One point was awarded for having met the benchmark of $< 10\%$ quantity not sufficient (QNS) when conducting pilocarpine iontophoresis (sweat test) for cystic fibrosis with infants (6 weeks-3 months of age) (J21); 1 additional point was awarded for meeting the benchmark of $< 5\%$ QNS for children over 3 months (J22).

Hospitals received up to 2 points for having higher rates of patients over age 10 with cystic fibrosis (not already taking insulin) who completed an oral glucose tolerance test in the previous 12 months (J23). One point was awarded for $\geq 50\%$ and $< 75\%$ of patients completing the test or 2 points were awarded for $\geq 75\%$ of patients completing the test.

Success With Asthma Inpatients (8 points). Success with asthma patients was measured by two factors: shorter inpatient stays and lower readmission rates for asthma-related symptoms. Hospitals were awarded up to 6 points total based on the percentage of asthma inpatients readmitted within 7 days and 30 days for exacerbation of asthma-related symptoms (J13). Hospitals were rewarded for lower percentages of inpatient readmissions: 1 point each for readmission rates $> 3\%$ and $\leq 5\%$, 2 points each for rates $> 1.5\%$ and $\leq 3\%$, or 3 points each for rates $\leq 1.5\%$. Up to 2 additional points are awarded for shorter lengths of stay for asthma inpatients (J12): 1 point for an average stay > 2 days and ≤ 4 days or 2 points for a stay ≤ 2 days.

Survival After Lung Transplant (5 points). Hospitals received up to 5 points based on the observed^{*****} survival rates at 1-year and 3-year for pediatric lung transplant patients (J47 and J48). Points were awarded in each group based on the observed probability of survival (unadjusted probability of survival). For 1-year survival rates, hospitals received 1 point for rates $\geq 50\%$ and $< 65\%$, 2 points for rates $\geq 65\%$ and $< 80\%$, or 3 points for rates $\geq 80\%$. For 3-year survival rates, hospitals received 1 point for rates $\geq 50\%$ and $< 60\%$, or 2 points for rates $\geq 60\%$.

Survival of Patients on Ventilators (3 points). Hospitals received up to 3 points for lower rates of inpatient deaths or cardiorespiratory arrests over the last 3 years for ventilator-dependent patients due to accidental obstruction, decannulation or tracheostomy (J41). Higher survival rates indicate better performance (i.e., lower rate of death of patients on ventilators) and were awarded more points, as follows: 1 point for survival $\geq 95\%$ and $< 97\%$, 2 points for survival $\geq 97\%$ and $< 99\%$, or 3 points for survival $\geq 99\%$.

Urology


Ability to Prevent Surgical Complications (17 points). This measure evaluated a number of complications and adverse outcomes in patients who received urologic surgical procedures in the last 3 years.

In an effort to create more data driven scores, the 2020-21 rankings implemented a funnel plot scoring methodology to assign points to hospitals for each element of the Ability to Prevent

***** The SRTR now uses “estimated” rather than “observed” survival in its public reports. This report uses “observed” for consistency with other Best Children’s Hospitals measures.

Surgical Complications score. To accomplish this, we calculated a value of 1 standard error from the national average dependent on patient volume across the 4-years (see **Figure 4**). The national average is the average rate of all hospitals who submitted the urology section of the survey. The standard errors were calculated using the standard statistical formula of $\sqrt{\frac{p*(1-p)}{n}}$, where n represents patient volume, and p represents the national ratio value of each metric. After establishing the ranges for scoring of using the funnel plot methodology, points were awarded based on the comparison of the hospital's metric to the national average.

Figure 4. Example funnel plot unplanned hospital admissions for urologic issue within 30 days of inpatient urological surgery and surgical volume with 1 standard error lines used for point boundaries



Complications for the following surgical procedures were tracked: distal hypospadias (K15a—National average: 4.26%), pyeloplasty (K15b—National average: 3.27%), and ureteral reimplantation, primary or revision surgery (K15c—National average: 0.73%). Hospitals received up to 9 points total for the three measures, with more points awarded for better performance (i.e., lower complication rates). The points were awarded as follows:

- 3 points were awarded to hospitals who had a complication rate less than or equal to the 1 standard error lower bound from the national average;
- 2 points were awarded to hospitals who had a complication rate greater than the 1 standard error lower bound and less than or equal to the 1 standard error upper bound from the national average;
- 1 point was awarded to hospitals who had a complication rate greater than 1 standard error from the national average.

For adverse events of unplanned hospital admissions for urologic issue within 30 days of inpatient urological surgery (K16.1a—National average: 0.19%) following a scheduled ambulatory urological surgical procedure (K16.1b—National average: 0.20%), hospitals received up to 3 points for each of the three measures (6 points total), with more points awarded for better performance (i.e., lower adverse event rates) as follows:

- 3 points were awarded to hospitals who had an adverse event rate less than or equal to the 1 standard error lower bound from the national average;
- 2 points were awarded to hospitals who had an adverse event rate greater than the 1 standard error lower bound and less than or equal to the 1 standard error upper bound from the national average;
- 1 point was awarded to hospitals who had an adverse event rate greater than 1 standard error from the national average.

For adverse events of unplanned reoperation for a urologic issue within 48 days of urological surgery (K16.1c—National average: 0.06%), hospitals received up to 2 points with more points awarded for better performance (i.e., lower adverse event rates) as follows:

- 2 points were awarded to hospitals who had an adverse event rate less than or equal to the 1 standard error upper bound from the national average;
- 1 point was awarded to hospitals who had an adverse event rate greater than the 1 standard error upper bound from the national average.

Speed in Treating Testicular Torsion (2 points). This measure evaluates how quickly patients who presented with torsion of the testis received care following their registration for care in the ED or outpatient clinic (K19). Hospitals received 1 point for $\geq 50\%$ and $< 90\%$ of patients

having an OR start time < 4 hours following check-in at the hospital or 2 points for $\geq 90\%$ of patients having an OR start time < 4 hours.

B. Normalization and Weighting

As with structural and process measures, individual outcomes measures were normalized to have a distribution between 0 and 1. *Table 16* shows the relative weight of each measure on the total outcomes score for that specialty. The outcome measures combined are worth 33.3% of the overall score in all specialties except for pediatric Cardiology & Heart Surgery, for which outcomes are worth 38.3%. To determine the percent of the overall score for a given measure, divide the individual measure relative weight by the total weight for that specialty and multiply by 33.3 (38.3 in Cardiology & Heart Surgery).

Table 16. Relative Weights of Outcomes Measures by Specialty

Measure	Cancer	Cardiology & Heart Surgery	Diabetes & Endocrinology	Gastroenterology & GI Surgery	Neonatology	Nephrology	Neurology & Neurosurgery	Orthopedics	Pulmonology & Lung Surgery	Urology
Ability to prevent biopsy-related complications						1				
Ability to prevent dialysis-related infections						1				
Ability to prevent infections in intensive-care units	0.38	0.98		0.56	2	0.74			1	
Ability to prevent pressure injuries	0.11	0.29		0.17		0.22			0.30	
Ability to prevent surgical complications							1.25	1		1.75
Five-year cancer survival	1									
Keeping breathing tube in place					1					
Matching breast milk with correct infants					1					
Median post-operative length of stay		1								
NICU temperature management					1					
Speed and success in treating complex fractures								1		
Speed in treating testicular torsion										1
Success in controlling epilepsy							1			
Success in hypothyroid management			1							
Success in managing cystic fibrosis patients									2	
Success in managing diabetes			2							
Success in managing dialysis patients						1				
Success of certain GI-related treatments				2						
Success with asthma inpatients									1.5	
Survival after (bone marrow/heart/kidney/liver/lung) transplant	1	1		1		1			0.5	
Survival after certain complex heart procedures		2								
Survival after congenital heart surgery		5								
Survival after Norwood/hybrid surgery		1								
Survival after surgery							1			
Survival of patients on ventilators									1	
Taking breast milk when discharged					1					
Total	2.49	11.27	3.00	3.73	6.00	4.96	3.25	2.00	6.30	2.75

VII. Calculation of the U.S. News Score

The U.S. News ranking score reflects the followings weights for each of the major components and the individual process measures as shown in **Table 17**. Starting with the 2017-18 rankings, individual component weights differ for Cardiology & Heart Surgery when compared with the other specialties. The differences can be seen in **Table 17**. This shift in scoring is due to the inclusion of risk-adjusted mortality measures, which are more reliable than the observed outcome measures used in the other specialties.

Relative structural measure weights can be found in Table 8, and the relative outcomes measure weights are shown in Table 17.

Table 17. Component Weighting

Component	All Specialties Except Pediatric Cardiology & Heart Surgery	Pediatric Cardiology & Heart Surgery
Structure	33.3%	33.3%
Process: Commitment to Best Practices	9.2%	9.9%
Process: Infection-preventing measures	9.2%	9.9%
Process: Expert opinion	15.0%	8.5%
Outcomes*	33.3%	38.3%

*Numbers do not add up to total due to rounding.

Although each measure represents a specific aspect of quality, a single score provides a result that is easy to use and understand and that portrays overall quality more accurately than any of the three components would individually. The rankings for the top 50 hospitals in each of the pediatric specialties, by U.S. News score, are shown in **Appendix D**. Starting with the 2012-13 rankings, hospitals with the same U.S. News rounded score have been considered to be tied.

The formula for calculating the U.S. News score for a given hospital is shown in Equation (2). The score can be thought of as a simple weighted sum of structural, process and outcome measures as shown below:

Equation (2) $Score = (\sum_{i=1}^{n_s} wts_i * s_i) + (\sum_{i=1}^{n_p} wtp_i * p_i) + (\sum_{i=1}^{n_o} wto_i * o_p)$,
 where

- $Score$ = raw hospital score in a given specialty,
- wts_i = weight assigned to structure measure i ,
- wtp_i = weight assigned to process (expert opinion) measure i ,
- wto_i = weight assigned to outcomes measure i ,
- s_i = normalized value for structural measure i ,
- p_i = normalized value for process measure i ,
- o_i = normalized value for outcomes measure i .

Please note that the formula is meant for illustrative purposes only; it *cannot* be used to directly calculate a score for an individual hospital. For presentation purposes, raw scores are transformed to a scale that assigns a U.S. News score of 100 to the top hospital. The formula for the transformation is shown in Equation (3):

Equation (3) U.S. News $Score = (score - minimum) / range$

VIII. Pediatric Honor Roll

In all, 88 different hospitals were ranked in at least one pediatric specialty in the 2020-21 rankings. The Best Children’s Hospitals Honor Roll, established in 2009, recognizes excellence across a broad range of pediatric specialties.

In each specialty, the No. 1-ranked hospital received 25 Honor Roll points and lower-ranked hospitals received progressively fewer points – the No 2 hospital receiving 24 points, the No. 3 hospital 23 points, and so on – with all hospitals ranked 21-50 receiving 5 points. A hospital ranked No. 1 in all 10 specialties would therefore have received 250 points. The 2020-21 Honor Roll recognizes the 10 hospitals that earned the most points out of 250 across the 10 specialties. The Honor Roll is ordered 1 through 10 based on total points (see Appendix E).

IX. 2020-21 Changes

- **Maintaining survey stability.** This year, U.S. News chose to make a limited number of changes in response to feedback from the working groups and hospitals. This was done to minimize burden to hospitals responding to the survey who must update their reporting each year in response to changes.
- **Providing guidance and requesting additional data documentation.** As with the past two surveys, U.S. News requested that hospitals provide deidentified data to support responses to key volume and outcome questions in the endocrinology and urology sections of the survey. In 2020, the survey also provided more explicit guidance to hospitals on how to pull data from their systems to answer questions. The survey also asked that hospitals provide some additional information about how they pulled numbers such as the sources of data, procedures, and data cleaning processes.
- **Outcomes.** Several changes were introduced in the rankings this year in outcomes as described below.
 - ***Funnel plot scoring.*** A new method of calculating scores for certain outcomes was expanded in the 2020-21 rankings to improve the reliability of hospitals' scores on these outcomes. The funnel plot methodology utilizes the available data to determine how well hospitals performed in avoiding negative outcomes such as death and complications. Both the rate of these events occurring as well as the volume of cases were taken into consideration when assigning points. In the 2019-20 rankings we introduced funnel plot scoring to the "Risk-adjusted survival after heart surgery" measure. This year, the funnel plot scoring approach was applied to the "Success in managing diabetes" measure in pediatric Diabetes and Endocrinology and the "Ability to prevent surgical complications" measure in pediatric Urology.
 - ***Temperature management in Neonatology.*** In recent years, a variety of process and outcome measures focused on temperature management in neonates have become common in Level IV NICUs. The rankings introduced new questions related to temperature management and reclassified several questions that had been used as process measures to be a part of a new outcome focused on temperature management.
 - ***Median post-operative length of stay in Cardiology and Heart Surgery.*** A new measure was introduced focusing on the success hospitals have in minimizing the time that congenital heart patients spend in the hospital

related to their heart condition for simple to moderately complex cases (i.e., STAT1 to STAT3). To accommodate this measure, we have reduced the weight assigned to the AMR.

- **Scoring CLABSI scoring.** The NHSN reporting program now includes tracking and reporting of risk-adjusted rates for CLABSI infections. The Standardized Infection Ratio (SIR) for CLABSI is available for most hospitals and represents a significant improvement in the ability to evaluate performance by hospitals in reducing infections in inpatient ICU settings. Hospital CLABSI SIR values were used when they were available; in cases where they were not available, standard CLABSI rates per 1,000 patient days were used.
- **Weighting.** With the changes introduced above related to new measures in Cardiology and Heart Surgery and Neonatology, the project has introduced minor weighting adjustments. The majority of weights for measures in the rankings in 2020-21 are consistent with their weight in the 2019-20 rankings.

X. Future Improvements

Continued refinements are anticipated during the next few years. They are likely to include the following:

- **Addressing the impact of COVID-19 on hospital care.** Future data collection and analysis will take consideration of the impact of the novel coronavirus pandemic on patients and hospital operations. The Best Children’s Hospitals project will seek input from pediatric clinicians, hospital leaders and measurement experts prior to determining necessary modifications to data collection and analysis. The working groups will also be asked to provide input on the survey measures and scoring values in preparation for the 2021 survey.
- **Consider further weighting changes.** We plan to review the survey with the working groups to consider additional changes to the weights used in the rankings to assess hospitals.
- **Expand outcome measures.** We plan, for example, to explore alternatives for collecting additional mortality data, infection rates, patient functional measures, and complication rates.
- **Explore risk adjustment.** We will continue to investigate methods for risk-adjusting pediatric mortality data to better reflect hospital-to-hospital differences in patient mix, severity and comorbidities. As more pediatric-specific databases are developed or further expanded to include more pediatric facilities, we will explore their possible use in creating risk-adjusted outcomes and performance measures of health care.
- **Identify additional structural measures.** External certifications of hospital quality, excellence in specialty areas and awards for high-quality care will be considered for incorporation in the rankings. Additional technologies, teams and practices that define high-quality pediatric services also will be evaluated.
- **Identify opportunities for data validation and auditing.** To ensure the integrity of the data used, the project team plans to continue to explore opportunities for employ data validation and possible auditing techniques to evaluate data submitted by hospitals for consideration in the rankings.

The project team will continue to work with advisory panels of physicians, nurses, hospital quality specialists and other healthcare professionals. RTI and U.S. News are grateful to these expert volunteers. Their recommendations and advice have been invaluable.

XI. Contact Information

We welcome suggestions and questions. Readers and users of the rankings are encouraged to contact the Best Children's Hospitals research team at BestHospitals@rti.org. This report and methodology reports for the adult rankings can be viewed or downloaded online in their entirety from the RTI International Web site at <http://www.rti.org/besthospitals>.

XII. References

1. Donabedian A. Evaluating the quality of medical care. *Milbank Memorial Fund Quarterly*. 1966; 44:166-203.
2. Donabedian A. Promoting quality through evaluating the process of patient care. *Medical Care*. 1968; 6:181.
3. Donabedian A. The quality of care: How can it be assessed? *Journal of the American Medical Association*. 1988; 260:1743-1748.
4. Donabedian A. The seven pillars of quality. *Archives of Pathology and Laboratory Medicine*. 1990; 114:1115-1118.
5. Donabedian A. The role of outcomes in quality assessment and assurance. *Quality Review Bulletin*. 1992; 18(11):356-360.
6. National Center for Health Statistics. *The international classification of diseases, ninth revision, clinical modification (ICD-9-CM)*. Hyattsville, MD: National Center for Health Statistics. Available at <http://www.cdc.gov/nchs/icd/icd9.htm>. Accessed on April 21, 2006.
7. Jacobs JP, Mavroudis C, Jacobs ML, et al. What is operative mortality? Defining death in a surgical registry database: a report from the STS Congenital Database Task Force and the Joint EACTS-STC Congenital Database Committee. *Ann Thorac Surg* 2006;81:1937–41.
8. Overman D, Jacobs JP, Prager RL, et al. Report from The Society of Thoracic Surgeons National Database Work Force: clarifying the definition of operative mortality. *World J Pediatr Congenit Heart Surg* 2013;4:10–2.

Appendix A
2020-21 Best Children's Hospitals
Working Groups and Members

The Pediatric Hospital Survey is informed, updated and enhanced every year with the help of more than 163 volunteer advisers in 12 working groups – one group for each of the 10 Best Children’s Hospitals specialties, plus infection control and radiology. Two senior advisers provide overarching expertise.

The working groups’ input is invaluable. However, U.S. News and RTI International make all final decisions concerning the content and wording of the survey and the analysis of the data it generates. Working group members do not have access to the data provided by participating hospitals and are not asked to endorse the decisions made by U.S. News.

The working group members and their institutions are listed below.

2020-21 Best Children’s Hospitals Working Groups and Advisors

Cancer

Dr. Amy Louise Billett, Dana-Farber/Boston Children's Cancer and Blood Disorders Center
Ms. Kamar Braish, CHOC Children's Hospital
Dr. Tim Cripe, Nationwide Children's Hospital
Dr. Jeffrey Dome, Children's National Medical Center
Dr. Douglas Hawkins, Seattle Children's Hospital
Dr. Jeffrey D. Hord, Akron Children’s Hospital
Dr. Jessica Kandel, University of Chicago, Comer Children's Hospital
Dr. Howard Katzenstein, Wolfson Children's Hospital
Dr. Eugenie Kleinerman, M.D. Anderson Children's Cancer Center
Dr. Andrew Kung, Memorial Sloan Kettering Cancer Center
Dr. Ellis Neufeld, St. Jude Children's Research Hospital
Dr. Rita Secola, Children’s Hospital Los Angeles
Dr. Sheri Spunt, Lucile Packard Children’s Hospital at Stanford University
Ms. Sherri Ward, Children's Hospital Colorado

Cardiology & Heart Surgery

Dr. Eric J. Devaney, Rainbow Babies & Children's Hospital
Dr. Charles Fraser, Jr., Dell Seton Medical Center at The University of Texas
Dr. Richard Friedman, Cohen Children's Medical Center
Dr. Jeffrey Jacobs, Professor of Surgery, University of Florida
Dr. Kathy Jenkins, Boston Children’s Hospital
Ms. Kristen Johnston, Cook Children's Medical Center
Dr. Richard Jonas, Children's National Medical Center
Dr. Jennifer Li, Duke Children's Hospital and Health Center
Dr. Bradley Marino, Ann & Robert H. Lurie Children's Hospital of Chicago
Dr. Sara Pasquali, C.S. Mott Children's Hospital
Dr. Christian Pizarro, Alfred I. duPont Children Hospital
Dr. Phil Saul, WVU Healthcare Children's Hospital
Dr. David Vener, Texas Children's Hospital
Dr. Robert Vincent, Children's Healthcare of Atlanta, Sibley Heart Center Cardiology
Dr. Karl F. Welke, Levine Children's Hospital

Diabetes & Endocrinology

Dr. Mark Clements, Children's Mercy Hospitals and Clinics
Dr. Fran Cogen, Children's National Medical Center
Dr. Mark Daniels, CHOC Children's Hospital
Dr. Emily Germain-Lee, Connecticut Children's Medical Center
Dr. Roy Kim, Cleveland Clinic Children's Hospital
Dr. Kenneth McCormick, Children's of Alabama, University of Alabama at Birmingham
Dr. Mary Murray, Primary Children's Hospital
Dr. David Repaske, University of Virginia Children's Hospital
Dr. Henry Rohrs, UF Health Shands Children's Hospital
Ms. Lydia Smith, Le Bonheur Children's Hospital
Dr. Anna Spagnoli, Children's Hospital & Medical Center
Dr. Doris Taha, Children's Hospital of Michigan
Dr. Steven Willi, Children's Hospital of Philadelphia

Gastroenterology & GI Surgery

Dr. Sari Acra, Monroe Carell Jr Children's Hospital at Vanderbilt
Dr. Bradley Barth, University of Texas Southwestern Medical Center/Children's Health
Dr. Mitchell B. Cohen, Children's of Alabama, University of Alabama at Birmingham
Dr. Eunice Huang, Le Bonheur Children's Hospital
Dr. Jeffrey Linzer Sr., Children's Healthcare of Atlanta
Ms. Macey Perkins, Dayton Children's Hospital
Dr. Timothy Sentongo, University of Chicago, Comer Children's Hospital
Dr. Thomas Sferra, Rainbow Babies & Children's Hospital
Dr. Mitchell Shub, Phoenix Children's Hospital
Dr. Robbyn Sockolow, Phyllis and David Komansky Center for Children's Health
Dr. Manu Sood, Children's Hospital of Wisconsin
Dr. Daniel von Allmen, Cincinnati Children's Hospital Medical Center
Dr. Barry K. Wershil, Ann & Robert H. Lurie Children's Hospital of Chicago

Infection Control

Dr. Margaret Aldrich, Children's Hospital at Montefiore Medical Center
Dr. Mallory Davis, Helen DeVos Children's Hospital at Spectrum Health
Dr. Sandra L. Fowler, Medical University of South Carolina Children's Hospital
Ms. Wendi Gornick, CHOC Children's Hospital
Dr. Judith Guzman-Cottrill, Doernbecher Children's Hospital
Ms. Michele Honeycutt, Arkansas Children's Hospital
Dr. W. Charles Huskins, Mayo Clinic Children's Center
Mr. Raed Khoury, Valley Children's Healthcare
Dr. Aaron Milstone, Johns Hopkins Children's Center
Ms. Elaine Whaley, Texas Children's Hospital
Ms. Jane Zbinden, St. Louis Children's Hospital
Dr. Danielle Zerr, Seattle Children's Hospital

Neonatology

Dr. Rishi Bakshi, C.S. Mott Children's Hospital
Dr. Beverly Brozanski, Children's Hospital of Pittsburgh, UPMC
Dr. David Burchfield, UF Health Shands Children's Hospital
Dr. F. Sessions Cole, St. Louis Children's Hospital
Dr. Jacquelyn Evans, Children's Hospital of Philadelphia
Dr. Theresa Grover, Children's Hospital Colorado
Dr. Andrew Herman, Levine Children's Hospital
Dr. Jeffrey Kaiser, Penn State Children's Hospital
Dr. Edward Liechty, Riley Hospital for Children at IU Health
Dr. Kris Reber, Nationwide Children's Hospital
Dr. Yao Sun, UCSF Benioff Children's Hospital
Dr. Jonathan Swanson, University of Virginia Children's Hospital
Dr. Michael Uhing, Children's Hospital of Wisconsin
Dr. Barry Weinberger, Cohen Children's Medical Center

Nephrology

Dr. Steve Alexander, Lucile Packard Children's Hospital at Stanford University
Dr. Paul Brakeman, UCSF Benioff Children's Hospital
Dr. Michael Braun, Texas Children's Hospital
Dr. Vikas Dharnidharka, St. Louis Children's Hospital
Dr. Joseph Flynn, Seattle Children's Hospital
Dr. Jens Goebel, Children's Hospital Colorado
Dr. Stuart Goldstein, Cincinnati Children's Hospital Medical Center
Dr. Chryso Katsoufis, Holtz Children's Hospital
Dr. Katherine MacRae Dell, Cleveland Clinic Children's Hospital
Dr. Tej Mattoo, Children's Hospital of Michigan
Dr. Jeffrey M. Saland, Mount Sinai Kravis Children's Hospital
Dr. Vijeta Salunkhe, Baylor Scott and White McLane Children's Hospital
Dr. Michael Somers, Boston Children's Hospital
Ms. Colleen Tipping, Children's Hospital of Pittsburgh, UPMC
Dr. Brad Warady, Children's Mercy Hospitals and Clinics

Neurology & Neurosurgery

Dr. Joshua Chern, Children's Healthcare of Atlanta
Dr. Bruce Cohen, Akron Children's Hospital
Dr. Mark Dias, Penn State Children's Hospital
Dr. Daniel Fain, Helen DeVos Children's Hospital at Spectrum Health
Dr. Mark D. Krieger, Children's Hospital Los Angeles
Dr. David Limbrick, St. Louis Children's Hospital
Dr. Tobias Loddenkemper, Boston Children's Hospital
Dr. Mohamad Mikati, Duke Children's Hospital and Health Center
Dr. Karin Muraszko, C.S. Mott Children's Hospital
Dr. Steve Roach, Dell Seton Medical Center at The University of Texas
Mr. Paul Sweitzer, Connecticut Children's Medical Center
Dr. Bradley Weprin, Children's Medical Center Dallas
Dr. Angus Wilfong, Phoenix Children's Hospital

Orthopedics

Dr. Laurel Blakemore, UF Health Shands Children's Hospital
Dr. Brian Brighton, Levine Children's Hospital
Dr. Kristen Carroll, Primary Children's Hospital/Shriners Hospital for Children Salt Lake City
Dr. Sumeet Garg, Children's Hospital Colorado
Dr. Ryan Goodwin, Cleveland Clinic Children's Hospital
Dr. J Eric Gordon, St. Louis Children's Hospital
Ms. Patricia Gust, St. Jude Children's Research Hospital
Dr. Gregory Hahn, Johns Hopkins All Children's Hospital
Dr. Lisa Ipp, Hospital for Special Surgery
Dr. Charles Mehlman, Cincinnati Children's Hospital Medical Center
Dr. Greg Mencio, Monroe Carell Jr Children's Hospital at Vanderbilt
Dr. Karl Rathjen, Texas Scottish Rite Hospital for Children-Children's Medical Center Dallas
Dr. Coleen Sabatini, UCSF Benioff Children's Hospital
Dr. Greg Schmale, Seattle Children's Hospital
Dr. David Skaggs, Children's Hospital Los Angeles

Pulmonology & Lung Surgery

Ms. Tiruayer Battle, Children's Hospital of Philadelphia
Dr. Scott Bickel, Norton Children's Hospital
Dr. John Carl, Cleveland Clinic Children's Hospital
Dr. Shailendra Das, Texas Children's Hospital
Dr. Charles Esther, North Carolina Children's Hospital at UNC
Dr. Thomas Lahiri, University of Vermont Children's Hospital
Dr. Brian McGinley, Primary Children's Hospital
Dr. Christopher Oermann, Children's Mercy Hospitals and Clinics
Dr. Jonathan Popler, Children's Healthcare of Atlanta
Dr. Jessica Rice, Johns Hopkins Children's Center
Dr. Kristie Ross, Rainbow Babies & Children's Hospital
Dr. Julie Ryu, Rady Children's Hospital-San Diego
Dr. Gregory S. Sawicki, Boston Children's Hospital
Dr. Pornchai Tirakitsoontorn, CHOC Children's Hospital

Radiology

Dr. Richard Barth, Lucile Packard Children's Hospital at Stanford University
Dr. Brian Coley, Cincinnati Children's Hospital Medical Center
Dr. Matthew Cooper, Riley Hospital for Children at Indiana University Health
Dr. Kassa Darge, Children's Hospital of Philadelphia
Dr. Jim Donaldson, Ann & Robert H. Lurie Children's Hospital of Chicago
Dr. Azam Eghbal, CHOC Children's Hospital
Dr. Sheila Moore, Children's Hospital of Pittsburgh, UPMC
Dr. Douglas Rivard, Children's Mercy Hospitals and Clinics
Dr. Michael Rubin, Akron Children's Hospital
Dr. Victor Seghers, Texas Children's Hospital
Dr. Stephen Simoneaux, Children's Healthcare of Atlanta

Urology

Dr. Glenn Cannon, Children's Hospital of Pittsburgh, UPMC

Dr. Patrick Cartwright, Primary Children's Hospital

Dr. Pasquale Casale, AdventHealth for Children

Ms. Kimberley Giles, Le Bonheur Children's Hospital

Dr. Nicholas Holmes, Rady Children's Hospital-San Diego

Dr. Micah Jacobs, University of Texas Southwestern Medical Center/Children's Health

Dr. Paul Merguerian, Seattle Children's Hospital

Dr. Natalie Plachter, Children's Hospital of Philadelphia

Dr. Pramod Reddy, Cincinnati Children's Hospital Medical Center

Dr. Jonathan Routh, Duke Children's Hospital and Health Center

Dr. Stacey Tanaka, Monroe Carell Jr Children's Hospital at Vanderbilt

Dr. Thomas Vates, K. Hovnanian Children's Hospital at Jersey Shore University Hospital

Dr. Vijaya Vemulakonda, Children's Hospital Colorado

Senior Advisors

Dr. Elaine Cox, Riley Hospital for Children at IU Health

Dr. Phil Saul, WVU Healthcare Children's Hospital

Appendix B
Glossary of Terms

Continuous video EEG monitoring with pediatric neurology support (F12a). EEG (electroencephalography) is a technology for measuring electrical activity produced by the brain, as recorded from electrodes placed on the scalp. EEG monitoring provides the ability to collect the brain's electrical activity continuously to help detect and diagnose neurological problems.

Cryoablation (B8f, E14b). This process uses cooled, thermally conductive gases and fluids circulated through hollow needles (cryoprobes) that are placed in contact with or inserted into diseased tissue to kill it.

EEG source localization (H5b). Source localization is the process of identifying the origin or site of seizure activity within the brain. The most common methods of doing this are the use of magnetoencephalography or EEG testing techniques.

Fast magnetic resonance imaging (MRI) shunt for hydrocephalus (F10i). Fast MRI shunt scans are exams performed in under 10 minutes as an alternative to CT scans to assess ventricular size when shunt tube malfunction is suspected.

Functional magnetic resonance (fMR) (B8b). fMR is a specialized type of MRI scan that measures changes in blood flow related to neural activity.

Genetic testing/counseling (A7d). A genetic testing/counseling service is equipped with the appropriate laboratory facilities and is directed by a physician qualified to advise parents and prospective parents on potential problems in cases of genetic defects. A genetic test is the analysis of human DNA, RNA, chromosomes, proteins and certain metabolites to detect heritable disease-related genotypes, mutations, phenotypes or karyotypes for clinical purposes. Genetic tests can have diverse purposes, including the diagnosis of genetic diseases in newborns, children and adults; the identification of future health risks; the prediction of drug responses; and the assessment of risks to future children.

Image-guided radiation therapy (IGRT) (A10e). IGRT is an automated system that produces high-resolution x-ray images to pinpoint tumor sites, adjust patient positioning and generally make treatment more effective and efficient.

Intensity-modulated radiation therapy (IMRT) (A10f). IMRT is a three-dimensional radiation therapy that improves the targeting of treatment delivery in a way that is likely to decrease damage to normal tissues and allows for varying intensities.

Intraoperative magnetic resonance imaging (ioMRI) (A10c). ioMRI uses a uniform magnetic field and radio frequencies to study tissue and structure of the body. It enables visualization of biochemical cellular activity in vivo without the use of ionizing radiation, radioisotopes or ultrasound.

Magnetic resonance cholangiopancreatography (MRCP) (D7a). MRCP is a noninvasive approach for imaging the biliary and pancreatic ducts using MRI.

Multidisciplinary pediatric acute pain/sedation service available onsite 24 hours a day (A8d). This service provides monitored anesthesia care and sedation within the hospital (but not within an operating room or PICU), as well as emergency airway management and acute and chronic pain management for neonates and pediatric patients on a 24-hour basis. A qualified program must

have at least an identified medical director (e.g., general pediatrician, pediatric subspecialist or anesthesiologist) with documented education in conscious sedation and a registered nurse coordinator (or pain management clinical nurse specialist).

Neonatal intensive care unit (NICU) (A7a). A NICU provides mechanical ventilation, neonatal surgery and special care for the sickest infants, including those with the lowest birth weights (below 1,500 grams), who are born in the hospital or transferred from another institution. The NICU is separate from the newborn nursery. A full-time neonatologist serves as director.

Neurophysiological intraoperative monitoring (H5a). This uses electrophysiological methods, including electroencephalography and electromyography, to monitor parts of the brain, spinal cord and peripheral nerves during surgery.

Palliative care program (A7e). A palliative care program is organized and staffed for children nearing the end of life or living with lifespan-limiting conditions. The program's purpose is to minimize pain and discomfort, provide emotional and spiritual support for children and their families, assist with financial guidance and social services and support decision making. Programs must include at least one physician providing direct patient care; a nurse coordinator; and a social worker, certified child-life specialist or pastoral counselor. All program staff must have training in palliative care.

Patient care rooms with protective environment (A7c). The Protective Environment incorporates the following: air exchanges ≥ 12 per hour; central or point-of-use high-efficiency particulate (HEPA) filters, consistent positive air pressure differentials between the patient's room and hallway and continuous monitoring of pressure differentials.

Pediatric anesthesia program available onsite 24 hours a day (A8a). This team provides anesthesia care for children before, during and after surgery (or other medical procedures). The team provides 24-hour coverage by board-certified anesthesiologists who specialize in pediatric anesthesia.

Pediatric intensive care unit (PICU) (A7b). A PICU is staffed with specially trained personnel and has monitoring and specialized support equipment for treating pediatric patients who, because of shock, trauma or other life-threatening conditions, require intensified, comprehensive observation and care.

Pediatric pain management program available onsite 24 hours a day (A8c). Administered by specially trained physicians and other clinicians, this is a recognized clinical service or program providing specialized medical care, drugs or therapies for the management of acute or chronic pain and other distressing symptoms among children suffering from an acute illness of diverse causes.

PET/computed tomography (PET/CT) scanning (A10b). PET/CT combines the capabilities of PET and CT scanning into a single, integrated device that provides metabolic functional information for monitoring chemotherapy, radiotherapy, and surgical planning.

Portable CT scanning unit (A10g). CT scanning unit that can be moved to where patient care is being provided rather than having a fixed unit in a single location. The portable unit is particularly helpful in delivering care in the ICU, emergency department, and in operating room environments.

Positron emission tomography (PET) (A10a). PET scanning is a computerized nuclear medicine imaging technology that uses radioactive (positron-emitting) isotopes created in a cyclotron or

generator to produce composite images of the brain and heart activity. The scans are sectional images depicting metabolic activity or blood flow rather than anatomy.

Radiofrequency ablation (B8f, E14c). This procedure involves placing probes that emit radiofrequency energy into the heart using a catheter. The radiofrequency energy is then used to destroy abnormal electrical activity in the heart tissue.

Rapid response team (A8a). A rapid response team, also known as a medical emergency team, is distinct from the hospital “code” team. It is available 24 hours a day and has three essential characteristics: (1) the team creates tools and provides staff education for recognizing an acute deterioration in patient condition; (2) the team follows the SBAR (situation, background, assessment, recommendation) method to communicate such a change in condition effectively and efficiently (i.e., escalation policy); and (3) the team responds to the change in condition with the goal of reducing/eliminating preventable “codes.”

Rehabilitation program and consultation service (A7f). This program provides either a rehabilitation unit and/or a consultation service within the pediatric program for patients requiring rehabilitation. The program must include a pediatric physiatrist (board certified/board eligible pediatric rehabilitation physician) as the director.

Therapeutic meta-iodine-benzyl-guanidine with I-131 radionuclide (I-131 MIBG) (B8a). I-131MIBG is a functional imaging and treatment agent used to help locate, diagnose and treat tumors of adrenergic tissues, such as neuroblastoma and pheochromocytoma. For this question, we are only interested in therapeutic use of I-131 MIBG to treat cancer.

Three-dimensional mapping (E14a). This includes the use of three-dimensional imaging systems such as MRI or ultrasound to guide ablation probes.

Vascular tumor program (A35). This program brings together a multidisciplinary team of specialists to diagnose and ensure the most effective treatment for optimal functioning and quality of life for children with vascular anomalies (tumors or malformations). To be eligible, a program must have at least one of each of the following as part of the team: pediatric surgeon, pediatric hematologist/oncologist, diagnostic radiologist with expertise in vascular anomalies, interventional radiologist with expertise in vascular anomalies, vascular pathologist, and support from physical or occupational therapy for rehabilitation following vascular surgery.

Virology laboratory with weekday 24-hour availability (F12b). This is a diagnostic laboratory that supports the NICU by conducting culture and tissue studies to determine patients’ virological conditions. Laboratory should be able to complete one or more of the following tests: HSV PCR from CSF, HSV PCR from blood, or direct HSV antigen testing for skin lesions.

3-Tesla magnetic resonance imaging (3T MRI) (A10d). 3T MRI is a higher-powered version of MRI that offers improved morphological and functional studies of the brain compared with the more common field strength of 1.5T.

Appendix C
2020-21 Sample Physician Questionnaire



Best Children's Hospitals

Your nominations will be reflected in the 2020-21
U.S. News & World Report <<specialty>> rankings.

Please name up to 10 U.S. hospitals that in your opinion provide the best care in <<specialty>> for patients who have the most challenging conditions or who need particularly difficult procedures.

Do not consider location or cost. For a hospital that is part of a health system or medical school, please name the individual hospital.

	Hospital	City	State
a.	<input type="text"/>	<input type="text"/>	<input type="text"/>
b.	<input type="text"/>	<input type="text"/>	<input type="text"/>
c.	<input type="text"/>	<input type="text"/>	<input type="text"/>
d.	<input type="text"/>	<input type="text"/>	<input type="text"/>
e.	<input type="text"/>	<input type="text"/>	<input type="text"/>
f.	<input type="text"/>	<input type="text"/>	<input type="text"/>
g.	<input type="text"/>	<input type="text"/>	<input type="text"/>
h.	<input type="text"/>	<input type="text"/>	<input type="text"/>
i.	<input type="text"/>	<input type="text"/>	<input type="text"/>
j.	<input type="text"/>	<input type="text"/>	<input type="text"/>

Thank you for your participation in this important study!

Please respond by returning your survey in the included postpaid envelope,
or by faxing your response to (800) 476-9721.



Conducted by:

RTI International
3040 Cornwallis Rd, PO Box 12194,
Research Triangle Park, NC 27709-2194

<<id>>
PPS01

Appendix D

2020-21 Best Children's Hospitals Rankings by Specialty

**Best Children's Hospital 2020-21:
Diabetes & Endocrinology**

Rank	Hospital	Overall Score	Success in managing diabetes	Success in hypothyroid management	Ability to prevent infections throughout hospital	Number of patients	Number of procedures	Nurse staffing	Advanced clinical services offered	Clinical support services offered	Advanced technologies available	Specialized clinics and programs available	Has fulltime subspecialists available	Recognized as Nurse Magnet hospital	Reputation with physicians in specialty	Commitment to best practices	Commitment to quality improvement	Adoption of health information technology	Active fellowship programs	Commitment to clinical research	Help for families	Enlists families in structuring care
1	Children's Hospital of Philadelphia	100	44	5	36	36	13	4.3	17	9	11	22	14	1	54.8	119	16	14	7	3	8	7
2	Boston Children's Hospital	94.7	39	6	34	35	9	3.9	17	9	11	21	14	1	46.0	103	16	14	7	3	8	7
3	Cincinnati Children's Hospital Medical Center	90.9	38	5	35	35	14	4.1	17	9	11	22	14	1	26.9	118	16	14	7	3	8	7
4	Children's Hospital Colorado	89.2	42	4	34	36	14	4.3	17	9	11	22	14	1	26.0	106	16	14	7	3	8	7
5	Texas Children's Hospital	87.5	40	4	36	36	14	4.6	17	9	11	22	14	1	18.8	114	16	14	7	3	8	7
6	Yale New Haven Children's Hospital	86.2	39	6	32	30	10	5.3	17	8	11	22	13	1	12.4	117	16	14	5	3	8	7
7	UPMC Children's Hospital of Pittsburgh	86.0	39	5	30	36	12	3.5	17	9	11	19	14	1	18.8	114	16	14	7	3	8	7
8	Children's Hospital Los Angeles	83.7	38	4	32	36	10	3.8	16	9	11	22	14	1	21.4	112	16	14	6	3	8	7
9	Lucile Packard Children's Hospital Stanford	81.5	42	3	35	32	13	3.9	17	9	11	18	14	1	14.9	95	16	14	7	3	8	7
10	Children's National Hospital	76.5	34	3	37	35	14	4.1	17	9	10	22	14	1	7.3	112	16	14	7	3	8	7
11	Nationwide Children's Hospital	76.2	31	5	34	36	9	3.3	17	9	11	20	14	1	6.8	108	16	14	7	3	8	7
12	Johns Hopkins Children's Center	75.8	36	4	34	26	11	3.3	16	9	11	18	14	1	8.6	107	16	14	7	2	8	7
13	North Carolina Children's Hospital at UNC	75.3	41	4	34	32	12	4.7	17	9	11	20	13	1	2.8	110	15	13	5	1	8	7
14	Rady Children's Hospital	75.1	44	3	33	34	13	3.2	17	9	11	19	14	1	3.3	101	16	14	6	2	8	7
15	Children's Medical Center Dallas	75.0	36	4	34	34	14	3.2	17	9	11	17	13	1	6.2	102	16	14	7	2	8	7
16	C.S. Mott Children's Hospital-Michigan Medicine	74.8	32	6	35	30	9	3.7	17	9	11	17	11	1	3.1	100	16	14	7	2	8	7
16	CHOC Children's Hospital	74.8	43	4	34	34	12	3.6	17	7	11	18	14	1	1.7	108	16	14	2	2	8	7
18	Mount Sinai Kravis Children's Hospital	74.6	36	6	30	29	14	3.7	17	8	11	20	12	1	2.7	112	15	14	3	1	8	7
18	Norton Children's Hospital	74.6	45	4	35	34	14	3.1	17	9	11	19	14	0	1.5	106	16	14	4	3	8	7
20	UCSF Benioff Children's Hospitals, San Francisco and Oakland	74.5	30	4	32	34	12	4.1	17	9	11	20	14	1	13.9	98	16	14	7	2	8	7
21	Rainbow Babies and Children's Hospital	74.1	31	5	34	31	12	3.3	18	9	11	22	12	1	4.6	111	16	14	4	3	8	7
21	Seattle Children's Hospital	74.1	32	3	35	35	13	3.4	17	9	11	22	14	1	10.7	111	16	14	7	2	8	7
22	UF Health Shands Children's Hospital	73.3	38	4	29	28	5	2.7	17	8	11	22	12	1	6.3	109	16	14	6	3	8	7
23	St. Louis Children's Hospital-Washington University	73.0	32	4	33	34	14	3.8	17	9	11	20	14	1	3.8	101	16	14	7	3	8	7
24	NewYork-Presbyterian Hospital-Columbia and Cornell	72.6	35	4	30	29	13	3.0	16	9	11	20	13	1	8.8	97	16	14	6	2	8	7
26	Riley Hospital for Children at IU Health	72.1	33	3	31	35	11	4.3	17	9	11	21	14	1	9.6	95	16	14	6	3	8	7
27	Cohen Children's Medical Center	71.6	36	4	34	33	13	3.9	16	9	11	15	14	1	2.4	108	16	14	5	1	8	7
28	MassGeneral Hospital for Children	70.7	27	5	31	28	14	3.5	15	9	11	20	11	1	4.9	111	16	14	5	3	8	7
29	Arnold Palmer Hospital for Children	69.9	37	6	32	31	12	3.0	14	8	8	21	12	1	0.2	103	16	14	1	1	7	7
30	Doernbecher Children's Hospital at Oregon Health and Science University	69.3	32	6	27	32	11	3.9	15	9	11	14	11	1	2.9	86	16	14	4	2	8	7
30	Monroe Carell Jr. Children's Hospital at Vanderbilt	69.3	34	3	33	33	9	3.1	17	9	11	22	14	1	5.3	101	16	14	6	2	8	7
32	Ann and Robert H. Lurie Children's Hospital of Chicago	69.2	33	4	31	33	10	3.0	17	9	10	16	14	1	7.3	96	16	14	6	1	8	7
33	Johns Hopkins All Children's Hospital	69.0	32	6	34	32	8	3.8	17	9	11	12	14	1	0.4	99	16	14	2	1	8	7
34	University of Minnesota Masonic Children's Hospital	68.3	40	4	30	31	9	4.0	15	9	11	17	13	0	1.8	102	15	14	6	2	8	7
35	Intermountain Primary Children's Hospital-University of Utah	67.6	39	5	33	32	9	3.4	16	9	11	22	13	0	1.4	96	16	13	3	1	7	7
36	Connecticut Children's Medical Center	67.3	35	4	33	32	11	3.1	17	8	11	21	14	1	1.5	104	16	14	2	1	8	7
36	Duke Children's Hospital and Health Center	67.3	28	5	33	29	5	3.5	17	9	11	20	14	1	1.8	102	16	14	6	2	7	7
38	University of Iowa Stead Family Children's Hospital	66.8	41	3	30	29	7	3.4	16	9	11	22	13	1	0.6	94	14	14	4	2	8	7
39	Phoenix Children's Hospital	66.6	26	6	34	36	14	3.0	17	9	11	22	14	0	0.8	104	16	14	6	2	8	7
40	Mayo Clinic Children's Center	66.5	32	3	30	30	14	4.1	16	9	11	20	12	1	2.6	110	16	14	4	2	8	7
40	University of Virginia Children's Hospital	66.5	31	4	32	26	10	3.1	17	8	11	22	12	1	1.7	106	16	14	3	3	8	7
42	Children's Healthcare of Atlanta	66.3	25	4	34	36	8	4.8	17	9	11	19	14	1	2.6	103	16	14	7	2	8	7
43	Children's Mercy Kansas City	66.2	28	3	34	33	7	4.6	17	9	11	19	14	1	5.2	101	16	14	6	2	8	7
44	Cleveland Clinic Children's Hospital	65.9	29	4	33	34	12	3.7	15	9	11	21	14	1	1.2	110	16	14	4	1	8	7
45	Wolfson Children's Hospital	65.7	33	4	32	32	11	2.6	17	9	10	19	14	1	1.4	98	16	14	4	2	7	7
46	Valley Children's Healthcare and Hospital	65.5	40	3	34	33	11	3.4	16	9	7	11	14	1	1.1	101	16	13	1	1	8	7
47	University of Chicago Comer Children's Hospital	65.3	32	4	30	23	12	2.8	15	9	11	21	11	1	1.6	98	16	14	6	2	8	7
48	Holtz Children's Hospital at UM-Jackson Memorial Medical Center	65.2	42	5	26	27	10	2.2	17	9	9	21	11	0	1.7	114	15	12	4	1	8	3
49	Nicklaus Children's Hospital	64.7	32	6	25	34	8	2.7	17	6	9	19	13	1	1.5	105	14	14	2	1	7	7
50	Spectrum Health Helen DeVos Children's Hospital	64.1	40	4	26	32	10	2.7	15	9	11	13	13	1	0.1	104	14	14	1	1	8	7
50	UCLA Mattel Children's Hospital	64.1	32	3	29	20	8	4.2	15	8	11	15	11	1	6.4	102	15	14	6	1	8	7

Rankings are based on all of the above measures.

NA: Service not provided by hospital. NR: Data not reported or unavailable.

Best Children's Hospital 2020-21: Neurology & Neurosurgery		Overall Score	Ability to prevent infections throughout hospital	Survival after surgery	Ability to prevent surgical complications	Success in controlling epilepsy	Number of surgeries	Number of epilepsy workups and treatments	Nurse staffing	Advanced clinical services offered	Clinical support services offered	Advanced technologies available	Specialized clinics and programs available	Has fulltime subspecialists available	Recognized as Nurse Magnet hospital	Reputation with physicians in specialty	Commitment to best practices	Commitment to quality improvement	Adoption of health information technology	Active fellowship programs	Commitment to clinical research	Help for families	Enlists families in structuring care
Rank	Hospital																						
1	Boston Children's Hospital	100	39	12	22	6	39	12	3.9	21	10	13	19	15	1	48.2	36	15	14	9	7	8	7
2	Texas Children's Hospital	96.8	41	12	22	6	41	12	4.6	21	10	13	19	15	1	26.7	37	15	14	9	6	8	7
3	Children's National Hospital	95.6	41	12	22	6	39	12	4.1	21	10	13	19	15	1	20.9	37	15	14	9	7	8	7
4	Children's Hospital of Philadelphia	94.4	40	12	19	5	35	12	4.3	20	10	13	18	15	1	41.4	37	15	14	9	7	8	7
5	Cincinnati Children's Hospital Medical Center	92.5	40	12	21	5	40	12	4.1	21	10	13	18	15	1	24.0	37	15	14	9	6	8	7
6	Nationwide Children's Hospital	91.3	39	12	21	6	34	12	3.3	21	10	13	19	15	1	15.3	37	15	14	9	7	8	7
7	St. Louis Children's Hospital-Washington University	89.3	39	12	16	6	38	12	3.8	21	10	13	19	15	1	19.9	37	15	14	9	6	8	7
8	Lucile Packard Children's Hospital Stanford	89.2	41	12	21	6	34	7	3.9	17	10	13	19	15	1	10.2	37	15	14	9	7	8	7
9	Children's Hospital Los Angeles	89.0	38	12	22	6	42	12	3.8	21	10	13	19	15	1	8.7	37	15	14	8	5	8	7
10	Children's Hospital Colorado	87.9	40	12	16	6	38	12	4.3	20	10	13	19	15	1	16.9	36	15	14	9	5	8	7
11	Ann and Robert H. Lurie Children's Hospital of Chicago	86.5	36	12	20	6	38	9	3.0	20	10	11	18	15	1	13.8	36	15	14	9	5	8	7
12	Children's Healthcare of Atlanta	85.5	40	12	21	6	34	10	4.8	21	10	13	19	15	1	3.6	36	15	14	9	5	8	7
13	Johns Hopkins Children's Center	85.1	39	12	21	4	32	7	3.3	20	10	13	19	15	1	18.4	36	15	14	9	5	8	7
13	UPMC Children's Hospital of Pittsburgh	85.1	35	11	20	6	27	10	3.5	20	10	12	19	15	1	9.8	37	15	14	9	7	8	7
15	UCSF Benioff Children's Hospitals, San Francisco and Oakland	84.9	37	11	16	6	33	11	4.1	20	10	13	19	15	1	12.0	37	15	14	9	7	8	7
16	Rady Children's Hospital	83.6	39	12	21	6	42	10	3.2	19	10	13	19	15	1	2.3	37	15	14	8	5	8	7
17	Children's Medical Center Dallas	83.3	40	12	22	5	41	12	3.2	20	10	12	19	14	1	3.5	37	15	14	8	5	8	7
17	Intermountain Primary Children's Hospital-University of Utah	83.3	38	12	22	6	39	12	3.4	21	10	13	19	15	0	9.2	34	15	13	7	6	7	7
19	NewYork-Presbyterian Hospital-Columbia and Cornell	82.9	36	12	21	6	31	9	3.0	19	10	13	17	15	1	6.4	36	15	14	8	4	8	7
20	C.S. Mott Children's Hospital-Michigan Medicine	82.5	40	12	20	6	31	7	3.7	19	10	12	15	13	1	4.9	32	15	14	9	7	8	7
20	Cohen Children's Medical Center	82.5	39	12	22	6	34	10	3.9	19	10	13	19	15	1	2.0	37	15	14	5	3	8	7
22	Riley Hospital for Children at IU Health	82.2	36	11	22	6	40	8	4.3	20	10	12	18	14	1	2.7	37	15	14	8	4	8	7
23	Nicklaus Children's Hospital	80.7	31	12	22	6	28	12	2.7	18	7	12	17	14	1	6.8	36	15	14	6	5	7	7
23	Seattle Children's Hospital	82.0	40	11	13	6	32	9	3.4	13	10	12	19	14	1	19.7	34	15	14	9	6	8	7
25	Cleveland Clinic Children's Hospital	79.7	38	12	20	4	26	11	3.7	13	10	13	19	15	1	7.6	35	15	14	9	5	8	7
26	Children's Mercy Kansas City	79.3	40	12	18	6	32	7	4.6	20	10	13	14	14	1	2.1	37	15	14	6	4	8	7
27	Johns Hopkins All Children's Hospital	78.9	40	12	21	6	30	7	3.8	17	10	13	17	15	1	0.9	34	15	14	4	4	8	7
28	Monroe Carell Jr. Children's Hospital at Vanderbilt	78.8	39	12	16	6	37	10	3.1	19	10	12	19	14	1	4.5	36	15	14	5	5	8	7
29	Phoenix Children's Hospital	77.0	40	11	16	6	36	12	3.0	21	10	13	19	15	0	4.3	37	15	14	9	6	8	7
30	Cook Children's Medical Center	76.8	31	12	21	6	29	12	3.4	15	10	13	19	15	1	2.8	35	13	14	0	5	8	7
31	Children's Memorial Hermann Hospital	76.7	36	12	20	6	26	6	3.4	19	10	12	16	15	1	1.4	33	15	14	6	5	7	7
32	Children's Hospital of Alabama at UAB	76.6	39	12	21	4	40	10	3.4	20	10	11	16	15	0	6.3	35	15	14	9	5	8	7
33	CHOC Children's Hospital	76.1	39	12	15	6	25	12	3.6	14	8	12	18	15	1	1.8	37	15	14	6	5	8	7
33	Duke Children's Hospital and Health Center	76.1	38	12	15	6	31	11	3.5	10	10	13	17	13	1	3.5	37	15	14	5	5	7	7
35	Le Bonheur Children's Hospital	75.1	39	11	14	6	25	9	3.0	18	10	13	11	14	1	5.2	35	15	14	8	5	8	7
36	Doernbecher Children's Hospital at Oregon Health and Science University	73.9	33	12	18	5	26	5	3.9	18	10	12	19	13	1	2.0	35	14	14	6	6	8	7
37	Valley Children's Healthcare and Hospital	73.8	38	12	22	6	20	8	3.4	15	10	11	7	14	1	0.9	34	15	13	1	2	8	7
38	Mayo Clinic Children's Center	73.5	36	12	12	5	22	9	4.1	21	10	13	19	13	1	5.9	37	15	14	4	5	8	7
39	Mount Sinai Kravis Children's Hospital	73.0	33	12	20	6	19	12	3.7	9	9	13	19	13	1	0.5	36	14	14	1	2	8	7
40	Children's Hospital of Wisconsin	72.9	36	12	13	6	25	10	4.3	19	10	13	18	15	1	2.3	33	15	12	5	4	8	7
40	UCLA Mattel Children's Hospital	72.9	33	12	17	4	17	7	4.2	18	9	13	4	14	1	6.3	36	14	14	8	7	8	7
42	Spectrum Health Helen DeVos Children's Hospital	72.7	32	12	18	6	23	11	2.7	21	10	13	18	14	1	0.4	36	15	14	1	4	8	7
43	Children's Hospital of Michigan	72.3	35	11	19	6	26	10	3.1	20	10	13	13	15	0	0.6	34	15	13	9	6	8	7
44	Levine Children's Hospital	71.9	39	12	19	5	26	7	3.0	21	10	13	15	14	1	0.9	34	15	14	1	3	8	7
45	Yale New Haven Children's Hospital	71.3	35	12	19	4	19	7	5.3	17	9	13	19	14	1	0.9	36	15	14	3	5	8	7
46	Nemours Alfred I. duPont Hospital for Children	71.0	33	11	19	6	16	9	3.5	18	10	11	14	14	1	0.7	31	15	14	5	3	8	7
47	Children's Hospital at Montefiore	70.7	38	12	22	2	17	10	4.6	15	10	13	19	14	0	3.7	36	15	14	9	5	8	7
48	Akron Children's Hospital	70.2	34	12	15	6	23	5	3.5	17	10	13	18	14	1	1.2	33	15	14	1	5	8	7
49	University of Iowa Stead Family Children's Hospital	69.8	35	12	16	5	25	5	3.4	18	10	13	19	14	1	1.2	33	13	14	4	5	8	7
50	Joseph M. Sanzari Children's Hospital at Hackensack University Medical Center	69.3	37	12	19	6	19	4	2.5	9	8	13	16	13	1	0.8	35	14	13	0	3	8	7

Rankings are based on all of the above measures.

NA: Service not provided by hospital. NR: Data not reported or unavailable.

Best Children's Hospital 2020-21: Orthopedics		Overall Score	Speed and success in treating complex fractures	Ability to prevent surgical complications	Ability to prevent infections throughout hospital	Number of patients	Number of procedures and surgeries	Nurse staffing	Advanced clinical services offered	Clinical support services offered	Advanced technologies available	Specialized clinics and programs available	Has fulltime subspecialists available	Recognized as Nurse Magnet hospital	Reputation with physicians in specialty	Commitment to best practices	Commitment to quality improvement	Adoption of health information technology	Active fellowship programs	Commitment to clinical research	Help for families	Enlists families in structuring care
1	Children's Hospital of Philadelphia	100	8	14	37	24	23	4.3	15	9	3	12	22	1	44.3	81	15	14	9	1	8	7
2	Boston Children's Hospital	97.8	8	13	36	22	26	3.9	14	9	3	12	22	1	49.9	76	15	14	9	1	8	7
3	Cincinnati Children's Hospital Medical Center	91.4	8	13	37	19	24	4.1	16	9	3	12	22	1	20.8	77	15	14	9	1	8	7
4	Children's Hospital Los Angeles	91.2	8	14	35	20	23	3.8	16	9	3	12	22	1	19.8	76	15	14	8	1	8	7
5	Children's Medical Center Dallas-Texas Scottish Rite Hospital for Children	89.8	8	9	37	21	26	3.2	16	9	3	12	21	1	42.3	81	15	14	9	1	8	7
6	St. Louis Children's Hospital-Washington University/Shriners Hospital	89.0	8	14	36	21	25	3.8	15	9	3	12	22	1	10.4	78	15	14	9	1	8	7
7	Rady Children's Hospital	88.2	8	10	36	22	20	3.2	14	9	3	12	22	1	36.5	79	15	14	8	1	8	7
8	Nationwide Children's Hospital	86.2	8	14	36	19	23	3.3	15	9	3	12	22	1	8.1	77	15	14	9	1	8	7
9	Children's National Hospital	86.1	8	15	38	21	15	4.1	14	9	3	12	22	1	5.7	71	15	14	9	1	8	7
10	Children's Healthcare of Atlanta	84.0	7	13	37	23	22	4.8	14	9	3	12	22	1	11.0	75	15	14	9	1	8	7
11	Rainbow Babies and Children's Hospital	82.1	8	14	35	19	19	3.3	14	9	3	12	21	1	6.7	77	15	14	4	1	8	7
12	Johns Hopkins Children's Center	81.3	8	14	36	12	12	3.3	13	9	3	12	22	1	5.7	76	15	14	9	1	8	7
13	Mayo Clinic Children's Center	80.1	8	15	33	8	13	4.1	16	9	3	12	21	1	4.0	74	15	14	6	1	8	7
14	Children's Hospital Colorado	79.7	6	12	37	20	24	4.3	16	9	3	11	22	1	13.1	74	15	14	9	1	8	7
14	Seattle Children's Hospital	79.7	8	12	37	21	18	3.4	5	9	3	12	20	1	10.2	65	15	14	9	1	8	7
15	Texas Children's Hospital	78.9	7	10	38	20	25	4.6	16	9	3	12	22	1	9.2	77	15	14	9	1	8	7
16	Nemours Alfred I. duPont Hospital for Children	78.0	7	10	31	21	22	3.5	16	9	2	12	22	1	18.5	78	15	14	8	1	8	7
18	UPMC Children's Hospital of Pittsburgh-Shriners Hospitals for Children Erie	77.4	8	14	33	20	16	3.5	15	9	3	10	22	1	0.9	68	15	14	9	1	8	7
19	Cohen Children's Medical Center	76.4	7	15	36	17	18	3.9	14	9	3	12	22	1	0.7	76	15	14	3	1	8	7
20	CHOC Children's Hospital	75.3	8	15	36	14	15	3.6	10	7	3	10	22	1	1.1	69	15	14	2	1	8	7
21	Valley Children's Healthcare and Hospital	74.9	8	15	35	18	18	3.4	9	9	2	11	21	1	0.3	73	15	13	1	1	8	7
22	C.S. Mott Children's Hospital-Michigan Medicine	74.5	7	12	37	18	15	3.7	14	9	3	12	21	1	2.9	73	15	14	9	1	8	7
23	Phoenix Children's Hospital	74.0	8	13	37	15	23	3.0	15	9	3	12	22	0	9.7	77	15	14	8	1	8	7
24	Intermountain Primary Children's Hospital-Shriners Hospitals for Children-Univ. of Utah	73.6	7	12	36	21	24	3.4	16	9	3	11	22	0	8.8	70	15	13	7	1	7	7
25	Joe DiMaggio Children's Hospital at Memorial	73.3	8	15	34	19	22	3.5	11	9	3	12	20	0	1.1	76	15	14	0	1	7	7
26	Cleveland Clinic Children's Hospital	73.2	8	14	35	12	10	3.7	7	9	3	9	22	1	1.4	70	15	14	4	1	8	7
27	Lerner Children's Pavilion-Hospital for Special Surgery	72.8	8	8	38	14	23	4.1	12	8	3	12	22	1	4.8	78	15	14	6	1	8	7
28	Children's Mercy Kansas City	72.5	8	8	37	22	21	4.6	15	9	3	11	20	1	4.3	77	15	14	4	1	8	7
29	Levine Children's Hospital	72.2	8	13	36	14	12	3.0	14	9	3	10	22	1	1.5	67	15	14	3	1	8	7
30	Monroe Carell Jr. Children's Hospital at Vanderbilt	71.7	7	10	36	16	18	3.1	13	9	3	12	21	1	6.0	74	15	14	8	1	8	7
31	Lucile Packard Children's Hospital Stanford	71.6	6	13	38	11	17	3.9	15	9	3	10	22	1	3.2	66	15	14	9	1	8	7
32	North Carolina Children's Hospital at UNC	71.2	7	14	37	11	12	4.7	15	9	3	6	22	1	1.1	68	14	13	5	1	8	7
33	UC Davis Children's Hospital/Shriners Hospitals for Children-Northern California	71.0	8	6	37	19	24	6.9	15	9	3	12	22	1	7.4	75	15	14	4	1	8	7
34	Le Bonheur Children's Hospital	70.6	8	9	36	15	16	3.0	14	9	3	11	21	1	4.0	73	15	14	7	1	8	7
35	Ann and Robert H. Lurie Children's Hospital of Chicago	69.9	7	12	34	16	23	3.0	14	9	2	8	22	1	4.6	67	15	14	4	1	8	7
36	Nicklaus Children's Hospital	69.6	8	11	28	22	17	2.7	15	6	2	11	21	1	2.7	78	15	14	6	1	7	7
37	University of Iowa Stead Family Children's Hospital	69.4	6	15	32	19	13	3.4	15	9	3	12	21	1	1.8	65	13	14	4	1	8	7
38	UCLA Mattel Children's Hospital	68.4	8	7	31	19	16	4.2	15	8	3	11	21	1	3.6	81	14	14	8	1	8	7
39	Dayton Children's Hospital	68.0	8	9	37	18	17	4.1	15	9	2	12	21	1	1.2	73	15	14	1	1	8	7
40	Kentucky Children's Hospital-Shriners Hospitals for Children	67.6	6	15	33	10	14	3.1	15	8	3	6	19	1	4.4	70	14	14	0	1	8	7
41	Spectrum Health Helen DeVos Children's Hospital	67.2	8	13	30	10	6	2.7	14	9	3	12	20	1	0.8	70	15	14	1	1	8	7
42	Children's Hospital of Michigan	66.3	7	13	33	20	12	3.1	13	9	3	11	22	0	0.3	74	15	13	7	1	8	7
43	University of Virginia Children's Hospital	65.8	8	10	35	8	9	3.1	15	8	3	10	20	1	0.3	79	15	14	3	1	8	7
44	Doernbecher Children's Hospital at Oregon Health and Science University	64.0	5	13	30	10	20	3.9	15	9	3	11	20	1	3.2	72	15	14	2	1	8	7
44	Riley Hospital for Children at IU Health	64.0	5	13	33	16	15	4.3	15	9	3	12	20	1	0.7	70	15	14	4	1	8	7
45	Children's Hospital of Wisconsin	63.7	7	9	33	15	17	4.3	15	9	3	12	22	1	0.7	71	15	12	4	1	8	7
47	Children's Hospital of Alabama at UAB	63.6	7	11	36	19	16	3.4	15	9	2	10	22	0	0.8	72	15	14	5	1	8	7
48	Wolfson Children's Hospital	63.1	6	14	33	11	11	2.6	14	9	2	5	21	1	1.8	72	15	14	2	1	7	7
49	UCSF Benioff Children's Hospitals, San Francisco and Oakland	62.7	5	10	35	16	15	4.1	15	9	3	10	22	1	3.0	72	15	14	7	1	8	7
50	Johns Hopkins All Children's Hospital	61.9	8	6	37	11	11	3.8	14	9	3	12	22	1	2.0	75	15	14	0	1	8	7
50	University of Minnesota Masonic Children's Hospital	61.9	8	12	32	8	9	4.0	5	9	3	10	22	0	0.2	63	14	14	6	1	8	7

Rankings are based on all of the above measures.

NA: Service not provided by hospital. NR: Data not reported or unavailable.

Best Children's Hospital 2020-21: Pulmonology & Lung Surgery		Overall Score	Success with asthma inpatients	Success in helping patients manage their asthma	Success in managing cystic fibrosis patients weakness disorder	Survival of patients on ventilators	Ability to prevent infections throughout hospital	Ability to prevent infections in intensive-care units	Ability to prevent pressure injuries	Survival after lung transplant	Number of patients	Number of tests and noninvasive procedures	Nurse staffing	Lung transplant program	Advanced clinical services offered	Clinical support services offered	Advanced technologies available	Has fulltime subspecialists available	Recognized as Nurse Magnet hospital	Reputation with physicians in specialty	Commitment to best practices	Commitment to quality improvement	Adoption of health information technology	Active fellowship programs	Commitment to clinical research	Help for families	Enlists families in structuring care	
Rank	Hospital																											
1	Boston Children's Hospital	100	8	12	15	3	3	48	5	5	2	17	12	3.9	5	38	9	2	12	1	46.4	47	15	14	6	3	8	7
2	Children's Hospital of Philadelphia	98.5	7	12	17	3	3	49	3	5	5	17	12	4.3	5	38	9	2	12	1	52.3	48	15	14	4	2	8	7
3	Texas Children's Hospital	97.8	8	13	14	3	3	49	3	5	5	17	12	4.6	5	38	9	2	12	1	37.7	48	15	14	6	3	8	7
4	Cincinnati Children's Hospital Medical Center	95.7	7	13	16	3	3	48	3	5	2	17	12	4.1	5	37	9	2	12	1	46.7	48	15	14	6	2	8	7
5	Children's Hospital Colorado	93.5	6	13	15	3	3	48	4	5	NA	15	12	4.3	NA	38	9	2	12	1	45.1	48	15	14	6	3	8	7
6	Nationwide Children's Hospital	92.1	7	13	15	3	3	48	5	5	3	17	12	3.3	4	37	9	2	12	1	20.0	48	15	14	6	2	8	7
7	Lucile Packard Children's Hospital Stanford	91.3	8	12	13	3	3	49	5	4	4	16	12	3.9	5	37	9	2	12	1	12.6	47	15	14	6	3	8	7
8	St. Louis Children's Hospital- Washington University	90.4	7	11	17	3	3	43	5	3	3	12	10	3.8	5	38	9	2	12	1	15.5	48	15	14	6	3	8	7
9	Children's National Hospital	90.1	8	13	17	3	3	50	5	5	NA	14	12	4.1	NA	38	9	2	12	1	8.6	48	15	14	6	3	8	7
10	Seattle Children's Hospital	88.4	8	11	14	3	3	48	5	4	NA	13	10	3.4	NA	30	9	2	11	1	25.5	46	15	14	6	2	8	7
10	UPMC Children's Hospital of Pittsburgh	87.4	6	8	15	3	3	44	5	2	5	15	10	3.5	5	33	9	2	12	1	17.7	48	15	14	6	2	8	7
12	Children's Hospital Los Angeles	85.4	8	13	13	3	3	47	5	4	NA	14	11	3.8	NA	37	9	2	12	1	13.3	48	15	14	5	2	8	7
13	Rainbow Babies and Children's Hospital	84.2	8	11	16	3	3	47	5	4	NA	12	10	3.3	NA	37	9	2	11	1	8.8	46	15	14	4	3	8	7
14	Riley Hospital for Children at IU Health	83.7	6	12	14	3	2	42	5	5	2	15	10	4.3	2	37	9	2	11	1	15.9	48	15	14	5	3	8	7
15	Children's Hospital of Wisconsin	81.1	7	13	18	3	3	44	5	5	NA	14	11	4.3	NA	37	9	2	12	1	1.6	47	15	12	5	2	8	7
16	Monroe Carell Jr. Children's Hospital at Vanderbilt	80.4	7	12	17	3	3	46	2	5	NR	13	12	3.1	1	38	9	2	11	1	10.2	47	15	14	5	2	8	7
17	Children's Healthcare of Atlanta	80.2	7	13	14	3	3	45	5	5	NR	15	12	4.8	1	37	9	2	12	1	2.5	47	15	14	6	2	8	7
18	C.S. Mott Children's Hospital-Michigan Medicine	79.5	7	11	15	3	3	47	5	3	NA	13	11	3.7	NA	38	9	2	11	1	3.1	43	15	14	6	3	8	7
18	Rady Children's Hospital	79.5	7	13	15	3	3	46	5	3	NA	12	8	3.2	NA	36	9	2	12	1	3.3	48	15	14	5	3	8	7
20	Johns Hopkins Children's Center	79.1	5	13	14	3	3	48	2	5	NR	15	10	3.3	1	37	9	2	12	1	16.5	47	15	14	6	2	8	7
21	Ann and Robert H. Lurie Children's Hospital of Chicago	78.4	6	12	15	3	3	43	3	4	NA	11	11	3.0	NA	38	9	2	12	1	9.2	47	15	14	6	3	8	7
22	North Carolina Children's Hospital at UNC	77.9	7	11	14	3	2	49	1	4	3	11	7	4.7	3	33	9	2	11	1	15.9	47	14	13	5	2	8	7
23	MassGeneral Hospital for Children	77.6	7	12	15	3	3	46	5	4	NA	8	11	3.5	NA	37	9	2	9	1	2.7	47	15	14	5	2	8	7
24	Cohen Children's Medical Center	76.7	8	13	15	3	3	48	4	5	NA	8	7	3.9	NA	35	9	2	12	1	0.6	47	15	14	4	2	8	7
25	Children's Medical Center Dallas	76.4	7	12	15	3	2	47	5	5	NA	14	10	3.2	NA	35	9	2	11	1	2.5	45	15	14	6	2	8	7
26	CHOC Children's Hospital	75.6	7	13	14	3	3	48	5	5	NA	12	11	3.6	NA	31	7	2	12	1	1.3	47	15	14	2	2	8	7
27	UF Health Shands Children's Hospital	75.1	8	13	11	3	3	43	5	5	4	12	7	2.7	5	34	8	2	9	1	0.6	43	15	14	5	2	8	7
28	NewYork-Presbyterian Hospital-Columbia and Cornell	74.9	7	7	11	3	3	40	3	2	5	14	10	3.0	4	37	9	2	12	1	5.5	48	15	14	5	2	8	7
29	Children's Hospital of Alabama at UAB	74.7	8	13	16	3	3	44	2	4	NA	14	11	3.4	NA	34	9	2	12	0	4.9	47	15	14	6	2	8	7
29	Children's Hospital of Richmond at VCU	74.7	8	13	17	3	2	47	5	3	NA	10	6	2.4	NA	37	8	2	11	1	1.7	47	15	14	4	2	7	7
31	Valley Children's Healthcare and Hospital	74.3	8	13	15	3	3	47	5	5	NA	9	9	3.4	NA	32	9	1	11	1	0.1	44	15	13	3	2	8	7
32	Cleveland Clinic Children's Hospital	73.6	4	13	18	3	3	43	4	4	NA	11	11	3.7	NA	29	9	2	12	1	1.8	48	15	14	4	2	8	7
32	Duke Children's Hospital and Health Center	73.6	7	11	14	3	3	46	5	3	NR	10	7	3.5	1	29	9	2	11	1	1.5	48	15	14	3	2	7	7
34	Arkansas Children's Hospital	73.4	8	13	15	3	3	34	5	4	NA	9	7	3.2	NA	35	9	2	12	1	1.2	44	14	14	5	2	8	7
35	Children's Mercy Kansas City	73.1	7	12	15	3	3	46	1	5	NA	10	9	4.6	NA	33	9	2	11	1	3.3	45	15	14	6	2	8	7
35	Nicklaus Children's Hospital	73.1	8	13	17	3	3	40	5	4	NA	11	5	2.7	NA	36	6	2	11	1	0.5	46	15	14	2	2	7	7
37	Phoenix Children's Hospital	72.7	8	13	14	3	3	44	5	2	NA	14	10	3.0	NA	35	9	2	12	0	1.3	48	15	14	6	1	8	7
38	Penn State Children's Hospital	72.6	7	12	15	2	3	40	5	4	NA	10	8	3.5	NA	33	9	2	11	1	0.4	46	14	14	5	2	8	7
39	Levine Children's Hospital	72.2	7	13	14	3	3	47	4	5	NA	11	7	3.0	NA	37	9	2	12	1	0.9	47	15	14	1	2	8	7
40	UCSF Benioff Children's Hospitals, San Francisco and Oakland	72.1	6	8	14	3	3	46	2	5	NA	8	8	4.1	NA	37	9	2	11	1	3.7	47	15	14	6	2	8	7
41	Dayton Children's Hospital	71.3	8	12	12	3	3	48	4	4	NA	10	11	4.1	NA	37	9	1	11	1	1.0	44	15	14	1	2	8	7
42	NYU Winthrop Hospital Children's Medical Center	71.2	8	13	11	3	3	45	5	5	NA	7	6	4.7	NA	28	8	2	8	1	1.0	46	14	14	2	2	8	7
43	Yale New Haven Children's Hospital	71.0	6	8	15	1	3	45	4	5	NA	8	7	5.3	NA	36	8	2	11	1	1.8	46	15	14	5	2	8	7
44	Children's Minnesota	70.8	7	13	16	3	3	38	3	3	NA	13	11	3.6	NA	38	9	2	12	1	1.7	40	15	14	3	2	8	7
45	Spectrum Health Helen DeVos Children's Hospital	70.5	6	10	17	3	3	35	5	4	NA	11	9	2.7	NA	37	9	2	11	1	1.2	47	15	14	1	2	8	7
46	Le Bonheur Children's Hospital	70.4	8	13	13	3	3	46	2	4	NA	10	5	3.0	NA	32	9	2	11	1	2.8	42	15	14	5	2	8	7
47	Johns Hopkins All Children's Hospital	70.3	3	13	16	3	3	48	5	5	NA	8	8	3.8	NA	25	9	2	12	1	1.0	46	15	14	2	2	8	7
48	SSM Health Cardinal Glennon Children's Hospital-St. Louis University	69.9	8	11	14	3	2	42	5	4	NA	11	8	3.0	NA	32	9	2	12	1	1.4	45	14	14	1	2	8	7
49	Miller Children's and Women's Hospital Long Beach	69.8	6	11	17	3	3	38	4	5	NA	12	8	4.2	NA	34	8	2	10	1	0.6	43	13	13	3	2	8	6
50	Children's Hospital and Medical Center	69.4	5	13	14	3	3	48	2	5	NA	11	11	4.5	NA	33	8	2	12	1	0.7	46	15	14	4	2	8	7

Rankings are based on all of the above measures.

NA: Service not provided by hospital. NR: Data not reported or unavailable.

Best Children's Hospital 2020-21: Urology		Overall Score	Ability to prevent surgical complications	Speed in treating testicular torsion	Ability to prevent infections throughout hospital	Number of patients	Number of surgeries	Number of minimally invasive procedures	Nurse staffing	Advanced clinical services offered	Clinical support services offered	Advanced technologies available	Specialized clinics and programs available	Has fulltime subspecialists available	Recognized as Nurse Magnet hospital	Reputation with physicians in specialty	Commitment to best practices	Commitment to quality improvement	Adoption of health information technology	Active fellowship programs	Commitment to clinical research	Help for families	Enlists families in structuring care
Rank	Hospital																						
1	Boston Children's Hospital	100	15	2	30	26	18	9	3.9	3	9	8	10	13	1	64.8	41	15	14	6	3	8	7
2	Riley Hospital for Children at IU Health	96.5	16	2	27	26	18	9	4.3	3	9	8	10	13	1	44.8	42	15	14	5	3	8	7
3	Children's Hospital of Philadelphia	93.7	12	2	31	26	19	9	4.3	3	9	8	10	13	1	66.4	42	15	14	6	3	8	7
4	Cincinnati Children's Hospital Medical Center	92.7	14	2	31	23	14	9	4.1	3	9	8	10	13	1	42.4	42	15	14	6	3	8	7
5	Monroe Carell Jr. Children's Hospital at Vanderbilt	86.6	14	2	30	21	17	8	3.1	3	9	8	10	13	1	33.3	41	15	14	5	3	8	7
6	Texas Children's Hospital	84.8	11	2	32	27	19	9	4.6	3	9	8	10	13	1	34.7	42	15	14	6	3	8	7
7	Ann and Robert H. Lurie Children's Hospital of Chicago	84.5	12	2	28	25	16	9	3.0	3	9	7	10	13	1	44.4	42	15	14	6	3	8	7
8	Children's Hospital Colorado	83.9	15	2	31	26	17	9	4.3	3	9	8	10	13	1	10.5	42	15	14	6	3	8	7
9	Seattle Children's Hospital	81.3	12	2	31	20	15	8	3.4	3	9	8	10	13	1	30.5	40	15	14	6	3	8	7
10	Nationwide Children's Hospital	78.0	11	2	30	23	18	9	3.3	3	9	8	10	13	1	26.6	40	15	14	6	3	8	7
11	UPMC Children's Hospital of Pittsburgh	76.6	14	2	27	21	15	9	3.5	3	9	8	10	13	1	10.0	42	15	14	6	3	8	7
12	Rady Children's Hospital	76.0	15	2	30	20	14	8	3.2	3	9	8	10	13	1	8.5	42	15	14	5	2	8	7
13	Children's Hospital Los Angeles	74.4	12	2	29	25	18	9	3.8	3	9	8	10	13	1	14.4	42	15	14	3	3	8	7
14	Children's Medical Center Dallas	74.1	12	2	31	19	15	9	3.2	3	9	8	10	13	1	15.0	42	15	14	6	2	8	7
15	St. Louis Children's Hospital-Washington University	73.9	15	2	30	19	14	8	3.8	3	9	8	10	13	1	3.2	39	15	14	6	3	8	7
16	UCSF Benioff Children's Hospitals, San Francisco and Oakland	72.2	13	2	29	24	14	9	4.1	3	9	8	10	13	1	8.1	40	15	14	6	2	8	7
17	Cohen Children's Medical Center	70.6	14	2	30	18	13	8	3.9	3	9	8	10	13	1	3.4	41	15	14	4	3	8	7
18	Mayo Clinic Children's Center	68.5	13	2	27	23	11	5	4.1	3	9	8	10	12	1	10.2	40	15	14	2	3	8	7
19	Children's Healthcare of Atlanta	68.4	10	2	31	23	18	9	4.8	3	9	8	10	13	1	9.4	39	15	14	6	3	8	7
19	Lucile Packard Children's Hospital Stanford	68.4	12	2	32	20	9	6	3.9	3	9	8	10	13	1	4.6	42	15	14	6	3	8	7
21	Children's Mercy Kansas City	68.1	13	2	31	18	16	8	4.6	3	9	8	10	13	1	1.8	42	15	14	4	3	8	7
22	C.S. Mott Children's Hospital-Michigan Medicine	68.0	13	2	31	19	10	7	3.7	3	9	8	10	12	1	4.7	37	15	14	6	3	8	7
23	Cleveland Clinic Children's Hospital	67.5	14	2	29	13	10	7	3.7	3	9	8	10	13	1	1.1	42	15	14	6	3	8	7
24	Le Bonheur Children's Hospital	67.2	15	2	30	11	12	8	3.0	3	9	7	8	12	1	3.7	39	15	14	5	2	8	7
25	Children's National Hospital	66.7	8	2	32	23	13	9	4.1	3	9	8	10	13	1	13.5	42	15	14	6	3	8	7
26	Johns Hopkins Children's Center	64.8	9	2	30	18	13	6	3.3	3	9	8	10	13	1	15.8	39	15	14	6	3	8	7
27	Children's Hospital and Medical Center	62.0	14	2	30	11	11	8	4.5	3	8	8	9	13	1	1.6	38	15	14	2	2	8	7
28	Duke Children's Hospital and Health Center	61.9	11	2	30	21	13	6	3.5	3	9	8	10	13	1	6.2	39	15	14	3	3	7	7
29	Children's Hospital of Richmond at VCU	60.8	15	2	30	14	14	8	2.4	3	8	8	9	12	1	1.7	37	13	14	2	2	7	7
30	CHOC Children's Hospital	60.5	11	2	30	20	15	7	3.6	3	7	7	10	13	1	3.3	40	15	14	4	3	8	7
31	Rainbow Babies and Children's Hospital	58.9	14	2	29	11	5	4	3.3	3	9	8	10	11	1	1.6	41	15	14	2	2	8	7
32	Children's Hospital of Michigan	57.4	14	2	27	12	11	7	3.1	3	9	7	10	13	0	2.3	39	15	13	4	3	8	7
33	Intermountain Primary Children's Hospital-University of Utah	57.0	11	2	30	26	16	8	3.4	3	9	8	10	13	0	3.9	38	15	13	5	3	7	7
34	Valley Children's Healthcare and Hospital	56.1	14	2	29	15	14	8	3.4	3	9	6	9	13	1	1.5	38	15	13	1	0	8	7
35	North Carolina Children's Hospital at UNC	55.8	11	2	31	9	12	6	4.7	3	9	8	10	12	1	1.4	38	14	13	5	2	8	7
36	UC Davis Children's Hospital/Shriners Hospitals for Children-Northern California	55.1	10	2	31	17	10	8	6.9	3	9	8	10	13	1	1.2	42	15	14	0	3	8	7
37	West Virginia University Children's Hospital	55.0	13	2	30	10	9	6	4.6	3	8	8	10	10	1	0.5	37	12	14	0	3	8	7
38	SSM Health Cardinal Glennon Children's Hospital-St. Louis University	54.8	13	2	28	11	8	5	3.0	3	9	7	10	13	1	1.0	38	14	14	1	3	8	7
39	Connecticut Children's Medical Center	53.8	13	2	30	10	8	4	3.1	3	8	8	10	13	1	0.2	41	15	14	0	2	8	7
40	Akron Children's Hospital	53.7	13	2	25	13	10	5	3.5	3	9	8	7	13	1	2.1	38	15	14	1	2	8	7
40	Norton Children's Hospital	53.7	13	2	31	18	12	4	3.1	3	9	8	10	13	0	0.4	39	15	14	4	2	8	7
40	Spectrum Health Helen DeVos Children's Hospital	53.7	14	2	24	13	12	8	2.7	3	9	7	8	12	1	0.2	39	15	14	1	2	8	7
43	University of Virginia Children's Hospital	52.7	12	2	29	8	12	7	3.1	3	8	8	7	11	1	2.5	38	15	14	1	2	8	7
44	Children's Hospital of Alabama at UAB	52.6	11	2	30	17	15	8	3.4	3	9	6	10	12	0	2.1	38	15	14	4	3	8	7
45	Phoenix Children's Hospital	52.5	10	2	31	22	12	8	3.0	3	9	8	10	13	0	2.5	41	15	14	6	2	8	7
46	Johns Hopkins All Children's Hospital	51.4	15	2	31	8	11	5	3.8	2	9	6	2	13	1	1.0	36	13	14	0	0	8	7
47	MassGeneral Hospital for Children	51.1	10	2	28	14	6	4	3.5	3	9	8	10	12	1	1.5	40	15	14	3	3	8	7
48	Arkansas Children's Hospital	51.0	11	2	21	15	13	7	3.2	3	9	8	10	13	1	0.7	41	14	14	3	3	8	7
49	Arnold Palmer Hospital for Children	50.8	12	2	30	12	8	4	3.0	3	8	7	10	11	1	0.0	38	15	14	1	3	7	7
50	Wolfson Children's Hospital	49.6	12	2	27	15	9	5	2.6	3	9	6	9	13	1	1.7	37	15	14	4	1	7	7

Rankings are based on all of the above measures.

NA: Service not provided by hospital. NR: Data not reported or unavailable.

Appendix E

2020-21 Best Children's Hospitals Honor Roll

2020-21 Best Children's Hospitals Honor Roll

Rank	Name	Points
1	Boston Children's Hospital	238
2	Children's Hospital of Philadelphia	227
3	Cincinnati Children's Hospital Medical Center	209
4	Texas Children's Hospital, Houston	208
5	Children's Hospital Los Angeles	183
6	Children's Hospital Colorado, Aurora	166
7	Children's National Hospital, Washington, D.C.	152
8	Nationwide Children's Hospital, Columbus, Ohio	150
9	UPMC Children's Hospital of Pittsburgh	146
10	Lucile Packard Children's Hospital Stanford, Palo Alto, Calif.	140

